

KUOPION YLIOPISTON JULKAISUJA D. LÄÄKETIEDE 417  
KUOPIO UNIVERSITY PUBLICATIONS D. MEDICAL SCIENCES 417

JAANA HARTIKAINEN

# Genetic Predisposition to Breast and Ovarian Cancer in Eastern Finnish Population

Doctoral dissertation

To be presented by permission of the Faculty of Medicine of the University of Kuopio  
for public examination in Auditorium L3, Canthia building, University of Kuopio,  
on Friday 12<sup>th</sup> October 2007, at 12 noon

Institute of Clinical Medicine,  
Pathology and Forensic Medicine, and Oncology,  
University of Kuopio  
Departments of Oncology and Pathology,  
Kuopio University Hospital



KUOPION YLIOPISTO

KUOPIO 2007

- Distributor:** Kuopio University Library  
P.O. Box 1627  
FI-70211 KUOPIO  
FINLAND  
Tel. +358 17 163 430  
Fax +358 17 163 410  
[www.uku.fi/kirjasto/julkaisutoiminta/julkmyyn.html](http://www.uku.fi/kirjasto/julkaisutoiminta/julkmyyn.html)
- Series Editors:** Professor Esko Alhava, M.D., Ph.D.  
Institute of Clinical Medicine, Department of Surgery
- Professor Raimo Sulkava, M.D., Ph.D.  
School of Public Health and Clinical Nutrition
- Professor Markku Tammi, M.D., Ph.D.  
Institute of Biomedicine, Department of Anatomy
- Author's address:** Institute of Clinical Medicine, Pathology and Forensic Medicine  
University of Kuopio  
P.O. Box 1627  
FI-70211 KUOPIO  
FINLAND  
Tel. +358 17 162 754  
Fax +358 17 162 753  
E-mail: [jaana.hartikainen@uku.fi](mailto:jaana.hartikainen@uku.fi)
- Supervisors:** Docent Arto Mannermaa, Ph.D.  
Institute of Clinical Medicine, Pathology and Forensic Medicine  
University of Kuopio and  
Department of Pathology, Kuopio University Hospital
- Professor Veli-Matti Kosma, M.D., Ph.D.  
Institute of Clinical Medicine, Pathology and Forensic Medicine  
University of Kuopio and  
Department of Pathology, Kuopio University Hospital
- Docent Vesa Kataja, M.D., Ph.D.  
Department of Oncology, Kuopio University Hospital and  
Department of Oncology, Vaasa Central Hospital
- Reviewers:** Docent Minna Pöyhönen, M.D., Ph.D.  
Department of Medical Genetics  
Biomedicum Helsinki  
University of Helsinki
- Docent Johanna Schleutker, Ph.D.  
Institute of Medical Technology  
University of Tampere and Tampere University Hospital
- Opponent:** Docent Kristiina Aittomäki, M.D., Ph.D.  
Department of Clinical Genetics  
Helsinki University Central Hospital

ISBN 978-951-27-0677-8

ISBN 978-951-27-0754-6 (PDF)

ISSN 1235-0303

Kopijyvä  
Kuopio 2007  
Finland

Hartikainen, Jaana. Genetic predisposition to breast and ovarian cancer in Eastern Finnish population. Kuopio University Publications D. Medical Sciences 417. 2007. 188 p.  
ISBN 978-951-27-0677-8  
ISBN 978-951-27-0754-6 (PDF)  
ISSN 1235-0303

## ABSTRACT

Twenty-five percent of the genetic susceptibility to breast cancer is explained by the susceptibility genes known so far. The aims of this study were to investigate the genetic background of familial breast/ovarian cancer and sporadic breast cancer in Eastern Finnish population.

The frequency and type of germ-line mutations in known high-risk breast/ovarian cancer susceptibility genes, *BRCA1* and *BRCA2*, were evaluated in 36 Eastern Finnish breast/ovarian cancer families. In addition, the autosomes were screened for linkage disequilibrium-based association using 435 microsatellite markers to identify new chromosomal regions and genes on them as genetic risk factors for sporadic breast cancer in a case-control set from the Eastern Finnish population.

Germ-line mutations in *BRCA1* and *BRCA2* genes were observed in 19.4 % of the studied Eastern Finnish breast/ovarian cancer families. The observed Eastern Finnish mutation spectrum differs from those observed in the Northern and Southern parts of the country as only two of the eight most common Finnish founder mutations were detected in the Eastern Finnish families. In addition, a novel *BRCA2* 4088insA mutation was found in one family. It has not been found elsewhere in Finland and it appears to be associated with a favourable clinical outcome of breast and ovarian cancer patients but is also highly penetrant.

In the two-staged autosome-wide scan altogether 27 microsatellite markers in 16 chromosomes showed association with breast cancer. In the haplotype analysis three chromosomal regions, 3p26, 11q23 and 22q12-q13 were further suggested as candidate locations for breast cancer associated genes. Breast cancer associated risk factors potentially locate also in the vicinity of single associated markers.

The associated region on chromosome 22q12-q13 was further studied using 10 SNP markers. Significant association was detected with one SNP located in the intronic sequence of *TMPRSS6* gene encoding matriptase-2. The heterozygous genotype and the minor allele were associated with increased risk of breast cancer. Matriptase-2 is a serine protease with functions in processes occurring in both normal and pathological conditions, including cancer progression. Thus, *TMPRSS6* is a potential novel candidate for breast cancer risk affecting gene.

In the AI analysis of 22q12-q13 abundant AI was detected more centromeric than previously reported, further supporting the existence of a tumour suppressor gene or genes in this region.

This study revealed that the proportion of mutations in high-penetrance genes predisposing to familial breast/ovarian cancer in Eastern Finland is similar to other parts of the country, but the mutation spectrum is different. This study also shows that several low-penetrance risk factors for sporadic breast cancer, some of which may be population specific, exist in the Eastern Finnish population.

National Library of Medicine Classification: QU 477, QZ 50, WP 322, WP 870

Medical Subject Headings: Breast Neoplasms/epidemiology; Breast Neoplasms/genetics; Chromosomes, Human, Pair 11/genetics; Chromosomes, Human, Pair 22/genetics; Chromosomes, Human, Pair 3/genetics; Female; Finland/epidemiology; Genes, *BRCA1*; Genes, *BRCA2*; Genetic Markers; Gene Frequency; Genetic Predisposition to Disease; Genotype; Haplotypes; Microsatellite Repeats/genetics; Ovarian Neoplasms/epidemiology; Ovarian Neoplasms/genetics; Genetics, Population; Risk Factors



## ACKNOWLEDGEMENTS

This study was carried out in the Institute of Clinical Medicine, Pathology and Forensic Medicine, and Oncology, University of Kuopio and in the Departments of Oncology and Pathology, Kuopio University Hospital during the years 1997-2007.

I owe my greatest gratitude to my main supervisor Docent Arto Mannermaa, Ph.D., for introducing me to the research of breast cancer genetics. His enthusiasm, guidance, encouragement and support, as well as expertise in cancer genetics, have been invaluable for the completion of this work.

I express my sincere thanks to my other supervisors, Professor Veli-Matti Kosma, M.D., Ph.D., Head of Pathology and Forensic Medicine, Institute of Clinical Medicine, for support and providing me the opportunity as well as the facilities to complete my work, and Docent Vesa Kataja, M.D., Ph.D., for his support and sharing his experience in clinical oncology.

Warm thanks belong to my co-authors, Hanna Tuhkanen, Ph.D., for support and friendship; the discussions and all the fun we have shared has been the resource for managing many bad days, and Airi Arffman, M.Sc., for support, friendship and encouragement; the occasional but persistent enquiries about the shoes was one of the driving forces. I also thank my co-author Mia Pirskanen, M.Sc., for support and friendship.

I wish to sincerely thank my other co-authors Professor Seppo Heinonen, M.D., Ph.D., Professor Matti Eskelinen, M.D., Ph.D., Professor Matti Uusitupa, M.D., Ph.D., Ulla Ristonmaa, M.Sc., Pia Vahteristo, Ph.D., and Professor Markku Ryyänen, M.D., Ph.D., for their collaboration and contribution to this work.

Special thanks belong to my co-authors and collaborators in the Department of Oncology, Strangeways Research Laboratory, University of Cambridge, Alison Dunning, Ph.D., for friendliness and teaching the secrets of linkage disequilibrium, Professor Douglas Easton, Ph.D., for expert knowledge in research of genetic predisposition to cancer, and Antonis Antoniou, Ph.D., and Paula Smith, M.Sc., for friendliness and expert help in statistical analyses.

I wish to sincerely thank the official reviewers Docent Minna Pöyhönen, M.D., Ph.D., and Docent Johanna Schleutker, Ph.D., for their time and valuable, developing and accurate comments during the final preparation of this thesis.

I owe my sincere gratitude to the personnel of Pathology and Forensic Medicine, Institute of Clinical Medicine, University of Kuopio, Ms. Eija Myöhänen, Ms. Irma Väänänen, Ms. Helena Kemiläinen, Ms. Tiina Tirkkonen, Ms. Aija Parkkinen, Ms. Kirsi Alisalo, Ms. Rauni Manninen, Ms. Anne Koivisto, Ms. Merja Fali, and Professor Ylermi Soini, M.D., Ph.D., for help, friendly support and daily discussions. Ms. Eija Myöhänen is especially acknowledged for technical assistance during the last part of my

work. I also thank the fellow researchers Kirsi Suhonen, M.D., Ph.D., Johanna Pellikainen, M.D., Ph.D., Sari Sillanpää, M.D., Maarit Anttila, M.D., Ph.D., Reijo Sironen, M.D., Ph.D., Timo Nykopp, M.B., Ivana Kholová, M.D., Ph.D., Tero Leinonen, B.Sc., Essi Hiltunen, M.D., and Jan Böhm, M.D., Ph.D., for friendly support.

I wish to express warm thanks to the permanent personnel and the visiting scientists of the Strangeways Research Laboratory, University of Cambridge, for friendliness, support, and nice moments spent punting on the river.

I warmly thank the personnel of the Department of Clinical Genetics, Kuopio University Hospital, especially Ms. Inkeri Happonen, Ms. Jaana Hoffren and Ms. Pirkko Jokela. The "old times" and the beginning of my research work in the DNA laboratory are among warm memories.

I wish to express sincere thanks to Ms. Marjo Laitinen, Ms. Petra Mäkinen, Docent Mikko Hiltunen, Ph.D., and Docent Seppo Helisalmi, Ph.D., and his family, for friendship and support during the years.

I also wish to thank my all dear friends for their support, encouragement, relaxing companionship and all the joyful moments that we have shared.

I owe my deepest gratitude to my parents Liisa and Taneli Hartikainen, for endless caring, love and support. I also wish to express special thanks to my little sister Anu and my brother-in-law Jani, for support and all the experiences we have shared. Thank you for being there.

My most loving thanks belong to my fiancé Marko Kauppinen for his love, care and support. You have brought so much joy and wonderful things into my life.

Financial support provided by the Special Government Funding (EVO) of Kuopio University Hospital, the Cancer Fund of Northern Savo, the Paavo Koistinen Fund, the Finnish Cultural Foundation, the University Foundation of Kuopio, the University of Kuopio, the Emil Aaltonen Foundation, and the Finnish Cancer Foundation are gratefully acknowledged.

Kuopio, October 2007

Jaana Hartikainen

## ABBREVIATIONS

|   |   |
|---|---|
| 3 $\beta$ -HSD                                | 3-betahydroxysteroid dehydrogenase  |
| 17 $\beta$ -HSD                               | 17-betahydroxysteroid dehydrogenase   |
| ADH   | alcohol dehydrogenase   |
| ADH1B , <i>ADH1B</i>                          | alcohol dehydrogenase type 2, alcohol dehydrogenase type 2 gene                     |
| ADH1C, <i>ADH1C</i>                           | alcohol dehydrogenase type 3, alcohol dehydrogenase type 3 gene                     |
| <i>ADPRT</i>                                  | poly(ADP-ribose) polymerase 1 gene  |
| AI  | allelic imbalance   |
| AIB1/SRC-3                                    | amplified in breast cancer 1/steroid receptor coactivator 3 protein                 |
| <i>AIB1</i> = <i>NCOA33</i>                   | amplified in breast cancer 1 gene   |
| Alu   | family of short interspersed nuclear elements                                       |
| Apal  | restriction fragment length polymorphism in <i>VDR</i> gene                         |
| APC   | adenomatous polyposis coli  |
| <i>APEX1</i> = <i>APE1</i> , APE1             | apurinic endonuclease gene, apurinic endonuclease                                   |
| AR, AR  | androgen receptor, androgen receptor gene   |
| A-T   | Ataxia-telangiectasia   |
| <i>ATM</i> , ATM                              | ataxia-telangiectasia mutated, gene and protein                                     |
| ATR   | ataxia-telangiectasia and rad3-related  |
| BAP1  | BRCA-interacting protein 1  |
| BARD1, <i>BARD1</i>                           | BRCA1-associated ring domain 1 protein, BRCA1-associated ring domain 1 protein gene |
| BCCIP $\alpha/\beta$                          | BRCA2-interacting protein alpha and beta  |
| BER   | base-excision repair  |
| BIC   | Breast Cancer Information Core  |
| BIR   | break-induced replication   |
| BLM   | DNA helicase recq-like type 2   |
| bp  | base pairs  |
| BRAF35  | BRCA2-associated factor, 35-kD  |
| BRC   | internal repeats in <i>BRCA2</i> gene   |
| <i>BRCA1</i> , BRCA1                          | breast cancer 1, gene and protein   |
| <i>BRCA2</i> , BRCA2                          | breast cancer 2, gene and protein   |
| BRCT  | BRCA1 carboxy-terminal  |
| <i>BRIP1</i> = <i>BACH1</i>                   | BRCA1-interacting protein 1 gene  |
| BsmI  | restriction fragment length polymorphism in <i>VDR</i> gene                         |
| BTF2-TFIIH                                    | basal transcription factor complex  |
| CA125   | cancer antigen 125  |
| <i>CACNG2</i>                                 | voltage-dependent calcium channel gamma-2 subunit gene                              |
| <i>CASP8</i>                                  | caspase 8 gene  |
| <i>CCND1</i>                                  | cyclin D1 gene  |
| Cdc25A  | cell division cycle 25A tyrosine phosphatase  |
| cdk4, cdk6                                    | cyclin dependent kinase 4, cyclin dependent kinase 6                                |
| <i>CDKN1A</i> , <i>CDKN1B</i> , <i>CDKN2A</i> | cyclin-dependent kinase inhibitor 1A, 1B and 2A genes                               |
| cDNA  | complementary DNA (deoxyribonucleic acid)   |
| Cds1  | Schizosaccharomyces pombe gene  |
| CE  | catechol oestrogens   |

|                                   |   |
|-----------------------------------|---|
| CE-SQ, CE-Q                       | catechol oestrogen semi-quinone, catechol oestrogen quinone   |
| <i>CETN2</i>                      | centrin 2 gene  |
| CHEK2, <i>CHEK2</i> , <i>CHK2</i> | checkpoint kinase 2, checkpoint kinase 2 gene   |
| Chk2                              | mouse checkpoint kinase 2 gene  |
| CI                                | confidence interval   |
| cM                                | centi-Morgan  |
| CMF/CNF                           | adjuvant chemotherapy containing cyclophosphamide, methotrexate, 5-fluorouracil and mitoxantrone                                |
| <i>c-MYC</i>                      | human homolog of avian v-myc myelocytomatosis viral oncogene  |
| COMT, <i>COMT</i>                 | catechol-O-methyltransferase, catechol-O-methyltransferase gene   |
| CS                                | Cowden syndrome   |
| CSA, <i>CSB</i>                   | excision-repair cross-complementing group 8 and 6 genes   |
| <i>CSF2RB</i>                     | granulocyte-macrophage colony-stimulating factor receptor beta gene   |
| CSGE                              | conformation-sensitive gel electrophoresis  |
| CYPs, CYP450                      | cytochrome P450 subfamily enzymes   |
| CYP1A1, <i>CYP1A1</i>             | cytochrome P450 subfamily I polypeptide 1/cytochrome P450A1, cytochrome P450 subfamily I polypeptide 1/cytochrome P450A1 gene   |
| <i>CYP1A2</i>                     | cytochrome P450 subfamily I polypeptide 2 gene  |
| CYP1B1, <i>CYP1B1</i>             | cytochrome P450 subfamily I polypeptide 1/cytochrome P4501B1, cytochrome P450 subfamily I polypeptide 1/cytochrome P4501B1 gene |
| CYP11A                            | cytochrome P450 subfamily XIA polypeptide 1   |
| CYP17, <i>CYP17</i>               | steroid 17- $\alpha$ hydroxylase, steroid 17- $\alpha$ hydroxylase gene   |
| CYP19, <i>CYP19</i>               | aromatase/oestrogen synthetase, aromatase gene  |
| <i>DDB1</i> , <i>DDB2</i>         | DNA damage-binding protein 1 and 2 genes  |
| DHEA                              | dehydroepiandrosterone  |
| DHPLC                             | denaturing high-pressure liquid chromatography  |
| <i>DMC1</i>                       | human homolog of yeast disrupted meiotic cDNA 1 gene  |
| Dmnk                              | Drosophila melanogaster gene  |
| DNA                               | deoxyribonucleic acid   |
| DNA-PK, <i>DNA-PK</i>             | DNA-activated protein kinase catalytic subunit, DNA-activated protein kinase catalytic subunit gene                             |
| dNTP                              | deoxyribonucleotide triphosphate  |
| DSB, DSDB                         | double-stranded DNA break   |
| DSS1                              | deleted in split-hand/split-foot 1 region   |
| E1, E2                            | oestrone, oestradiol  |
| E2F1                              | E2F transcription factor 1  |
| <i>EDH17B2</i>                    | 17-beta-hydroxysteroid dehydrogenase I gene   |
| EDTA                              | ethylenediamine tetra-acetic acid   |
| EM                                | expectation maximization algorithm  |
| EMSY                              | protein encoded by EMSY gene  |
| ER                                | oestrogen receptor  |
| <i>ERBB2</i> = <i>HER2</i> /neu   | human homolog to avian v-erb-b2 erythroblastic leukaemia viral oncogene   |
| <i>ERCC1</i>                      | excision-repair complementing defective in Chinese hamster 1 gene   |



|                                   |   |
|-----------------------------------|---|
| <i>ERCC4=XPF</i>                  | excision-repair complementing defective in Chinese hamster 4 gene                         |
| <i>ESR1, ESR2</i>                 | oestrogen receptor alpha and beta genes   |
| EtBr                              | ethidium bromide  |
| FA                                | Fanconi anemia  |
| FA-N                              | Fanconi anemia complementation group N  |
| <i>FANCD2, FANCD2</i>             | Fanconi anemia complementation group D2, gene and protein                                 |
| FDH                               | Finnish disease heritage  |
| <i>FEN1</i>                       | flap structure-specific endonuclease 1 gene   |
| <i>FGFR2, FGFR2</i>               | fibroblast growth factor receptor 2, fibroblast growth factor receptor 2 gene             |
| FIGO                              | International Federation of Gynaecology and Obstetrics                                    |
| FokI                              | restriction fragment length polymorphism in <i>VDR</i> gene                               |
| GH, <i>GH1</i>                    | growth hormone, growth hormone gene   |
| <i>GHRH</i>                       | growth hormone releasing hormone gene   |
| <i>GHRHR, GHRHR</i>               | growth hormone releasing hormone receptor, growth hormone releasing hormone receptor gene |
| <i>GHRL</i>                       | growth hormone secretagogue receptor ligand gene  |
| <i>GHSR</i>                       | growth hormone secretagogue receptor gene   |
| GSH                               | glutathione   |
| GST                               | glutathione S-transferase   |
| <i>GSTM1, GSTM3, GSTP1, GSTT1</i> | glutathione S-transferase M1, M3, P1 and T1 genes   |
| GSTM1                             | glutathione S-transferase M1  |
| GSTM3                             | glutathione S-transferase M3  |
| GSTP1                             | glutathione S-transferase P1  |
| GSTT1                             | glutathione S-transferase T1  |
| <i>GTF2H1-4</i>                   | general transcription factor IIH polypeptide 1-4 genes                                    |
| GWS                               | genome-wide scan  |
| <i>H19</i>                        | H19 gene  |
| HA                                | heteroduplex analysis   |
| HBCC                              | hereditary breast and colorectal cancer   |
| HDAC1, HDAC2                      | histone deacetylase 1, histone deacetylase 2  |
| <i>HER2/NEU=ERBB2</i>             | human homolog to avian v-erb-b2 erythroblastic leukaemia viral oncogene                   |
| HLA                               | major histocompatibility complex  |
| <i>hMSH3</i>                      | human homolog 3 of <i>E. coli</i> MutS gene   |
| HNPCC                             | hereditary nonpolyposis colon cancer syndrome   |
| <i>hPMS1, hPMS2</i>               | human homologs of <i>S cerevisiae</i> postmeiotic segregation increased 1 and 2 genes     |
| HR                                | homologous recombination  |
| <i>HRAS1</i>                      | v-ha-ras Harvey rat sarcoma viral oncogene homolog gene                                   |
| htSNP                             | haplotype-tagging single nucleotide polymorphism  |
| HWE                               | Hardy-Weinberg equilibrium  |
| <i>ICAM5</i>                      | intercellular adhesion molecule 5 gene  |
| ID4                               | inhibitor of DNA binding 4  |
| IGF                               | insulin-like growth factor  |
| <i>IGF-I, IGF-II</i>              | insulin-like growth factor I and II genes   |
| IGF-I, IGF-II                     | insulin-like growth factor I and II   |
| IGF-IR, IGF-IIR                   | insulin-like growth factor I and II receptors   |

|                                   |  |
|-----------------------------------|--|
| IGFBP                             | insulin-like growth factor binding protein                           |
| <i>IGFBP3</i>                     | insulin-like growth factor binding protein 3 gene                    |
| IGFBP3                            | insulin-like growth factor binding protein 3                         |
| <i>IL2RB</i>                      | interleukin 2 receptor beta gene                                     |
| kb                                | kilobases  |
| KBCP                              | Kuopio Breast Cancer Project   |
| <i>KCTD17</i>                     | potassium channel tetramerization domain gene                        |
| kD                                | kilo Dalton  |
| <i>KU70</i>                       | thyroid autoantigen, 70-kD   |
| LD                                | linkage disequilibrium   |
| LFS, LFL                          | Li-Fraumeni syndrome, Li-Fraumeni-like                               |
| <i>LIG1, LIG3</i>                 | DNA ligase 1 and DNA ligase 3 genes                                  |
| <i>LIG4</i>                       | DNA ligase 4 gene  |
| <i>LKB1=STK11</i>                 | serine-threonine protein kinase 11 gene                              |
| LOD                               | logarithm of odds  |
| LOH                               | loss of heterozygosity   |
| <i>LSP1</i>                       | lymphocyte-specific protein gene                                     |
| Mb                                | mega bases   |
| MLH1, <i>MLH1</i> = <i>hMLH1</i>  | human homolog of E. coli MutL, human homolog of E. coli MutL gene    |
| MMR                               | mismatch repair  |
| MnSOD, <i>MnSOD</i> = <i>SOD2</i> | manganese superoxide dismutase, manganese superoxide dismutase gene  |
| MPO, <i>MPO</i>                   | myeloperoxidase, myeloperoxidase gene                                |
| <i>MPST</i>                       | mercaptopyruvate sulfurtransferase                                   |
| <i>MRE11</i>                      | homolog of S. cerevisiae meiotic recombination 11 gene               |
| mRNA                              | messenger ribonucleic acid   |
| MSH2, <i>MSH2</i> = <i>hMSH2</i>  | human homolog 2 of E. coli MutS, homolog 2 of E coli MutS gene       |
| MSH6, <i>hMSH6</i>                | human homolog 6 of E. coli MutS and homolog 6 of E coli MutS gene    |
| <i>MUTYH</i>                      | human homolog of E. coli MutY gene                                   |
| NADPH                             | nicotinamide adenine dinucleotide phosphate                          |
| NAT1, <i>NAT1</i>                 | N-acetyltransferase 1, N-acetyltransferase 1 gene                    |
| NAT2, <i>NAT2</i>                 | N-acetyltransferase 2, N-acetyltransferase 2 gene                    |
| NBR2                              | neighbour of BRCA1 gene 2  |
| NBS1, <i>NBS1</i>                 | nibrin, nibrin gene  |
| NCBI                              | National Center for Biotechnology Information                        |
| <i>NCF4</i>                       | neutrophil cytosolic factor 4 gene                                   |
| NER, GG-NER                       | nucleotide-excision repair, global genome nucleotide-excision repair |
| NHEJ                              | non-homologous end-joining   |
| NLS, NLS1                         | nuclear localization signal, nuclear localization signal 1           |
| NQO1, <i>NQO1</i>                 | NAD(P)H:quinone oxidoreductase, NAD(P)H:quinone oxidoreductase gene  |
| nt                                | nucleotide(s)  |
| <i>NTH1</i>                       | endonuclease III-like 1 gene   |
| <i>NUMA1</i>                      | nuclear mitotic apparatus protein 1 gene                             |
| OB1, OB2, OB3                     | oligonucleotide/oligosaccharide binding folds 1, 2 and 3             |
| OCCR                              | ovarian cancer cluster region  |

|                               |   |
|-------------------------------|---|
| <i>OGG1</i>                   | 8-oxoguanine DNA glycosylase gene   |
| OR                            | odds ratio  |
| PARP-1                        | poly(ADP-ribose)polymerase 1  |
| p16                           | cyclin-dependent kinase inhibitor 2A  |
| p21                           | cyclin-dependent kinase inhibitor 1A  |
| p53                           | tumour protein p53  |
| p73                           | tumour protein p73  |
| p185                          | <i>ERBB2</i> gene product, tumour antigen p185  |
| PAHs                          | polyaromatic hydrocarbons   |
| PALB2, <i>PALB2</i>           | partner and localizer of BRCA2, partner and localizer of<br>BRCA2 gene                                  |
| P/CAF                         | p300/CBP-associated factor  |
| <i>PCNA</i>                   | proliferating cell nuclear antigen gene   |
| PCR                           | polymerase chain reaction   |
| <i>PGR</i>                    | progesterone receptor gene  |
| Plk1                          | polo-like kinase 1  |
| PML                           | promyelotic leukaemia protein   |
| <i>POLB</i>                   | DNA polymerase $\beta$ gene   |
| poly-A                        | variable length poly(A) polymorphism in <i>VDR</i> gene   |
| POU1F1                        | POU domain class 1 transcription factor   |
| PR                            | progesterone receptor   |
| <i>PTEN=MMAC1=TEP1</i>        | phosphatase and tensin homolog gene   |
| PTT                           | protein truncation test   |
| <i>PVALB</i>                  | parvalbumin gene  |
| <i>RABL4</i>                  | RAB, member of RAS oncogene family-like 4gene   |
| <i>RAD23A, RAD23B</i>         | human homolog A and B of yeast Rad23 genes  |
| <i>RAD50</i>                  | human homolog of <i>S. cerevisiae</i> Rad50 gene  |
| <i>RAD51, RAD51</i>           | human homolog of <i>S. cerevisiae</i> Rad51, human homolog of <i>S.</i><br><i>cerevisiae</i> Rad51 gene |
| Rad51                         | <i>S.cerevisiae</i> Rad51   |
| <i>RAD51B, RAD51C, RAD51D</i> | human homolog B, C and D of <i>S. cerevisiae</i> Rad51 genes  |
| <i>RAD52</i>                  | human homolog of yeast Rad52 gene   |
| Rad53                         | <i>Saccharomyces cerevisiae</i> Rad53 gene  |
| <i>RAD54</i>                  | human homolog-like of <i>S. cerevisiae</i> Rad54 gene   |
| RB, <i>RBI</i>                | retinoblastoma protein, retinoblastoma gene   |
| RefSNP                        | reference single nucleotide polymorphism database   |
| RFC1, RFC2, RFC4              | replication factor C1, replication factor C subunit 2 and<br>replication factor C4                      |
| RFLP                          | restriction fragment length polymorphism  |
| RING-finger                   | a distinct zinc-chelating domain involved in mediating<br>protein-DNA and protein-protein interactions  |
| ROS                           | reactive oxygen species   |
| <i>RPA, RPA1-3</i>            | replication protein A1-A3 genes   |
| RT-PCR                        | reverse-transcriptase polymerase chain reaction   |
| SDS-PAGE                      | sodium dodecylsulphate polyacrylamide gel electrophoresis   |
| SHBG, <i>SHBG</i>             | sex hormone-binding globulin, sex hormone-binding globulin<br>gene                                      |
| SNP                           | single nucleotide polymorphism  |
| SOD                           | superoxide dismutase  |
| <i>SOD2</i>                   | manganese superoxide dismutase gene   |

|  |   |
|--|---|
| SSA  | single-strand annealing   |
| SSCP                                       | single-stranded conformation polymorphism                                   |
| SST, <i>SST</i>                            | somatostatin, somatostatin gene   |
| <i>SSTR1</i> , <i>SSTR2</i> , <i>SSTR5</i> | somatostatin receptor 1, 2 and 5 genes                                      |
| SSTR1-5                                    | somatostatin receptors 1-5  |
| <i>STK15=AURORA2=AURKA=BTAK=ARK1</i>       | serine threonine kinase gene  |
| STR  | short tandem repeat, microsatellite   |
| SULT                                       | sulfotransferase  |
| SULT1A1, <i>SULT1A1</i>                    | sulfotransferase 1A1, sulfotransferase 1A1 gene                             |
| SULT1E1, <i>SULT1E1</i>                    | sulfotransferase 1E1, sulfotransferase 1E1 gene                             |
| TaqI                                       | restriction fragment length polymorphism in <i>VDR</i> gene                 |
| TCR  | transcription-coupled repair  |
| TGFβ1, <i>TGFB1</i>                        | transforming growth factor beta, transforming growth factor beta gene       |
| <i>TMPRSS6</i>                             | transmembrane protease, serine 6, gene                                      |
| <i>TNFα</i> , <i>TNFβ</i>                  | tumour necrosis factor alpha and beta genes                                 |
| TNM  | tumour-node-metastasis classification                                       |
| <i>TNRC9=TOX3</i>                          | TOX high mobility group box family member 3 gene                            |
| <i>TP53</i>                                | tumour protein 53 gene  |
| <i>TST</i>                                 | thiosulphate sulfurtransferase  |
| TTK  | phosphotyrosine-picked threonine kinase                                     |
| UCSC                                       | University of California Santa Cruz   |
| UICC                                       | International Union Against Cancer  |
| USP11                                      | ubiquitin-specific protease 11  |
| UTR  | untranslated region   |
| UV   | ultraviolet   |
| <i>VDR</i> , <i>VDR</i>                    | vitamin D receptor, vitamin D receptor gene                                 |
| <i>VEGF</i> , <i>VEGF</i>                  | vascular endothelial growth factor, vascular endothelial growth factor gene |
| WHO  | World Health Organization   |
| VNTR                                       | variable number of tandem repeats   |
| <i>XAB2</i>                                | XPA-binding protein 2   |
| <i>XPA</i> , <i>XPG</i>                    | xeroderma pigmentosum complementation group A and G genes                   |
| <i>XPB=ERCC2</i>                           | excision-repair complementing defective in Chinese hamster 2 gene           |
| <i>XRCC1</i>                               | X-ray repair complementing defective in Chinese hamster 1                   |
| <i>XRCC1</i>                               | X-ray repair complementing defective in Chinese hamster 1 gene              |
| <i>XRCC2</i>                               | X-ray repair complementing defective in Chinese hamster 2 gene              |
| <i>XRCC3</i>                               | X-ray repair complementing defective in Chinese hamster 3 gene              |
| <i>XRCC4</i>                               | X-ray repair complementing defective in Chinese hamster 4 gene              |
| <i>XRCC5</i>                               | X-ray repair complementing defective in Chinese hamster 5 gene              |
| <i>ZNF350=ZBRK1</i>                        | zinc finger protein 350 gene  |

## LIST OF ORIGINAL PUBLICATIONS

This thesis is based on the following original publications referred to in the text by their Roman numerals I-V.

- I Hartikainen JM, Kataja V, Pirskanen M, Arffman A, Ristonmaa U, Vahteristo P, Ryyänänen M, Heinonen S, Kosma V-M, Mannermaa A: Screening for *BRCA1* and *BRCA2* mutations in Eastern Finnish breast/ovarian cancer families. *Clinical Genetics* 2007, 72:311-320.
- II Hartikainen JM, Mannermaa A, Heinonen S, Kosma VM, Kataja V: A *BRCA2* mutation, 4088insA, in a Finnish breast and ovarian cancer family associated with favourable clinical course. *Anticancer Research* 2007, in press.
- III Hartikainen JM, Pirskanen MM, Arffman AH, Ristonmaa UK, Mannermaa AJ: A Finnish *BRCA1* exon 12 4216-2nt A to G splice acceptor site mutation causes aberrant splicing and frameshift, leading to protein truncation. *Human Mutation* 2000, 15:120.
- IV Hartikainen JM, Tuhkanen H, Kataja V, Dunning AM, Antoniou A, Smith P, Arffman A, Pirskanen M, Easton DF, Eskelinen M, Uusitupa M, Kosma VM, Mannermaa A: An autosome-wide scan for linkage disequilibrium-based association in sporadic breast cancer cases in Eastern Finland: three candidate regions found. *Cancer Epidemiology, Biomarkers & Prevention* 2005, 14:75-80.
- V Hartikainen JM, Tuhkanen H, Kataja V, Eskelinen M, Uusitupa M, Kosma VM, Mannermaa A: Refinement of the 22q12-q13 breast cancer-associated region: evidence of *TMPRSS6* as a candidate gene in an Eastern Finnish population. *Clinical Cancer Research* 2006, 12:1454-1462.

This thesis includes also unpublished data. The original papers in this thesis have been reproduced with the permission of the publishers.



## CONTENTS

|  |           |
|--|-----------|
| <b>1. INTRODUCTION.....</b>  | <b>19</b> |
| <b>2. REVIEW OF THE LITERATURE.....</b>  | <b>22</b> |
| <b>2.1 Breast cancer .....</b>   | <b>22</b> |
| 2.1.1 Incidence of breast cancer .....   | 22        |
| 2.1.2 Epidemiology and risk factors for breast cancer .....                              | 22        |
| 2.1.3 Prognostic factors in breast cancer .....  | 24        |
| <b>2.2 Ovarian cancer .....</b>  | <b>24</b> |
| 2.2.1 Incidence of ovarian cancer.....   | 24        |
| 2.2.2 Epidemiology and risk factors for ovarian cancer .....                             | 25        |
| 2.2.3 Prognostic factors in ovarian cancer .....   | 25        |
| <b>2.3 Familial breast cancer and high-risk genes.....</b>                               | <b>26</b> |
| 2.3.1 <i>BRCA1</i> and <i>BRCA2</i> genes .....  | 27        |
| 2.3.2 The <i>BRCA1/2</i> proteins and their functions.....                               | 28        |
| 2.3.2.1 Protein structure and functional motifs .....                                    | 28        |
| 2.3.2.2 Proposed functions .....   | 29        |
| 2.3.3 Cancer risks conferred by mutations in the <i>BRCA</i> genes.....                  | 32        |
| 2.3.3.1 Variation in cancer risks conferred by <i>BRCA</i> mutations.....                | 33        |
| 2.3.4 Clinical features of <i>BRCA</i> gene mutation-associated cancers.....             | 33        |
| 2.3.5 Mutation spectrum and founder mutations in the <i>BRCA</i> genes.....              | 35        |
| 2.3.6 <i>BRCA</i> mutations in Finland.....  | 36        |
| 2.3.7 Common polymorphisms and <i>BRCA</i> genes in sporadic breast/ovarian cancer ..... | 38        |
| 2.3.8 Other genes involved in familial breast cancer .....                               | 40        |
| 2.3.8.1 <i>PTEN</i> and Cowden syndrome .....  | 40        |
| 2.3.8.2 <i>TP53</i> and Li-Fraumeni syndrome .....                                       | 40        |
| 2.3.8.3 <i>ATM</i> .....   | 42        |
| 2.3.8.4 Checkpoint kinase 2, <i>CHEK2</i> ( <i>CHK2</i> ).....                           | 43        |
| 2.3.8.4.1 <i>CHEK2</i> gene and protein function.....                                    | 43        |
| 2.3.8.4.2 <i>CHEK2</i> germline mutations and risk of breast cancer .....                | 44        |
| 2.3.8.5 <i>LKB1</i> and Peutz-Jeghers syndrome.....                                      | 48        |
| 2.3.8.6 <i>PALB2</i> and Fanconi anemia.....   | 48        |
| <b>2.4 Sporadic breast cancer and low-penetrance susceptibility genes.....</b>           | <b>50</b> |
| 2.4.1 Steroid hormone biosynthesis, signalling and metabolism genes .....                | 54        |
| 2.4.1.1 Steroid hormone biosynthesis genes .....   | 54        |
| 2.4.1.2 Steroid hormone signalling genes .....   | 54        |
| 2.4.1.3 Steroid hormone metabolism genes .....   | 56        |
| 2.4.2 Carcinogen and xenobiotics metabolism .....  | 60        |
| 2.4.3 Other hormone- and growth factor-related genes.....                                | 61        |
| 2.4.3.1 Insulin-like growth factor family .....  | 61        |
| 2.4.3.2 Vitamin D receptor, <i>VDR</i> .....   | 62        |
| 2.4.3.3 Vascular endothelial growth factor, <i>VEGF</i> .....                            | 64        |

|            |  |           |
|------------|--|-----------|
| 2.4.3.4    | Transforming growth factor beta, <i>TGFB1</i> .....  | 64        |
| 2.4.4      | Genes involved in double-stranded DNA break repair .....   | 65        |
| 2.4.4.1    | <i>BARD1</i> .....   | 66        |
| 2.4.5      | Genes involved in single-stranded DNA damage repair mechanisms .....                                       | 67        |
| 2.4.5.1    | Base-excision repair .....   | 67        |
| 2.4.5.2    | Nucleotide-excision repair .....   | 68        |
| 2.4.5.3    | Mismatch repair .....  | 68        |
| 2.4.6      | Genes involved in cell cycle control.....  | 69        |
| 2.4.7      | Proto-oncogenes .....  | 70        |
| 2.4.8      | Other genes studied for breast cancer association .....  | 71        |
| <b>2.5</b> | <b>Strategies for finding new breast cancer susceptibility genes.....</b>                                  | <b>73</b> |
| 2.5.1      | Linkage disequilibrium .....   | 73        |
| 2.5.2      | Positional cloning of Mendelian genes .....  | 76        |
| 2.5.3      | Strategies for finding genes for complex diseases.....   | 77        |
| 2.5.3.1    | Case-control association study.....  | 77        |
| 2.5.3.1.1  | Genome-wide approach.....  | 78        |
| 2.5.3.1.2  | Candidate-gene approach .....  | 78        |
| 2.5.4      | The Finnish population .....   | 80        |
| 2.5.4.1    | The Finnish Disease Heritage .....   | 82        |
| <b>2.6</b> | <b>Methods for mutation detection.....</b>   | <b>83</b> |
| 2.6.1      | Heteroduplex analysis, HA.....   | 83        |
| 2.6.2      | Single-strand conformation polymorphism, SSCP/Conformation-sensitive gel electrophoresis, CSGE.....        | 84        |
| 2.6.3      | Protein truncation test, PTT.....  | 84        |
| 2.6.4      | Restriction fragment length polymorphism, RFLP.....  | 85        |
| 2.6.5      | Sequencing .....   | 85        |
| <b>3.</b>  | <b>AIMS OF THE STUDY .....</b>   | <b>87</b> |
| <b>4.</b>  | <b>MATERIAL AND METHODS.....</b>   | <b>88</b> |
| <b>4.1</b> | <b>Study subjects.....</b>   | <b>88</b> |
| 4.1.1      | Breast and ovarian cancer families (I-III) .....   | 88        |
| 4.1.2      | Population-based breast cancer case-control material (IV-V).....   | 90        |
| 4.1.2.1    | Cases and controls in the autosome-wide scan for LD (IV).....  | 91        |
| 4.1.2.2    | Cases and controls in refining of the breast cancer-associated region on 22q12-q13 (V) .....               | 91        |
| <b>4.2</b> | <b>Extraction of DNA from blood lymphocytes (I-V).....</b>   | <b>91</b> |
| <b>4.3</b> | <b><i>BRCA</i> gene mutation detection (I-III) .....</b>   | <b>92</b> |
| 4.3.1      | PCR amplification and protein truncation test (PTT) of <i>BRCA1</i> and <i>BRCA2</i> exon 11 (I, II) ..... | 92        |
| 4.3.2      | Heteroduplex analysis of <i>BRCA1</i> and <i>BRCA2</i> genes (I, III).....                                 | 94        |
| 4.3.3      | Sequencing (I-III, V).....   | 94        |
| 4.3.4      | RFLP analysis, extraction of RNA, and cDNA analysis of the <i>BRCA1</i> 4216-2nt A→G mutation (III) .....  | 97        |
| 4.3.5      | Analysis of mutation-associated haplotypes (I-III) .....   | 99        |



|         |   |     |
|---------|---|-----|
| 4.4     | Genotyping of microsatellites in the autosome-wide scan for LD (IV).....  | 99  |
| 4.5     | Validating and selecting of SNPs for the refining of the breast cancer-associated region on 22q12-q13 (V) ..... | 101 |
| 4.6     | Genotyping of SNPs for the refining of the breast cancer-associated region on 22q12-q13 (V) .....               | 105 |
| 4.7     | Allelic imbalance (AI) analysis (V) .....   | 105 |
| 4.7.1   | Paraffin-embedded samples and extraction of DNA for AI analysis .....   | 105 |
| 4.7.2   | Genotyping of microsatellites for AI analysis.....  | 107 |
| 4.7.3   | Calculation of AI values.....   | 108 |
| 4.8     | Statistical analyses (IV, V) .....  | 108 |
| 4.8.1   | Microsatellite and SNP allele and genotype frequencies (IV, V).....   | 108 |
| 4.8.2   | Hardy-Weinberg equilibrium (IV, V) .....  | 108 |
| 4.8.3   | Breast cancer risk estimations (Odds ratios, OR) (IV, V).....   | 109 |
| 4.8.4   | Haplotype estimation (IV, V).....   | 109 |
| 4.8.5   | Pairwise linkage disequilibrium (V) .....   | 110 |
| 4.8.6   | Power estimations (V).....  | 110 |
| 4.9     | Ethical aspects .....   | 110 |
| 5.      | RESULTS .....   | 111 |
| 5.1     | <i>BRCA1</i> and <i>BRCA2</i> mutations in breast/ovarian cancer families (I-III) .....                         | 111 |
| 5.1.1   | <i>BRCA1</i> mutations found (I, III) .....   | 111 |
| 5.1.1.1 | <i>BRCA1</i> exon 12 4216-2nt A→G (III) .....   | 112 |
| 5.1.1.2 | <i>BRCA1</i> exon 20 5370 C→T (I).....  | 113 |
| 5.1.2   | <i>BRCA2</i> mutations found (I, II) .....  | 113 |
| 5.1.2.1 | <i>BRCA2</i> exon 9 999del5 (I).....  | 114 |
| 5.1.2.2 | <i>BRCA2</i> exon 11 4088insA (II).....   | 114 |
| 5.1.2.3 | <i>BRCA2</i> exon 11 6503delTT (I) .....  | 114 |
| 5.1.3   | Haplotype analysis (I, II).....   | 115 |
| 5.2     | Autosome-wide scan for linkage disequilibrium-based association with breast cancer (IV).....                    | 116 |
| 5.2.1   | Allelic association.....  | 116 |
| 5.2.2   | Genotype association and Hardy-Weinberg equilibrium (HWE).....  | 117 |
| 5.2.3   | Haplotypes.....   | 117 |
| 5.3     | Further refinement of chromosome 22q12-q13 breast cancer-associated region (V).....                             | 118 |
| 5.3.1   | SNP allelic and genotype association .....  | 118 |
| 5.3.2   | Hardy-Weinberg equilibrium .....  | 118 |
| 5.3.3   | Pairwise linkage disequilibrium .....   | 120 |
| 5.3.4   | Haplotypes.....   | 120 |
| 5.3.5   | Allelic imbalance analysis.....   | 121 |
| 6.      | DISCUSSION .....  | 122 |
| 6.1     | High-risk susceptibility genes in Eastern Finnish breast and ovarian cancer families (I-III) .....              | 122 |

|            |  |            |
|------------|--|------------|
| 6.1.1      | Eastern Finnish <i>BRCA1</i> and <i>BRCA2</i> mutation frequency is not different from other parts of Finland (I)..... | 122        |
| 6.1.2      | Limited <i>BRCA1</i> and <i>BRCA2</i> mutation spectrum in Eastern Finland (I)..                                       | 124        |
| 6.1.3      | Geography and haplotypes in the <i>BRCA</i> mutation-carrier families (I, II)  | 125        |
| 6.1.4      | A novel Eastern Finnish <i>BRCA2</i> mutation and favourable prognosis (II)  | 126        |
| <b>6.2</b> | <b>Sporadic breast cancer and new low-penetrance susceptibility genes in Eastern Finland (IV, V).....</b>              | <b>128</b> |
| 6.2.1      | Autosome-wide scan for LD-based breast cancer association (IV).....  | 128        |
| 6.2.1.1    | Candidate-genes for further studies on breast cancer association (IV).....   | 130        |
| 6.2.2      | Breast cancer association on chromosome 22q12-q13 (V).....   | 132        |
| 6.2.2.1    | <i>TMPRSS6</i> as a candidate-gene (V).....  | 132        |
| 6.2.2.2    | Other genes on chromosome region 22q12-q13 (V).....  | 134        |
| 6.2.2.3    | Allelic imbalance on chromosome region 22q12-q13 (V).....  | 134        |
| <b>7.</b>  | <b>SUMMARY AND CONCLUSIONS.....</b>  | <b>135</b> |
| <b>8.</b>  | <b>REFERENCES.....</b>   | <b>137</b> |
| <b>9.</b>  | <b>ORIGINAL PUBLICATIONS.....</b>  | <b>187</b> |

## 1. INTRODUCTION

Breast cancer is the most common cancer among women in industrialised countries. Approximately one in ten women in Finland will develop breast cancer during her lifetime. Currently more than 4000 new breast cancer cases are diagnosed yearly and the number is increasing (Finnish Cancer Registry). At year 2005 ovarian cancer was the ninth most common cancer in the Finnish female population accounting 424 new ovarian carcinomas (Finnish Cancer Registry).

Many risk factors (hormonal, environmental) for breast cancer are known but most of the genetic background and molecular mechanisms still remain to be elucidated. Although mutations in *BRCA1* (Hall et al. 1990, Miki et al. 1994) and *BRCA2* (Wooster et al. 1994, Wooster et al. 1995) genes are known to confer a high lifetime risk of breast cancer, together with the other susceptibility genes so far identified, *ATM* (Swift et al. 1987, Easton 1994, Athma et al. 1996), *CHEK2* (Bell et al. 1999, CHEK2-Breast Cancer Consortium 2002), *TP53* (Easton 1999), *PALB2* (Rahman et al. 2007), and *BRIP1* (Seal et al. 2006), these genes explain only  $\leq 25\%$  of the familial aggregation of breast cancer (Antoniou and Easton 2006).

Several linkage studies in large family material have been conducted but other *BRCA1/2*-like, high-risk, high-penetrance breast cancer susceptibility genes have not been found. Therefore, the yet unidentified genes presumably are numerous and confer a moderate risk (Thompson and Easton 2004). In a study investigating the genetic models of the non-*BRCA1/2* familial clustering of breast cancer the findings suggest that several common, low-penetrance genes may account for the residual familial aggregation of breast cancer (Antoniou et al. 2002). It is most likely that low-penetrance genes for breast cancer susceptibility are present in the general population (Pharoah et al. 1997).

In Finland, 26 different *BRCA1* and *BRCA2* mutations have been found among female and male breast cancer and ovarian cancer patients (Vehmanen et al. 1997a, Vehmanen et al. 1997b, Roth et al. 1998, Huusko et al. 1998, Sarantaus et al. 2000, Syrjäkoski et al. 2000, Sarantaus et al. 2001a, Sarantaus et al. 2001b, Pääkkönen et al. 2001, Eerola et al. 2001, Vahteristo et al. 2001a, Syrjäkoski et al. 2004a). Fourteen of

these, seven in each gene, are considered as founder mutations as they are recurrent and account for the majority of all detected *BRCA1* and *BRCA2* mutations in Finland. Some of these mutations are unique to the Finns. Due to the population history the Finns are genetically isolated homogeneous population and founder effects can be seen in several autosomal recessive diseases and cancer predisposition syndromes (Norio et al. 1973, Nyström-Lahti et al. 1994, Moisio et al. 1996). This effect has been observed among *BRCA1* and *BRCA2* mutations also and it has an impact on diagnostics as well (Sarantaus et al. 2000). The proportion of *BRCA1* and *BRCA2* mutations among Southern, Western and Northern Finnish breast/ovarian cancer families has been studied but in Eastern Finnish population it has not been reported yet.

Linkage disequilibrium (LD) -based genetic association studies are suitable tools for detecting low-penetrance susceptibility genes that likely interact with environmental and lifestyle factors as well as with other genetic factors to cause disease. Young, isolated populations (e.g. the Eastern Finns) may provide more help by reducing the genetic heterogeneity. LD, or allelic association, is based on the assumption that the affected share a genetic variant/mutation, which is so close to the marker that the probability of a recombination event to occur between them is minimal. In young (15-25 generations) isolated populations the number of meioses is relatively low and LD is thought to extend further than in older populations (Ophoff et al. 2002). LD analysis has been used to discover genes for several diseases of the Finnish disease heritage e.g. neuronal ceroid lipofuscinosis and diastrophic dysplasia (Järvelä 1991, Hästbacka et al. 1992), as well as in other isolated populations and complex diseases like severe bipolar disorder and congenital muscular dystrophy (Ophoff et al. 2002, Toda et al. 1996). In detection of susceptibility genes for complex disease also SNP haplotypes have been successfully used in the Finnish population (Laitinen et al. 2004).

In the present study the proportion of *BRCA1* and *BRCA2* germ-line mutations was evaluated in breast/ovarian cancer families of Eastern Finnish origin. Also, to find new genetic medium or low-penetrance risk factors for sporadic breast cancer, the autosomes were screened utilizing linkage disequilibrium-based association and 435 microsatellites in a breast cancer case-control set from Eastern Finland. One of the associated regions

(22q12-q13) was further investigated by SNP association analysis to identify the putative breast cancer related gene.

## **2. REVIEW OF THE LITERATURE**

### **2.1 Breast cancer**

#### **2.1.1 Incidence of breast cancer**

Breast cancer is the most common cancer among women in industrialised countries. Each year over one million new cases are diagnosed in the world and it is the leading cause of cancer death (Globocan 2002). The incidence is highest in Northern America, Western and Northern Europe, and Australia/New Zealand (Globocan 2002). In Finland the proportion of breast cancer is approximately one third (31.8 % in 2005) of all female cancers (Finnish Cancer Registry). One in ten women in our country will be affected with breast cancer at some point in her life. Every year more than 3 500 new cases are diagnosed and the number is increasing, the number of new cases in 2005 already being 4 027 (Finnish Cancer Registry). The age-adjusted incidence of breast cancer was 86.7 per 100 000 person years in the year 2005 in Finland (Finnish Cancer Registry). In 2005 the number of breast cancer deaths was 804, i.e. the age-adjusted mortality rate was 14.7 per 100 000 person years. The age-adjusted mortality rate has remained rather unchanged since the 1950's (14 - 16 / 100 000 person years). Thus, a relative decline in breast cancer mortality has been observed following the initiation of the population based mammography screening programme in 1987 (Finnish Cancer Registry 2002 and 2004, Botha et al 2003). Male breast cancer in Finland is rare, accounting for less than 1 % of all breast cancers annually (Finnish Cancer Registry 2002).

#### **2.1.2 Epidemiology and risk factors for breast cancer**

Oestrogens have an important role in the development and progression of breast cancer. The life-time exposure to oestrogen is related to the risk of developing the disease (Pike et al 1979). Thus, the risk of breast cancer rises throughout a woman's lifetime (Finnish Cancer Registry 2002). Breast cancer before the age of 25 years is rare, except in certain familial cases (Finnish Cancer Registry 2002). Risk factors for breast cancer are early age at menarche (MacMahon et al 1982), nulliparity or delayed first childbirth (Layde et al. 1989, Ewertz et al. 1990, Kelsey et al. 1993), short duration of breast-feeding (Layde

et al. 1989, Kelsey et al. 1993), low number of children (Layde et al. 1989, Ewertz et al. 1990), late menopause (Trichopoulos et al. 1972, Collaborative Group on Hormonal Factors in Breast Cancer 1997), postmenopausal obesity (Folsom et al. 1990, Hunter and Willett 1993), extended use of oral contraceptives (Collaborative Group on Hormonal Factors in Breast Cancer 1996) and long-term oestrogen replacement therapy (Collaborative Group on Hormonal Factors in Breast Cancer 1997), which all are surrogates for oestrogen exposure. Higher education and socio-economic status has been found to associate with increased breast cancer risk, which is largely explained by nulliparity, later age at first childbirth and greater use of synthetic hormones (Heck and Pamuk 1997, Pukkala and Weiderpass 1999).

Increased risk of breast cancer is associated with carcinoma of the contralateral breast, a history of benign breast disease e.g. atypical hyperplasia and sclerosing adenosis (Dupont and Page 1985, Carter et al. 1988, Wang et al. 2004, Bernstein et al. 1992), and ionising radiation (Howe and McLaughlin 1996, Preston et al. 2002, Pukkala et al. 2006). Also cigarette smoke has been associated with an increase in risk for breast cancer (Gram et al. 2005, Johnson 2005) and accumulation of environmental oestrogens, xeno-oestrogens, may have a role in breast cancer aetiology (Davis et al. 1993). Physical inactivity and several dietary factors such as fat or energy intake and high consumption of cooked red meat (Bernstein et al. 1994, Bartsch et al. 1999, Nair et al. 1999, Zheng et al. 1998) as well as high alcohol consumption enhance breast cancer risk (Garfinkel et al. 1988, Smith-Warner et al. 1998). Many epidemiological studies have observed a positive association between adult height and risk of breast cancer, although the explanation for this is not clear (Tretli 1989, Vatten and Kvinnsland 1990, Hunter and Willett 1993). Also race and ethnicity, national origin and geographical location have an influence on breast cancer risk (Ziegler et al. 1993).

Family history of breast/ovarian cancer is a known risk factor. It is recognized that a woman with a first-degree relative with breast cancer has twice the risk of developing the disease herself (Pharoah et al. 1997). Genetic susceptibility accounts for 5-10 % of all cases of breast cancer (Claus et al. 1996). Mutations in *BRCA1* and *BRCA2* genes account for approximately 15 % of the excess familial relative risk of breast cancer (Peto et al. 1999, Anglian Breast Cancer Study Group 2000) and in total the so far

known breast cancer susceptibility genes account for no more than 25 % of the familial aggregation of breast cancer (Easton 1999). In addition, numerous low-penetrance genetic variants exist and these contribute together with environmental and lifestyle factors and with other genes to cause breast cancer (Antoniou et al. 2002).

### **2.1.3 Prognostic factors in breast cancer**

Breast cancer is predominantly a postmenopausal disease and appearance at young age, <35 years, is related to aggressive disease and reduced survival (Shannon and Smith 2003). Tumour stage is the most important prognostic factor in breast cancer and it is the basis for selecting patients for different treatment strategies. Breast cancer stage is assessed according to the TNM classification of the UICC and it involves tumour size, lymph node status and distant metastases (International Union Against Cancer 2002). Histological grade, based on tubule formation, nuclear pleiomorphism, and the number of mitoses, has independent prognostic value in breast cancer (WHO 2003). Expression of oestrogen and progesterone receptors in the tumour tissue predicts a good response to hormonal therapy of breast cancer (Clarke et al. 2004), whereas *ERBB2* (*HER2/neu*) gene amplification and/or protein overexpression in the tumour tissue is associated with poor prognosis (Ross et al. 2003). Gene expression profiling of breast tumours has enabled classification of tumours by their gene expression patterns, and provided insights into development of new potential diagnostic and prognostic markers (Perou et al. 2000, van't Veer et al. 2002, Sorlie et al. 2003, Chang et al. 2005).

## **2.2 Ovarian cancer**

### **2.2.1 Incidence of ovarian cancer**

Ovarian cancer is among the most common cancers in women worldwide with over 200,000 new cases annually. The incidence rates are highest in Europe, North America and Australia/New Zealand (Globocan 2002). In Finland, ovarian cancer was the ninth most common of female cancers accounting for 424 new cases in year 2005 (Finnish Cancer Registry). In 2005 the ovarian cancer incidence was 8.4 and mortality rate 5.0



per 100 000 person years. In addition to ovarian carcinomas, approximately 110 borderline tumours are diagnosed annually. The estimation of the number of new cases of ovarian cancer in 2006 is 507. (Finnish Cancer Registry).

### 2.2.2 Epidemiology and risk factors for ovarian cancer

The risk factors for ovarian cancer include genetic and environmental (hormonal and lifestyle) factors, many of them shared with breast cancer risk. Ageing and family history of ovarian or breast cancer increase the risk. Also hormone replacement therapy has been associated with increased risk. The number of full-term pregnancies and use of oral contraceptives are inversely correlated with risk, and late age at menarche, early age at menopause and breastfeeding modestly reduce the risk. Also tubal sterilisation has been associated with reduction in risk. The effects of obesity, physical activity and fertility treatment are unclear. (Bertone-Johnson 2005).

It is established that at least 10 % of all epithelial ovarian cancers are hereditary, with the *BRCA* genes contributing to at least 90 % of these cases and small percentages attributable to hereditary nonpolyposis colorectal cancer (HNPCC) syndrome (*MSH2* and *MLH1* genes) and perhaps some yet to be discovered susceptibility gene (Boyd 2003). It is presumed that genetic and environmental factors affect the penetrance of *BRCA* gene mutations for ovarian cancer but very few factors have been identified to date. One such factor is the *HRAS1* variable number of tandem repeat (VNTR). The rare alleles have been suggested to increase the ovarian (but not breast) cancer risk in *BRCA1* mutation heterozygotes (Phelan et al. 1996). *HRAS1* VNTR rare alleles have been shown to increase the relative risk of ovarian cancer in general population also (Weitzel et al. 2000).

### 2.2.3 Prognostic factors in ovarian cancer

The most important prognostic factor in ovarian cancer is the FIGO (International Federation of Gynaecology and Obstetrics) stage. Stage I tumours are limited to ovaries, stage II to ovaries with pelvic extension, stage III to ovaries with peritoneal implants and stage IV to ovaries with distant metastasis. (Cancer Committee of the FIGO, 1986).

Histological grade describing the tumour differentiation is also associated with prognosis (WHO 2003). The best-known and well-described serum marker for ovarian cancer is the CA125 antigen (Whitehouse and Solomon 2003).

### 2.3 Familial breast cancer and high risk genes

Breast cancer attributable to family history of the disease has been reported to account for 5-10 % of all cases of breast cancer (Claus et al. 1996). A number of genes exist with a proven high penetrance to familial breast cancer when mutated. High-risk breast and ovarian cancer susceptibility genes, *BRCA1* (Hall et al. 1990, Miki et al. 1994) and *BRCA2* (Wooster et al. 1994, Wooster et al. 1995), are estimated to explain ~15-20 % of the excess familial relative risk of breast cancer (Table 1) (Peto et al. 1999, Anglian Breast Cancer Study Group 2000). Other loci for breast cancer susceptibility have also been identified, including *PTEN*, *TP53*, *ATM*, *CHEK2*, *BRIP1* and *PALB2*, but these are of lower frequency in the general population (Table 1) (Easton 1999, CHEK2-Breast Cancer Consortium 2002, Seal et al. 2006, Rahman et al. 2007). Moderate family history of breast cancer has been shown to independently predict breast cancer risk without associated *BRCA1* or *BRCA2* mutations (Pharoah et al. 1997). Overall, breast

**Table 1.** Breast cancer risks conferred by inactivating mutations in known susceptibility genes.

| Gene         | Overall relative risk of breast cancer | Contribution to   |                   | References  |
|--------------|--|-------------------|-------------------|---|
|              |  | Breast cancer (%) | Familial risk (%) |   |
| <i>BRCA1</i> | >30 at <40 years<br>14 at >60 years    | 1-2               | *                 | Thompson and Easton 2004, Antoniou et al. 2003    |
| <i>BRCA2</i> | 11                                     | 1-2               | *                 | Thompson and Easton 2004, Antoniou et al. 2003    |
| <i>ATM</i>   | 2.37                                   | 0.86              | †                 | Renwick et al. 2006                               |
| <i>CHEK2</i> | 2.34                                   | 0.7               | 0.5 <sup>†</sup>  | CHEK2 Breast Cancer Case-Control Consortium, 2004 |
| <i>PALB2</i> | 2.3                                    | 0.23              | 0.24 <sup>†</sup> | Rahman et al. 2007                                |
| <i>BRIP1</i> | 2.0 (3.5 at <50)                       | 0.2               | †                 | Seal et al. 2006                                  |


\*Collectively these account for ~15-20 % of the overall familial risk of breast cancer (Peto et al. 1999, Anglian Breast Cancer Study Group 2000).

<sup>†</sup>Collectively these account for ~2.3 % of the overall familial relative risk (Rahman et al. 2007).

cancer is approximately twice as common in women with an affected first-degree relative; this risk increases with the number of affected relatives and is greater for women with relatives affected at a young age (Collaborative Group on Hormonal Factors in Breast Cancer 2001). However, in families with both breast and ovarian cancers, the aggregation of these two cancers appears to be explained by *BRCA1/BRCA2* mutation-carrier probability (Claus et al. 1998).

### 2.3.1 *BRCA1* and *BRCA2* genes

In 1990 linkage to marker *D17S74* on chromosome 17q21 was detected in a study using 23 families with 146 breast cancer cases (Hall et al. 1990). In most of the families the typical features for hereditary breast cancer were seen; early age of onset, bilateral cases and male breast cancer. Later the *BRCA1* (Breast Cancer 1 gene) gene was identified and cloned on 17q21 (Miki et al. 1994). In another linkage study using 22 families with at least one male breast cancer case strong negative LOD score values were obtained suggesting that male breast cancer is not linked to the *BRCA1* locus on 17q (Stratton et al. 1994). An analysis using 15 families with no linkage to *BRCA1* locus revealed linkage to 13q12-q13, which was presumed to be the locus for *BRCA2* (Breast Cancer 2 gene) (Wooster et al. 1994). *BRCA2* gene was cloned on chromosome 13q12 in 1996. Both genes encode charged and exceptionally large proteins. (Tavtigian et al. 1996). (Figure 1).



**Figure 1.** Genomic structure of the *BRCA1* and *BRCA2* genes. *BRCA1* gene spans over 80 kb of genomic sequence consisting of 24 exons and *BRCA2* approximately 70 kb with 27 exons (Miki et al. 1994, Tavtigian et al. 1996). Modified from Håkansson et al. 1997.

## 2.3.2 The BRCA1/2 proteins and their functions

### 2.3.2.1 Protein structure and functional motifs


Twenty-two of the 24 exons of *BRCA1* gene encode an 1863 amino-acid-protein (Miki et al. 1994). The protein is expressed in several tissues e.g. in ovary, testis and mammary gland epithelial cells (Miki et al. 1994). The 3418 amino-acid BRCA2 protein is encoded by 25 of the 27 exons of *BRCA2* gene (Tavtigian et al. 1996). The BRCA2 protein does not share sequence homology with any other protein (Wooster et al. 1995, Tavtigian et al. 1996). Like BRCA1, the BRCA2 protein is expressed in several tissues, including mammary gland, spleen, ovary, lung, testis and thymus (Tavtigian et al. 1996). Although BRCA1 and BRCA2 proteins have only little resemblance to each other or other proteins of known function, they contain structural motifs that give clues of their biochemical functions. (Figure 2).

In its amino-terminal region BRCA1 contains a zinc binding RING-finger motif that is found in regulatory proteins and appears to be involved in protein-protein interactions (Miki et al. 1994, Saurin et al. 1996). This motif has a role in e.g. the interaction of BARD1 (BRCA1-associated RING domain protein) and BAP1 (BRCA1-interacting protein 1) with BRCA1, as well as in BRCA1 homodimer formation (Wu et al 1996, Jensen et al. 1998, Brzovic et al. 1998). Two nuclear localization signals (NLS) locate in BRCA1 exon 11 but presumably only one (NLS1) of them is needed for the nuclear localization of BRCA1 (Chen et al. 1996, Thakur et al. 1997). The carboxy terminal-region of BRCA1 contains a conserved BRCT (BRCA1 C-terminal) domain that acts as a protein-binding site and interacts with multiple transcription factors (Koonin et al. 1996), and a region that has been shown to have transcriptional activation potential (Chapman & Verma 1996). (Figure 2).


In the C-terminal region of BRCA2 protein locates three oligonucleotide/oligosaccharide binding folds (OB1, OB2, OB3) that bind DSS1 and single-stranded DNA (Yang et al. 2002a). This provided a structural and biochemical basis for understanding the loss of recombination-mediated double-strand break repair in BRCA2-associated cancers (Yang et al. 2002a). Also in the amino-terminal region within exon 3 of BRCA2 has been shown to have transcriptional activation potential (Milner et al. 1997). In its' central part, in exon 11, BRCA2 contains eight conserved sequence motifs, the

BRC repeats that interact directly with RAD51 protein (Bork et al. 1996, Bignell et al. 1997, Wong et al. 1997). Within the final 156 residues in the carboxy-terminal of BRCA2 resides two nuclear localization signals (NLS) that are required for the nuclear localization and proper function of the BRCA2 protein (Spain et al. 1999). (Figure 2). In addition, both *BRCA1* and *BRCA2* have been suggested to have properties of granins as they include a motif similar to the granin consensus at the C-terminus of the protein (Jensen et al. 1996).

### BRCA1 1863 aa



### BRCA2 3418 aa



**Figure 2.** Structural domains and functional motifs of BRCA1 and BRCA2 proteins. The borders of exon 11 are shown with black lines. NLS=nuclear localization signal; OB1, OB2, OB3=oligonucleotide/oligo-saccharide binding folds; OCCR=ovarian cancer cluster region. Some of the most important interacting proteins are shown below the proteins. See text for details. Modified from Borg 2001.

#### 2.3.2.2 Proposed functions

BRCA1 and BRCA2 proteins interact with a number of regulatory proteins and they seem to have multiple fundamental functions. BRCA1 interacts directly and indirectly with numerous molecules, including tumour suppressors, oncogenes, DNA damage

repair proteins, cell cycle regulators, transcriptional activators and repressors, e.g. BRCA2, BARD1, ATM, ATR, CHEK2, BLM, MSH2, MSH6, MLH1 and RFC1-RFC2-RFC4 complex (Wang et al. 2000, Borg 2001, Welcsh and King 2001). BRCA2 interacts with Rad51, BRCA1, DSS1, EMSY, P/CAF, BRAF35, Plk1, USP11, FANCD2 and G, androgen receptor (AR), BCCIP $\alpha/\beta$  and PALB2 (Xia et al. 2006). Both BRCA1 and BRCA2 proteins localize to the nucleus of dividing cells and work in pathways that are required for the maintenance of chromosome structure (Chen et al. 1996, Thakur et al. 1997, McAllister et al. 1997, Scully et al. 1997a, Chen et al. 1998a, Welsch et al. 2000). BRCA1 and BRCA2 have major roles in maintaining genomic integrity e.g. through tumour suppression, transcription regulation, cell proliferation and differentiation, cell cycle control and participating in DNA repair. Recently it has been reported that BRCA2 is the Fanconi anaemia D1 protein and, as such, plays a key role in complex series of nuclear events that promote DNA crosslink repair (Howlett et al. 2002, Taniguchi and D'Andrea 2006). Also BRCA1 has been proposed to be involved in the Fanconi anaemia-BRCA pathway (Garcia-Higueira et al. 2001, D'Andrea and Grompe 2003). In their role as tumour suppressors BRCA1 and BRCA2 behave as caretakers, suppressing genome instability as they are essential for preserving the chromosome structure (Xu et al. 1999, Tirkkonen et al. 1997, Patel et al. 1998, Gretarsdottir et al. 1998).

*BRCA1* and *BRCA2* are considered as gatekeeper tumour suppressor genes that are involved in multiple tumour types, and as caretaker genes (Schutte et al. 1995, Thompson et al 1995, Holt et al. 1996, Wang et al. 2002, Kinzler and Vogelstein 1997). Gatekeepers directly regulate the growth of tumours by inhibiting growth or promoting death (Kinzler and Vogelstein 1997). *BRCA* genes have been shown to suppress breast cancer cell growth (Thompson et al. 1995, Holt et al. 1996, Wang et al. 2002). According to Knudson's double hit hypothesis the inactivation of a tumour suppressor (or gatekeeper) gene needs two mutations, usually one germ-line and one somatic, and leads to tumour development (Knudson 1971). Mutations in caretaker genes do not directly result in tumour formation but instead they cause genetic instability which leads to increased mutations in the genome and eventually to tumour formation (Kinzler and Vogelstein 1997). Tumours arising in women carrying a single germ-line mutant

*BRCA1* or *BRCA2* allele exhibit LOH at this locus, losing the wild-type allele and retaining the mutant copy of the gene, implying that *BRCA1* and *BRCA2* proteins function as tumour suppressors (Smith et al. 1992, Collins et al. 1995, Cornelis et al. 1995). How these proteins exert the tumour suppressor functions is incompletely understood.

*BRCA1* and *BRCA2* participate in the biological response to DNA damage, which involves the activation of cell cycle checkpoints and the recruitment of the machinery for DNA repair. Failure to activate these checkpoints or DNA repair following DNA damage manifests as increased sensitivity to genotoxic agents (Patel et al. 1998, Sharan et al. 1997, Xu et al. 2001, Scully et al. 1997b). In addition to the role in DNA repair after damage caused by exposure to exogenous agents, *BRCA1* and *BRCA2* proteins function in DNA damage response after endogenous damage that arise during processes such as DNA replication or transcription (Yarden and Brody 1999, Milner et al. 1997, Chapman et al. 1996, Scully et al. 1997c, Fuks et al. 1998, Bochar et al. 2000, Venkitaraman 2002, Welch et al. 2000, Scully and Livingston 2000). Although both are essential for error-free homologous recombination (HR) *BRCA1* and *BRCA2* have distinct roles in DSB repair. Disruption of *BRCA2* prevents efficient homologous recombination (HR) but single-stranded annealing (SSA) and non-homologous end-joining (NHEJ), which are more error-prone, are used (Xia et al. 2001, Moynahan et al. 2001, Tutt et al. 2001), whereas in *BRCA1* deficient cells SSA and HR appear to be decreased and NHEJ predominates as the mechanism for DSB repair (Moynahan et al. 1999). The major role of *BRCA2* is through control of the RAD51 recombinase (Chen et al. 1998b, Wong et al. 1997, Davies et al. 2001, Pellegrini et al. 2002, Yang et al. 2002a, Venkitaraman 2002, Pellegrini and Venkitaraman 2004), while *BRCA1* performs a distinct and more general function as a link between the sensing/signaling and effector components of the mammalian response to DNA damage, helping to ensure that the ensuing response is appropriate to the initiating lesion (Wang et al. 2000, Zhong et al. 1999, Cortez et al. 1999, Lee et al. 2000, Venkitaraman 2002). Chromosomal instability provoked by *BRCA* deficiency is the result of incorrect routing of DSB processing down inappropriate pathways, rather than the failure of repair per se (Venkitaraman 2002).

### 2.3.3 Cancer risks conferred by mutations in the *BRCA* genes

In a recent analysis of pooled pedigree data unselected for family history with breast or ovarian cancer from 22 studies, the average cumulative risks in *BRCA1* mutation carriers by age 70 years were 65 % for breast cancer and 39 % for ovarian cancer. The corresponding estimates for *BRCA2* were 45 % and 11 % (Antoniou et al. 2003). In high risk families the risks are somewhat higher, especially for breast cancer in *BRCA2* mutation carriers; ~85 % by age 70 and the risk for ovarian cancer ~27 % (Easton et al. 1995, Ford et al. 1998). The risk in *BRCA1*-mutation carriers by age 70 is ~70 % for breast cancer and ~40 % for ovarian cancer (Easton et al. 1995, Ford et al. 1998). The risk for subsequent breast or ovarian cancer is also high in breast cancer patients carrying *BRCA1* and *BRCA2* mutations (Easton et al. 1995, Breast Cancer Linkage Consortium 1999). The cumulative risk for male breast cancer in *BRCA2* mutation carriers has been estimated to be 3-6 % by age 70 years (Easton et al. 1997, Thompson and Easton 2001). Although the majority of families with male breast cancer are connected with germ-line *BRCA2* mutations, also *BRCA1* is suggested to contribute a larger fraction to male breast cancer than previously thought (Couch et al. 1996, Frank et al. 2002).

In addition to the high risks of breast and ovarian cancers, *BRCA2* mutations may be associated with increased risks of several other cancers. Statistically significant increases in risks were observed for prostate cancer, pancreatic cancer, stomach cancer, gallbladder and bile duct cancer and malignant melanoma (Breast Cancer Linkage Consortium 1999). The overall increases in risks for other than breast and ovarian cancer are quite small, except for prostate cancer in *BRCA2* mutation carriers where the cumulative risk has been estimated to be 10-34 % by age 80 years (Breast Cancer Linkage Consortium 1999, Thompson and Easton 2001). In *BRCA1* mutation carriers the overall increased risk of cancer at sites other than breast and ovary is small and is observed in women but generally not in men. *BRCA1* mutations may confer increased risks of other abdominal cancers in women and increased risks of pancreatic cancer in men and women (Thompson et al. 2002a).



### 2.3.3.1 Variation in cancer risks conferred by *BRCA* mutations

Depending on the *BRCA* mutation different risks of breast, ovarian, and other cancers, and possibly different probabilities for particular tumour types or tumour progression are conferred (Gayther et al. 1995, Gayther et al. 1997a, Scott et al. 2003). Different hypotheses have been introduced as the biological explanation for the variation in cancer risk by mutation location but none of them has been confirmed (Thompson and Easton 2001, Thompson et al. 2002b, Ware et al. 2006). The risk of cancer among the carriers may also differ considerably, due to e.g. chance, genetic, hormonal, dietary, and environmental modifying factors (Antoniou et al. 2000, Antoniou et al. 2002, Thompson and Easton 2004).

A lower risk of ovarian cancer in relation to breast cancer risk has been reported to be associated with mutations in the 3' end of the *BRCA1* gene, whereas the mutations in the central part of the gene confer a lower breast cancer risk than other mutations (Risch et al. 2001, Gayther et al. 1995, Thompson et al. 2002b). For *BRCA2*, mutations in the central part of the gene, in a region known as the ovarian cancer cluster region, OCCR (Figure 2), at nucleotides 3035-6629, appear to be associated with a lower breast cancer risk and higher ovarian cancer risk than other *BRCA2* mutations (Gayther et al. 1997a, Thompson and Easton 2001). Later, the OCCR was slightly modified on statistical grounds to comprise nucleotides 3059-4075 and 6503-6629 (Thompson and Easton 2001). Also splice-site mutations in both *BRCA1* and *BRCA2*, as well as substitutions in the *BRCA2* gene have been suggested to confer higher risks than substitutions in the *BRCA1* gene (Scott et al. 2003).

Possible genetic factors that modify the effect of *BRCA* gene mutations, besides *BRCA* genes themselves, include rare *HRAS1* alleles and a polymorphism in the *RAD51* gene (Phelan et al. 1996, Levy-Lahad et al. 2001, Wang et al. 2001a).

### 2.3.4 Clinical features of *BRCA* gene mutation-associated cancers

Several histopathological and clinical features differ among *BRCA1*, *BRCA2* and sporadic breast cancers. The pathological characteristics of *BRCA2*-associated breast tumours are less clear than for *BRCA1*, and overall their behaviour seems to be more similar to those in sporadic cases. (Table 2).

**Table 2.** Clinical features of *BRCA1* and *BRCA2* mutation-associated breast and ovarian cancers.

| <b><i>BRCA1</i>-associated</b>   | <b><i>BRCA2</i>-associated</b>  | <b>References</b>  |
|--|---|--|
| Breast carcinomas diagnosed at younger age   | Breast carcinomas diagnosed at younger age  | Hall et al. 1990, Easton et al. 1993, Miki et al. 1994, Wooster et al. 1994, Ford et al. 1994, Easton et al. 1995, Boyd et al. 2000, Risch et al. 2001       |
| Bilateral cancer more common   | Bilateral cancer more common  |  |
| Ovarian carcinomas diagnosed at younger age  |   |  |
| Often ER- and PR-  | Predominantly ER+ and PR+   | Johannsson et al. 1997, Loman et al. 1998, Lakhani et al. 2000   |
| High histological grade  | Evidence of higher average grade (effect weaker than for <i>BRCA1</i> )                     | Marcus et al. 1996, Breast Cancer Linkage Consortium 1997, Johannsson et al. 1997, Lakhani et al. 1998   |
| High mitotic count   |   |  |
| More often highly proliferating and aneuploid  |   |  |
| Overrepresentation of medullary or atypical medullary tumours                                      |   | Breast Cancer Linkage Consortium 1997  |
| Distinguished gene expression patterns in tumours  | Distinguished gene expression patterns in tumours   | Hedenfalk et al. 2001  |
| A trend towards worse survival in breast cancer found in some studies                              | Less data but not likely to be very different from sporadic<br>Worse survival also reported | Foulkes et al 1997, Verhoog et al. 1998, 2000, Loman et al. 2000, Stoppa-Lyonnet et al. 2000, Goode et al. 2002a, Robson et al. 2003, Brekelmans et al. 2006 |
| Ovarian carcinomas typically of serous histology, moderate to high grade and advanced stage        |   | Rubin et al. 1996, Boyd et al. 2000, Moslehi et al. 2000, Risch et al. 2001, Lakhani et al. 2004   |
| Mucinous subtype underrepresented  |   |  |
| A trend towards better survival in ovarian cancer (due to enhanced susceptibility to chemotherapy) | Little data available, better survival when combined analysis with <i>BRCA1</i>             | Rubin et al. 1996, Pharoah et al. 1999, Boyd et al. 2000, Ben David et al. 2002, Narod and Boyd 2002, Cass et al. 2003                                       |

Significantly different groups of genes are expressed by breast cancers with *BRCA1* mutations and breast cancers with *BRCA2* mutations, and sporadic breast tumours, and thus, these three groups can be distinguished from each other. This supports further the molecular difference between cancers with underlying germ-line mutations in *BRCA1* and *BRCA2* genes as well as cancers without such mutations. (Hedenfalk et al. 2001). Also ovarian tumours segregate into "*BRCA1*-like" and "*BRCA2*-like" subgroups according to the gene expression profiles. Mutations in *BRCA1* and *BRCA2* may lead to carcinogenesis through distinct molecular pathways that also appear to be involved in

sporadic cancers. (Jazaeri et al. 2002).

### 2.3.5 Mutation spectrum and founder mutations in the *BRCA* genes

More than 1 500 different mutations or polymorphisms have been identified in the *BRCA1* gene and almost 1 900 in the *BRCA2* gene (BIC, February 2006). The majority of mutations are frameshift mutations caused by small insertions or deletions, or alterations affecting the splice-site (BIC). Most of these mutations hamper the ability of the plausible tumour-suppressor or caretaker gene to block the development of cancer.

A mutational hotspot in the *BRCA1* or *BRCA2* gene has not been described but frequent founder mutations have been discovered in some populations, of which the Icelandic and Ashkenazi Jewish populations are the most studied. In Iceland the preponderance of a single *BRCA2* 999del5 mutation has been observed (Gudmundsson et al. 1996, Thorlacius et al. 1996). (Table 3). It is estimated to account for up to 76 % of families with multiple cases of female breast cancer and/or male breast cancer (Thorlacius et al. 1996). In addition to the 999del5 mutation in the *BRCA2* gene, only one rare mutation in the *BRCA1* gene has been identified in the Icelandic population (Bergthorsson et al. 1998).

**Table 3.** Predominant founder mutations in *BRCA1* and *BRCA2* genes.

| Population | Founder mutation      | Freq. in population | Breast cancer risk in female carriers | Reference  |
|------------|-----------------------|---------------------|---------------------------------------|--|
| Icelandic  | <i>BRCA2</i> 999del5  | 0.6 %<br>(1/170)    | 37 % at age 70                        | Johannesdottir et al. 1996,<br>Thorlacius et al. 1997,<br>Thorlacius et al. 1998 |
| Ashkenazim | <i>BRCA1</i> 185delAG | ~1 %                | 36 % (life-time)                      | Fodor et al. 1998  |
|            | <i>BRCA1</i> 5382insC | 0.25 %              |                                       | Fodor et al. 1998  |
|            | <i>BRCA2</i> 6174delT | ~1 %                | 36 % (life-time)                      | Fodor et al. 1998  |

The Ashkenazi Jews have three ancestral founder mutations, 185delAG and 5382insC in *BRCA1* (Simard et al. 1994) and 6174delT in *BRCA2* (Tavtigian et al. 1996) (Table 3). These mutations account for almost all the *BRCA1/2* mutations found in this population. The life-time risk for breast cancer among Ashkenazi Jewish carriers (regardless of age of onset or family history of cancer) of the *BRCA1* 185delAG or

*BRCA2* 6174delT mutations is approximately 3 times the overall risk for the general population (Fodor et al. 1998). Different estimations of higher risks of breast and ovarian cancer among the carriers of the Ashkenazi Jewish founder mutations in any population have been also published (Struewing et al. 1997, King et al. 2003, Antoniou et al. 2005).

In addition to the Ashkenazi Jews, the 5382insC mutation and another *BRCA1* mutation, 4153delA, are frequent among breast/ovarian cancer families of Eastern European, e.g. Russian, Hungarian, Latvian and Polish origin (Neuhausen et al. 1996, Gayther et al. 1997b, Ramus et al. 1997a, Csokay et al. 1999, Gorski et al. 2000). Patients being heterozygous carriers of two Ashkenazi Jewish founder mutations, 185delAG/6174delT and 5382insC/6174delT, have been detected (Ramus et al. 1997b, Friedman et al. 1998). The observations suggest that the phenotypic affects of double heterozygosity for *BRCA1* and *BRCA2* germ-line mutations are not cumulative (Friedman et al. 1998). A patient of a non-Jewish ancestry carrying both *BRCA1* and *BRCA2* mutations, G2508T/3295insA, as well as a patient homozygous for *BRCA1* 2800delAA mutation have also been observed (Liede et al. 1998, Boyd et al. 1995).

In the Dutch population large Alu-mediated deletions in the *BRCA1* gene are recurrent and are among the founder mutations (Petrij-Bosch et al. 1997, Puget et al. 1997, Swensen et al. 1997). Alu-mediated duplications of the *BRCA1* gene have been found (Puget et al. 1999), and a genomic deletion in the *BRCA2* gene in a Swedish breast/ovarian cancer family has also been published (Nordling et al. 1998).

Recurrent founder mutations are being identified in a growing number of populations as the data on *BRCA1* and *BRCA2* mutations accumulates. Founder effect is also evident in Finland (Vehmanen et al. 1997a, Huusko et al. 1998, Sarantaus et al. 2000).

### **2.3.6 *BRCA* mutations in Finland**

In Finland 15 different *BRCA1* and 11 different *BRCA2* mutations have been found among female and male breast cancer and ovarian cancer patients so far (Vehmanen et al. 1997a, Vehmanen et al. 1997b, Roth et al. 1998, Huusko et al. 1998, Sarantaus et al. 2000, Syrjäkoski et al. 2000, Sarantaus et al. 2001a, Sarantaus et al. 2001b,

Pääkkönen et al. 2001, Eerola et al. 2001, Vahteristo et al. 2001a, Syrjäkoski et al. 2004a). Fourteen of these, seven in each gene, are recurrent and are considered as founder mutations in Finland as they account for the majority of all detected mutations. Some of these mutations are unique to the Finns. The most recurrent mutation in *BRCA2* is the Icelandic founder mutation 999del5 (Table 4). Interestingly, the Icelandic and Finnish *BRCA2* 999del5 mutations could have an ancient common origin and in addition to Iceland and Finland the 999del5 mutation has not been reported in other Nordic countries (Barkardottir et al. 2001). Large genomic alterations have not been found in *BRCA1* or *BRCA2* gene in the Finnish population (Lahti-Domenici et al. 2001, Laurila et al. 2005, Karhu et al. 2006).

**Table 4.** The four most recurrent founder mutations in the *BRCA1* and *BRCA2* gene in Finland.

| Gene         | Mutation          |
|--------------|-------------------|
| <i>BRCA1</i> | 4216-2nt A→G      |
|              | 3604delA          |
|              | 3744delT/3745delT |
|              | 4446C→T           |
| <i>BRCA2</i> | 999del5           |
|              | 7708C→T           |
|              | 8555T→G           |
|              | 9346-2ntA→T       |

In the previous studies conducted in 88 Northern and 100 Southern Finnish families *BRCA1/2* mutations were found in 12.5 % and 21 % of the families, in 8-43 % depending on the strength of family history and being the highest in true high-risk families with both breast and ovarian cancers. *BRCA1* mutations were observed in 7 % and 10 % of studied breast/ovarian cancer families. The corresponding figures for *BRCA2* were 6 % and 11 %. (Huusko et al. 1998, Vehmanen et al. 1997a). The proportion of *BRCA2* mutations was slightly higher than *BRCA1* mutations. The highest (80 %) frequencies of *BRCA* mutations in Finland have been observed in families with both breast and ovarian cancer and early-onset breast cancer, while in families with later-onset breast cancer only, the mutation frequency has been the lowest, 1.5 % (Vahteristo et al. 2001a). Among 1035 unselected Finnish breast cancer cases *BRCA2* mutations were considerably more frequent with the frequency

of 1.4 %, than *BRCA1* mutations with the frequency of 0.4 % (Syrjäkoski et al. 2000). In Finnish ovarian carcinoma families *BRCA1/2* founder mutations were found in 26 % (Sarantaus et al. 2001a) and *BRCA2* mutations were slightly more frequent than *BRCA1* mutations. However, among unselected ovarian carcinoma patients *BRCA1/2* founder mutations account for an approximate 5 % of ovarian carcinomas (Sarantaus et al. 2001b) and *BRCA1* mutations are more frequent than *BRCA2* mutations.

### **2.3.7 Common polymorphisms and *BRCA* genes in sporadic breast/ovarian cancer**

The most common polymorphisms which cause amino acid substitutions in the *BRCA1* gene, Gln356Arg, Pro871Leu, Glu1038Gly and Ser1613Gly, do not make a significant contribution to breast and ovarian cancer risk (Dunning et al. 1997, Wenham et al. 2003, Auranen et al. 2005). However, the homozygous 356Arg genotype was suspected to be protective against breast cancer (Dunning et al. 1997). Cox and co-workers (2005a) reported a haplotype in the *BRCA1* gene that was associated with a small (18 %) increase in risk of sporadic breast cancer. However, the functional variants responsible for the association were unclear. They also reported no association with Gln356Arg and breast cancer. In another recent multiethnic study with breast cancer case-control material no evidence for significant associations between risk of sporadic breast cancer and common variation in *BRCA1* was found (Freedman et al. 2005).

The only common nonsynonymous SNP in the *BRCA2* gene is Arg372His (N372H). In a joint analysis of British, German and Finnish population the rare homozygotes (HH) were first associated with a 30 % increase in risk of breast cancer, with the strongest association observed in women <45 years of age (Healey et al. 2000a). This was also observed in an Australian study (Spurdle et al. 2002) but not in studies conducted in the US (Freedman et al. 2004, Cox et al. 2005b, Garcia-Closas et al. 2006). The rare homozygous genotype HH has also been shown to associate with increased risk of ovarian cancer in Australian and British population (Auranen et al. 2003), whereas in a US study no association was detected (Wenham et al. 2003). In the largest collaborative study conducted so far, no significant association was found

between increased risk of breast cancer and the HH homozygotes, and only a marginally statistically significant interaction of N372H with age (Breast Cancer Association Consortium 2006).

Somatic inactivating mutations in *BRCA1* and *BRCA2* gene are very rare and thus do not play a major role in the tumorigenesis of sporadic breast or ovarian cancer (Futreal et al. 1994, Meravjer et al. 1995, Miki et al. 1996, Lancaster et al. 1996, Teng et al. 1996, Weber et al. 1996). However, allelic deletions have been observed in sporadic breast and ovarian cancers in both genes, which imply that *BRCA* genes may play roles as tumour suppressor genes in sporadic breast and ovarian cancers also (Futreal et al. 1994, Takahashi et al. 1995, Beckmann et al. 1996, Gras et al. 2001, Yang-Feng et al. 1993). Also inactivation and reduction in expression levels of *BRCA1* in sporadic breast and ovarian cancers have been detected (Thompson et al. 1995, Taylor et al. 1998, Wilson et al. 1999, Esteller et al. 2000, Russell et al. 2000). The silencing of the *BRCA1* gene in primary breast and ovarian carcinomas occurs epigenetically by promoter hypermethylation, especially in the presence of LOH (Dobrovic and Simpfendorfer 1997, Baldwin et al. 2000, Rice et al. 1998, Bianco et al. 2000, Esteller et al. 2000). Hypermethylation of the *BRCA2* promoter region has not been observed in breast cancer (Collins et al. 1997) and only very rarely in ovarian cancers (Hilton et al. 2002). Instead, it has been suggested that *BRCA2* is silenced by overexpression of EMSY (Hughes-Davies et al. 2003). EMSY binds to exon 3 of *BRCA2* and thereby suppresses the transcriptional activity of *BRCA2* (Hughes-Davies et al. 2003).

Other possible mechanisms leading to low *BRCA1* expression are regulation of protein stability, allele specific expression, p53 or ID4 mediated expression, activation of the negative regulatory site in the first intron, loss of activation by transcription factors or activation of *NBR2* expression (reviewed in Mueller and Roskelley 2003). *BRCA1* protein has also been observed to be mislocalised to cytoplasm (Chen et al. 1995). However, the extent to which *BRCA1* and *BRCA2* contribute to the pathogenesis of sporadic breast and ovarian cancer remains unclear as *BRCA1* and *BRCA2* may affect through other more indirect mechanisms.

### 2.3.8 Other genes involved in familial breast cancer

In addition to the hereditary breast and ovarian cancer syndrome caused by mutations in the *BRCA1* and *BRCA2* genes, breast cancer occurs as a component of other inherited cancer predisposition syndromes. Germ-line mutations in these predisposing genes are however, very rarely seen in breast cancer patients without manifestations of other features of the syndromes. Thus, the contribution of these genes (*PTEN*, *TP53*, *ATM*, *LKB1*) to familial aggregation of site-specific breast cancer is marginal. Low-penetrance genes accounting for the familial aggregation of breast cancer have been also identified. These include *CHEK2* and a recently identified *PALB2*.

#### 2.3.8.1 *PTEN* and Cowden syndrome

Cowden syndrome (CS) is an autosomal dominant disorder, characterised by the development of hamartomas and benign tumours in multiple organs, e.g. skin, intestine, thyroid and breast, and an increased risk for breast, thyroid and endometrial carcinomas. Mutations in the tumour suppressor *PTEN* gene (also known as *MMAC1* or *TEP1*), on chromosome 10q22-q23, are present in over 80 % of Cowden syndrome families (Nelen et al. 1996, Liaw et al. 1997, Eng 2003). Truncating *PTEN* mutations in Cowden syndrome families are associated with cancer and cause a 25-50 % lifetime risk of breast cancer in women and the age at diagnosis of breast cancer in patients with CS is approximately 10 years younger than in sporadic cases (Starink et al. 1986, Longy and Lacombe 1996). The risk of male breast cancer might also be increased (Marsh et al. 1998, Fackenthal et al. 2001). No mutations in *PTEN* gene have been detected in breast cancer families without CS features (Tsou et al. 1997, Carroll et al. 1999, Lauge et al. 1999). In sporadic breast cancer *PTEN* mutations or polymorphisms seem not to play a significant role either (FitzGerald et al. 1998, de Jong et al. 2002, Haiman et al. 2006).

#### 2.3.8.2 *TP53* and Li-Fraumeni syndrome

The *TP53* gene is located on chromosome 17p13 (McBride et al. 1986). The p53 protein encoded by *TP53* is a 53-kD phosphoprotein that has four distinct domains: an N-terminal transactivation domain, a central DNA-binding domain, a tetramerization domain and a C-terminal regulatory domain (Soussi and May 1996). p53 is a



transcription factor and it is activated and stabilized in response to different forms of cellular stress, e.g. ionizing radiation, chemotherapeutic drugs and UV light, via post-translational modifications and protein-protein interactions, after which it transactivates or transrepresses the genes involved in cell-cycle control, DNA repair, apoptosis, and the formation of blood vessels (Vogelstein et al. 2000). p53 has been named as the "guardian of the genome" due to the multiple roles of the wild-type p53 protein in controlling the cell-cycle and proliferation (Lane 1992).

Germ-line mutations in the *TP53* gene cause Li-Fraumeni syndrome (Malkin et al. 1990). Li-Fraumeni syndrome (LFS) is a rare autosomal dominant disorder, characterised by an increased risk of soft tissue and osteosarcomas, leukaemias, brain tumours, adrenocortical carcinomas, and breast cancer (Li and Fraumeni 1969, Li et al. 1988). The risk of developing breast cancer before the age of 45 is 18-fold higher among LFS females compared to the general population (Garber et al. 1991). Germline *TP53* mutations have been found in approximately 70 % of LFS and 25 % of LFL (Li-Fraumeni-like) families (Varley et al. 1997, Eng et al. 1997). Germ-line mutations in the *TP53* gene have been estimated to account for less than 1 % of breast cancer but recent studies suggest a proportion higher than that, especially among cases with disease onset at the age 30 years or younger (Börresen et al. 1992, Sidransky et al. 1992, Lalloo et al. 2003). Germ-line mutations in *TP53* have also been found in Finnish breast cancer families meeting the LFS/LFL criteria (Huusko et al. 1999, Rapakko et al. 2001).

Germ-line mutations of *TP53* in familial, non-LFS or non-LFL, early-onset, or bilateral breast cancer cases are rare (Börresen et al. 1992, Lidereau and Soussi 1992, Sidransky et al. 1992, Balz et al. 2002, Ohayon et al. 2005, Lalloo et al. 2006). An Arg72Pro polymorphism is located in a proline-rich region of the *TP53* gene that is required for the growth suppression activity of p53 (Walker et al. 1996). After the initial finding of increased risk of breast cancer in homozygous carriers of the proline allele Arg72Pro has been examined in numerous studies for the possible role in breast cancer risk (Sjalander et al. 1996). The largest case-control studies so far have found no association between Arg72Pro and breast cancer risk (Wang-Gohrke et al. 2002, Tammiska et al. 2005, Breast Cancer Association Consortium 2006).

### 2.3.8.3 *ATM*

Ataxia-telangiectasia (A-T) is a rare autosomal recessive degenerative disorder caused by biallelic mutations of the *ATM* gene on chromosome 11q22-q23 (Savitsky et al. 1995). A-T is characterized by progressive cerebellar ataxia, a weakened immune system, hypersensitivity to ionizing radiation, highly increased susceptibility to malignancies, primarily of lymphoid origin, and distinctive dilated blood vessels in the eyes and skin. The ATM protein encoded by *ATM* is a large serine-threonine kinase that is activated in response to DNA damage by ionizing radiation (Canman et al. 1998, Kastan and Lim 2000). Upon activation ATM phosphorylates a wide array of downstream targets that regulate cell cycle checkpoints, apoptosis, and DNA repair (Kastan and Lim 2000, Bakkenist and Kastan 2003). ATM also regulates several tumour suppressors, including p53, BRCA1 and CHEK2 (Canman et al. 1998, Banin et al. 1998, Khanna et al. 1998, Cortez et al. 1999, Matsuoka et al. 1998).

Studies based on relatives of A-T patients have shown that heterozygous carriers of variant *ATM* are clinically unaffected but are shown to be at increased risk of cancer, especially the female carriers who have a two- to seven-fold increased risk of breast cancer (Swift et al. 1987, Pippard et al. 1988, Borresen et al. 1990, Swift et al. 1991, Easton 1994, Athma et al. 1996, Olsen et al. 2001, Thompson et al. 2005). In contrast, mutation screening of *ATM* in breast cancer case-controls sets outside A-T families have produced mixed results and have suggested allelic heterogeneity of *ATM* and that only a specific class of variants contribute to breast cancer risk in heterozygotes (FitzGerald et al. 1997, Teraoka et al. 2001, Gatti et al. 1999, Chenevix-Trench et al. 2002). Recently it has been shown in a UK study using familial breast cancer case-control population that A-T-causing biallelic *ATM* mutations are breast cancer susceptibility alleles in monoallelic carriers outside A-T families, and the combined *ATM* mutation prevalence and contribution to breast cancer incidence is similar to *CHEK2* 1100delC; both confer an estimated two-fold risk of breast cancer (Renwick et al. 2006). Association of some of the missense variants (7271T>G, Ser49Cys) has been observed in other large case-control studies also (Bernstein et al. 2006, Stredrick et al. 2006). In the Breast Cancer Association Consortium study the Ser49Cys was not significantly associated with overall breast cancer risk, but a modest association was not excluded either, and this

SNP increased the risk of PR positive breast cancer (Cox et al. 2007). Results from studies of association between common polymorphisms (SNPs) in *ATM* and risk of breast cancer have been controversial so far (Angele et al. 2003, Tamimi et al. 2004a, Lee et al. 2005a). In the Finnish population germ-line mutations in *ATM* have an apparent but overall limited contribution to familial and sporadic breast cancer predisposition outside A-T families (Pylkäs et al. 2007).

#### **2.3.8.4 Checkpoint kinase 2, *CHEK2* (*CHK2*)**

##### **2.3.8.4.1 *CHEK2* gene and protein function**

The cloning of the gene encoding the checkpoint kinase 2, *CHEK2*, was reported by five laboratories in 1998/1999 (Matsuoka et al. 1998, Blasina et al. 1999, Chaturvedi et al. 1999, Brown et al. 1999, Tominaga et al. 1999). The *CHEK2* (also known as *CHK2*) gene locates on chromosome 22q12.1 and spans 50 kb of genomic DNA. Fourteen exons encode a 543-amino-acid protein with 83 % identity to mouse *Chk2* and 34 % identity to *Drosophila* *Dmnk* (Matsuoka et al. 1998). *CHEK2* is also the human homolog of the *Saccharomyces cerevisiae* Rad53 and *Schizosaccharomyces pombe* Cds1 protein kinases that have crucial roles in controlling cell-cycle checkpoints after DNA damage and stalled replication (Matsuoka et al. 1998, Blasina et al. 1999, Chaturvedi et al. 1999, Brown et al. 1999). At the protein level the structure is dominated by several evolutionally conserved elements (Matsuoka et al. 1998, Bartek et al. 2001).

*CHEK2* is a G2 checkpoint protein kinase that plays a critical role in maintaining genomic stability (DNA repair) and coordinates cell cycle responses. In mammalian cells *CHEK2* is activated in response to replication blocks and DNA damage caused by UV light or ionising radiation (Matsuoka et al. 1998). *ATM* phosphorylates *CHEK2* which in turn phosphorylates other key cell cycle proteins, including *BRCA1* and *p53* (Matsuoka et al. 1998, Chehab et al. 2000, Lee et al. 2000, Shieh et al. 2000). In addition to *ATM*-dependent activation *CHEK2* is activated in an *ATM*-independent, *ATR*-dependent manner (Matsuoka et al. 1998, Chaturvedi et al. 1999, Blasina et al. 1999, Hirao et al. 2002). *CHEK2* has been also shown to regulate apoptosis through a *p53*-independent pathway by phosphorylating the promyelocytic leukemia protein, *PML*

(Yang et al. 2002b). Furthermore, CHEK2 has been reported to regulate also E2F1 transcription factor activity in response to the DNA-damaging agent etoposide (Stevens et al. 2003). On the other hand, E2F1 expression results in increase in CHEK2 protein levels and it may be essential for p53 activation and apoptosis induction (Rogoff et al. 2004). Ectopic expression of E2F1 induces the ATM-dependent phosphorylation of CHEK2 and stimulates the kinase activity of CHEK2 (Powers et al. 2004). NBS1 is also required for the induction of CHEK2 phosphorylation induced by E2F1 (Powers et al. 2004). Moreover, CHEK2 plays a critical role in the pro-apoptotic transcription factor p73 protein induction following DNA damage and modulation of E2F1 makes an important contribution (Urist et al. 2004). A recent study suggests that a mitotic checkpoint kinase TTK participates in the regulation of DNA damage by functioning upstream of CHEK2 and phosphorylating it (Wei JH et al. 2005). DNA-PK is also suggested to be involved in activation of CHEK2 in response to DNA damage (Li and Stern 2005).

Checkpoint mechanisms and DNA-repair machinery are needed for temporary halting of cell-cycle progression in proliferating cells, in order to provide time for repair and prevent fixation and propagation of harmful mutations (Hartwell and Weinert 1989, Elledge 1996, Weinert 1998, Walworth 2000, Bartek et al. 2001). When DNA damage occurs, activated CHEK2 targets several cell cycle modulators. These events ultimately lead to cell cycle arrest, DNA repair, or programmed cell death. Failures in the checkpoint machinery can result in developmental malformations, embryonic lethality or the accumulation of mutations that could potentially lead to genetic disease, including cancer (Hartwell 1992, Hartwell and Kastan 1994, Weinert 1997, Dasika et al. 1999). CHEK2 mutations have been detected in tumours and it has been suggested that constitutive activation of ATM-CHEK2 pathway occurs at pre-invasive stages of major types of human tumours, including colon, breast and lung carcinomas (Bartkova et al. 2005).

#### **2.3.8.4.2 *CHEK2* germ-line mutations and risk of breast cancer**

The first germ-line *CHEK2* mutations were reported as early as in 1999 in both sporadic and hereditary human cancers. Germ-line mutations in *CHEK2*, including 1100delC,

were detected in Li-Fraumeni syndrome patients without *TP53* mutations (Bell et al. 1999). The findings suggested that *CHEK2* is a tumour suppressor gene conferring predisposition to sarcoma, breast cancer and brain tumours, and provided a link between the central role of p53 inactivation in human cancer and the well-defined G2 checkpoint in yeast (Bell et al. 1999). The 1100delC mutation was also detected in Finnish families suggestive of Li-Fraumeni syndrome (Vahteristo et al. 2001b). Recent studies have confirmed that *CHEK2* is mutated in a subset of familial breast cancers (CHEK2-Breast Cancer Consortium 2002, Vahteristo et al. 2002, Meijers-Heijboer et al. 2003, Oldenburg et al. 2003). Both truncated and missense mutants of *CHEK2* have lost their kinase activity i.e. the ability to interact with, and efficiently phosphorylate, substrates such as Cdc25A and p53 (Falck et al. 2001a, Falck et al. 2001b, Wu et al. 2001). It has been suggested that the range of cancers associated with mutations of the *CHEK2* may be much greater than previously thought and they may increase the risk of cancer in many different organs (Cybulski et al. 2004). However, this was not confirmed in a large multicenter study (Thompson et al. 2006). Significant association to ovarian cancer has not been detected either (Baysal et al. 2004).

The *CHEK2* 1100delC was shown to be a relatively common low-penetrance allele conferring susceptibility to breast cancer (CHEK2-Breast Cancer Consortium 2002, Vahteristo et al. 2002, CHEK2 Breast Cancer Case-Control Consortium 2004) (Table 5). The 1100delC variant allele has been estimated to result in an approximately 2-fold increase of breast cancer risk in women and a 10-fold increase of risk in men. In *BRCA1* or *BRCA2* mutation carriers the 1100delC allele did not confer increase in cancer risk, possibly because of functional redundancy (CHEK2-Breast Cancer Consortium 2002, CHEK2 Breast Cancer Case-Control Consortium 2004). The estimated 10-fold risk for male breast cancer or the high carrier frequency among males (CHEK2-Breast Cancer Consortium 2002) has not been confirmed in other studies and is likely to be smaller (Syrjäkoski et al. 2004b, Neuhausen et al. 2004, Sodha et al. 2004, Ohayon et al. 2004).

Vahteristo and co-workers (2002) suggested that *CHEK2* acts as a low-penetrance tumour-suppressor gene in breast cancer and that it makes a significant contribution to familial clustering of breast cancer - including families with only two affected relatives, which are more common than families that include larger numbers of affected women.

**Table 5.** Observed prevalence of the *CHEK2* 1100delC mutation.

| Observed frequency of <i>CHEK2</i> 1100delC mutation in |              |   |   |   |  |
|---|--------------|---|---|---|--|
| Population  | Controls     | Breast cancer cases in <i>BRCA1/2</i> neg. families | Breast cancer cases with family history | Breast cancer cases in families with <i>BRCA1/2</i> mutations | Population-based breast cancer cases                 |
| UK, Netherlands, North America                          | 1.1 %        | 5.1 %   |   | 1 %   | 1.4 %  |
| Finland   | 1.4 %        | 5.5 %   | 3.1 %                                   |   | 2.0 %  |
| UK, Netherlands, Finland, Germany, Australia            | 0.7 %        |   |   |   | 1.9 %  |
| Netherlands   | 2.8 %        | 11.4 %  |   |   |  |
| Germany   | 0.9 %        |   | 1.5 %, 1.4 %                            |   | 0.45 %, 0.8 %  |
| Spain   | 0 %          | 0 %   |   |   |  |
| Czech and Slovak  |              |   | 0.2 %                                   |   | 0.4 %  |
| Italy   | 0 %          | 0 %   | 0.11 %                                  |   |  |
| Australia   |              |   | 0.6 %                                   |   |  |
| US  | 0.4 %, 0.5 % |   | 1.1 %                                   |   | 1.1 %, 1.2 %   |
| Ashkenazi Jews  | 0.3 %        |   | 1.0 %                                   |   |  |
|   |              |   |   |   | References   |
|   |              |   |   |   | CHEK2-Breast Cancer Consortium 2002                  |
|   |              |   |   |   | Valteristo et al. 2002                               |
|   |              |   |   |   | CHEK2 Breast Cancer Case-Control Consortium 2004     |
|   |              |   |   |   | Oldenburg et al. 2003                                |
|   |              |   |   |   | Dufault et al. 2004, Rashid et al. 2005              |
|   |              |   |   |   | Osorio et al. 2004                                   |
|   |              |   |   |   | Kleibl et al. 2005                                   |
|   |              |   |   |   | Caligo et al. 2004                                   |
|   |              |   |   |   | Jekimovs et al. 2005                                 |
|   |              |   |   |   | Mateus-Pereira et al. 2004, Friedrichsen et al. 2004 |
|   |              |   |   |   | Offit et al. 2003                                    |

In the collaborative analysis the prevalence of 1100delC was somewhat greater in women reporting a first-degree relative with breast cancer and also some evidence was seen that the prevalence in cases decreased with increasing age at diagnosis (CHEK2 Breast Cancer Case-Control Consortium 2004). 1100delC has also been observed to associate with bilateral breast cancer (Vahteristo et al. 2002). A strong association of the *CHEK2* 1100delC with a putative hereditary breast and colorectal cancer phenotype (HBOC) was detected, suggesting a subtype of familial breast cancer (Meijers-Heijboer et al. 2003). It was also suggested that *CHEK2* 1100delC mutation acts in synergy with another as-yet-unknown cancer-susceptibility gene or genes (Meijers-Heijboer et al. 2003, Oldenburg et al. 2003, Johnson et al. 2005).

Variation in 1100delC carrier frequencies in different populations have been observed (CHEK2 Breast Cancer Case-Control Consortium 2004), and it is either absent or occurs in similar frequencies among breast cancer cases and controls in many populations. Thus, it is evident that the contribution of *CHEK2* mutations varies by ethnic groups and from country to country. (Table 5). Other variants in *CHEK2* have not been detected to make a similar contribution to familial breast cancer susceptibility as the 1100delC does (Allinen et al. 2001, Schutte et al. 2003, Dufault et al. 2004, Friedrichsen et al. 2004). Though, significant associations have been found in some study populations where the 1100delC is rare. (Table 6). Common polymorphisms in *CHEK2* are unlikely to confer an increased risk of breast cancer (Kuschel et al. 2003).

**Table 6.** Observed variants in *CHEK2*, other than 1100delC.

| Variant                    | Population                            | Association with                         | References   |
|----------------------------|---------------------------------------|--|--|
| Ser428Phe                  | Ashkenzi Jews                         | familial breast cancer                   | Shaag et al. 2005  |
| Ile157Thr                  | Byelorussian                          | familial breast cancer                   | Bogdanova et al. 2005  |
| Ile157Thr                  | Finnish, Polish, German, Byelorussian | sporadic breast cancer                   | Kilpivaara et al. 2004, Cybulski et al. 2004, 2006a, Bogdanova et al. 2005 |
| deletion of exons 9 and 10 | Czech and Slovak, Polish              | 2-fold increase in risk of breast cancer | Walsh et al. 2006, Cybulski et al. 2007                                    |

The carriers of *CHEK2* 1100delC appear to develop ER and PR receptor positive tumours more frequently than non-carriers (de Bock et al. 2004, Schmidt et al. 2007), and an association with lobular histology and a *CHEK2* variant has been suggested (Huzarski et al. 2005, Cybulski et al. 2006). Tumours from 1100delC carriers also tend

to be of higher grade than those of non-carriers (Kilpivaara et al. 2005). Tumours with reduced CHEK2 expression had more often larger primary tumour size compared to tumours with normal expression (Kilpivaara et al. 2005) and CHEK2 expression is often reduced in tumours from patients with germ-line 1100delC (Vahteristo et al. 2002). Otherwise a characteristic *CHEK2* signature in breast tumours has not been discovered yet (Cybulski et al. 2006, Schmidt et al. 2007). A worse disease-free survival has also been suggested for the mutation carriers (de Bock et al. 2004, Schmidt et al. 2007).

#### **2.3.8.5 *LKB1* and Peutz-Jeghers syndrome**

Peutz-Jeghers syndrome is a rare autosomal dominant disorder characterised by mucocutaneous pigmentation and gastrointestinal polyposis. In addition to an elevated risk of gastrointestinal cancers, an increased risk of cancers at other sites, such as breast, ovary, uterus, cervix, lung, and testis, has been described. (Giardiello et al. 1987, Tomlinson and Houlston 1997, Giardiello et al. 2000). The cumulative risk of breast cancer by age 70 years has been reported to be as high as 45 % (95 % CI=27-68) (Hearle et al. 2006). The Peutz-Jeghers syndrome is caused by germ-line mutations in the serine-threonine kinase gene *LKB1* (= *STK11*) located on chromosome 19p13 (Hemminki et al. 1998, Jenne et al. 1998). Germ-line mutations in the *LKB1* gene have not been reported in other than Peutz-Jeghers families, and this gene seems to play a role in breast cancer susceptibility only in patients with Peutz-Jeghers syndrome (Chen and Lindblom 2000, de Jong et al. 2002).

#### **2.3.8.6 *PALB2* and Fanconi anemia**

Fanconi anemia (FA) is a rare autosomal recessive syndrome characterized by congenital abnormalities, progressive bone-marrow failure, and cancer susceptibility. (D'Andrea and Grompe 2003). FA is a genetically heterogeneous condition and thus far the disease includes 13 subtypes (complementation groups), 12 of which have been attributed to mutations in distinct genes, suggesting that their gene products function in a common pathway (Taniguchi and D'Andrea 2006, Reid et al. 2007, Patel et al. 2007). The most recently discovered of FA genes is *PALB2* (Reid et al. 2007).



*PALB2* (partner and localizer of *BRCA2*) gene locates on chromosome 16p12 and encodes a recently discovered protein that interacts with *BRCA2* (Xia et al. 2006). *PALB2* colocalizes with *BRCA2* in nuclear foci, promotes its localization and stability in key nuclear structures (e.g. chromatin and nuclear matrix), and enables its recombinational repair and checkpoint functions (Xia et al. 2006). Biallelic *PALB2* germ-line mutations are responsible for a subset of Fanconi anemia (FA-N) cases characterized by a phenotype similar to that caused by biallelic *BRCA2* mutations (Reid et al. 2007, Xia et al. 2007). Monoallelic *PALB2* mutations were also found in individuals with breast cancer from familial breast cancer pedigrees that were negative for *BRCA1* and *BRCA2* mutations. *PALB2* mutations showed an incomplete segregation in affected relatives and were estimated to confer a 2.3-fold increase in breast cancer risk and it was suggested that the risks of breast cancer associated with *PALB2* mutations may be age dependent. The breast cancer population-attributable fraction of *PALB2* mutations in this UK population was estimated to be 0.23 % and the percentage of familial relative risk due to *PALB2* to be 0.24 %. (Rahman et al. 2007). A truncating mutation, being a possible founder mutation, in *PALB2* was also detected in familial and sporadic breast cancer cases and in one prostate cancer family in the Finnish population (Erkko et al. 2007). *PALB2* is the first gene that encodes for a protein that interacts with *BRCA2* and confers an increased risk of breast cancer when mutated (Xia et al. 2006, Rahman et al. 2007).

The phenotypes associated with biallelic *BRCA2* and *PALB2* mutations are markedly similar to each other and differ from the other ten known FA genes (Reid et al. 2007, Xia et al. 2007). Biallelic mutations in *BRIP1* cause one of the FA subtypes and heterozygous mutations in *BRIP1* confer an elevated risk of breast cancer (Levrin et al. 2005, Seal et al. 2006). However, it is suggested that breast cancer susceptibility is associated with only a subset of Fanconi anemia genes which is consistent with the negative results of mutational screens of other FA genes (Seal et al. 2003). The currently known FA genes that associate with breast cancer, *BRCA2*, *PALB2* and *BRIP1*, are not part of the FA core complex and are the only known FA genes that act downstream of *FANCD2* (Rahman et al. 2007).

## **2.4 Sporadic breast cancer and low-penetrance susceptibility genes**

Familial aggregation of breast cancer is thought to account for 5-10 % of all cases of breast cancer (Claus et al. 1996). Currently known high-risk and high-penetrance susceptibility genes confer approximately 20 % of this (Thompson and Easton 2004). However, in twin studies heritable factors are observed even in one fourth of breast cancer cases (Lichtenstein et al. 2000). It is assumed that several common, low to moderate penetrance genes are present in the general population and account also for the residual familial aggregation of breast cancer (Pharoah et al. 1997, Antoniou et al. 2002). The disease causing alleles of high-penetrance genes are relatively rare in the general population, whereas the low-penetrance disease-associated alleles are common. High-penetrance genes predispose to both earlier onset of the disease and multifocal tumours. In contrast to these familial cases, sporadic breast cancers caused by low-penetrance genes usually appear unilaterally and have relatively late age at diagnosis. Low-penetrance genes are likely to interact with environmental and life-style factors as well as with other genes to cause cancer. (Rebbeck 1999). As a result of this etiologic complexity, malignant diseases that do not follow a simple Mendelian pattern of inheritance in a substantial number of families have remained refractory to mapping by linkage. (Le Marchand 2005). It has been observed that subtle effects of some genetic variants may be magnified and only become detectable in the presence of certain exposures (Low et al. 2006). The identification of such low or moderate-penetrance susceptibility alleles is an extensive task. Utilising the candidate-gene approach, polymorphisms in several genes have been tested in breast cancer case-control association studies. In these studies the genes have been selected in pathways of known or suspected breast cancer-associated factors, such as genes affecting the oestrogen levels and metabolism, DNA repair or tumour growth. The genes studied for breast cancer association in the Eastern Finnish population (KBCP) are listed in Table 7.

**Table 7.** Genes studied for association with breast cancer risk in the Eastern Finnish (KBEP) population.

| Gene                          | Polymorphism              | Association   | Significant association when stratified  | OR (95% CI)   | Reference              |
|-------------------------------|---------------------------|---|--|---|------------------------|
| <b>Oestrogen biosynthesis</b> |                           |   |  |   |                        |
| <i>CYP17</i>                  | 5'UTR T1931C              | No overall association                                    |  | 0.92 (0.69-1.22) for C allele containing genotypes                | Mitrunen et al. 2000   |
| <b>Oestrogen metabolism</b>   |                           |   |  |   |                        |
| <i>CYP1A1</i>                 | Ile462Val                 | No overall association                                    |  | 0.76 (0.50-1.14) for Ile/Val and Val/Val combined                 | Sillanpää et al. 2007  |
|                               | Thr461Asn                 | No overall association                                    |  | 1.67 (0.50-5.48) for Thr/Asn                                      | Sillanpää et al. 2007  |
| <i>CYP1B1</i>                 | Leu432Val                 | No overall association                                    | Increased risk among smokers of 1-9 cigarettes/day                                       | 2.6 (1.07-6.46) for Leu/Val<br>5.1 (1.30-19.89) for Val/Val       | Sillanpää et al. 2007  |
| <i>COMT</i>                   | Val108/158Met             | No overall association                                    |  | 0.81 (0.58-1.13) for Val/met and Met/Met combined                 | Mitrunen et al. 2001c  |
| <i>SULT1A1</i>                | Arg213His                 | No overall association                                    | Decreased risk in premenopausal women with at least 3 pregnancies and one His allele     | 0.23 (CI=0.09-0.63)   | Sillanpää et al. 2005b |
| <i>GSTM1</i>                  | Allele deletion           | No overall association                                    | Increased risk among postmenopausal GSTM1 null genotypes                                 | 1.49 (1.03-2.15)  | Mitrunen et al. 2001a  |
| <i>GSTM3</i>                  | 3 bp deletion in intron 6 | No overall association                                    |  | 1.00 (0.73-1.37) for variant allele containing genotypes          | Mitrunen et al. 2001a  |
| <i>GSTP1</i>                  | Ile105Val                 | No overall association                                    |  | 0.88 (0.66-1.18) for Ile/Val<br>0.57 (0.31-1.04) for Val/Val      | Mitrunen et al. 2001a  |
| <i>GSTT1</i>                  | Allele deletion           | No overall association                                    |  | 1.18 (0.8-1.76) for null genotypes                                | Mitrunen et al. 2001a  |
| <i>MnSOD</i>                  | Val16Ala                  | Ala allele carriers at increased risk compared to Val/Val |  | <b>1.5 (1.1-2.0)</b>  | Mitrunen et al. 2001b  |
| <b>Carcinogen metabolism</b>  |                           |   |  |   |                        |
| <i>NAT2</i>                   | C282T<br>T341C            |   | Increased risk associated with slow acetylator genotypes 282T/282T, 341C/341C, 282T/341C | 1.32 (1.01-1.73)<br>Risk higher among women smoking 5< pack-years | Sillanpää et al. 2005a |
| <i>ADH1B</i>                  | 3'UTR A>G                 | No overall association                                    |  | 0.79 (0.59-1.05) for AG<br>1.38 (0.71-2.69) for GG                | Cox et al. 2007        |

Table 7. (continued)

| Gene  | Polymorphism | Association  | Significant association when stratified                       | OR (95% CI)   | Reference  |
|---|--------------|--|---|---|--|
| <b>DNA repair</b>                               |              |  |   |   |  |
| <i>BRCA2</i>                                    | Asn372His    | No overall association                             |   | 1.04 (0.83-1.30) for Asn/His<br>1.07 (0.61-1.87) for His/His                  | Healey et al. 2000a  |
| <i>XRCC1</i>                                    | Arg280His    | No overall association                             |   | 1.15 (0.80-1.66) for Arg/His and His/His combined                             | Metsla et al. 2005   |
|   | Arg399Gln    | No overall association                             | Increased risk among ever smokers and at least one Gln allele | 2.33 (1.30-4.19)<br>Risk higher among women smoking 5< pack-years             | Metsla et al. 2005   |
| <i>XPB</i>                                      | Lys751Gln    | No overall association                             | Increased risk among ever smokers and Gln/Gln genotype        | 2.52 (1.27-5.03)<br>Risk higher among women smoking 5< pack-years             | Metsla et al. 2005   |
| <i>ATM</i>                                      | Ser49Cys     | No overall association                             |   |   | Cox et al. 2007  |
| <b>Hormone- and growth factor-related genes</b> |              |  |   |   |  |
| <i>IGFBP3</i>                                   | -202C>A      | No overall association                             |   | 1.20 (0.89-1.62) for CA<br>0.91 (0.61-1.36) for AA<br><b>0.73 (0.54-0.98)</b> | Cox et al. 2007<br>Sillanpää et al. 2004                     |
| <i>VDR</i>                                      | Apal         | a allele carriers at decreased risk compared to AA |   |   | Sillanpää et al. 2004  |
|   | TaqI         | No overall association                             |   | 0.68 (0.41-1.12) for Tt and tt combined                                       | Sillanpää et al. 2004  |
| <i>TGFBI</i>                                    | Leu10Pro     | No overall association                             |   | 0.91 (0.71-1.17) for Leu/Pro<br>1.14 (0.68-1.91) for Pro/Pro                  | Dunning et al. 2003  |
| <b>Cell cycle control</b>                       |              |  |   |   |  |
| <i>STK15</i>                                    | Phe3Ile      | No overall association                             |   | 1.02 (0.78-1.34) for Phe/Ile<br>0.61 (0.35-1.07) for Ile/Ile                  | Breast Cancer Association Consortium 2006<br>Cox et al. 2007 |
| <i>CDKN1A</i>                                   | Ser31Arg     | No overall association                             |   | 1.27 (0.81-2.00) for Ser/Arg<br>3.15 (0.33-30.43) for Arg/Arg                 | Cox et al. 2007  |
| <b>Other genes</b>                              |              |  |   |   |  |
| <i>CASP8</i>                                    | Asp302His    | No overall association                             |   | 0.82 (0.57-1.16) for Asp/His<br>6.53 (0.34-126.92) for His/His                | Cox et al. 2007  |
| <i>ICAM5</i>                                    | Val301Ile    | No overall association                             |   | 0.94 (0.60-1.48) for Val/Ile<br>1.04 (0.66-1.63) for Ile/Ile                  | Cox et al. 2007  |
| <i>NUMA1</i>                                    | Ala794Gly    | No overall association                             |   | 0.69 (0.36-1.33) for Ala/Gly and Gly/Gly combined                             | Cox et al. 2007  |


|                             |                            |   |                                    |                    |
|-----------------------------|----------------------------|---|------------------------------------|--------------------|
| <i>FGFR2</i>                | intron 2 G>A<br>rs2981582  | GA and AA at increased risk<br>compared to GG | <b>1.36 (1.11-1.66)</b> per allele | Easton et al. 2007 |
| <i>LSP1</i>                 | intron 10 T>C<br>rs3817198 | No overall association                        | 1.03 (0.82-1.30) per allele        | Easton et al. 2007 |
| <i>H19</i>                  | C>T rs2107425              | No overall association                        | 1.03 (0.83-1.28) per allele        | Easton et al. 2007 |
| <i>TNRC9/<br/>LOC643714</i> | intron 1 C>T<br>rs8051542  | CT and TT at increased risk<br>compared to CC | <b>1.37 (1.11-1.67)</b> per allele | Easton et al. 2007 |
|                             | intron 1 A>G<br>rs12443621 | AG and GG at increased risk<br>compared to AA | <b>1.26 (1.03-1.55)</b> per allele | Easton et al. 2007 |
|                             | C>T<br>rs380362            | No overall association                        | 1.03 (0.83-1.27) per allele        | Easton et al. 2007 |

### 2.4.1 Steroid hormone biosynthesis, signalling and metabolism genes

Most of the known risk factors for breast cancer relate to the increased, prolonged or life-time oestrogen exposure. Oestrogens also undergo extensive oxidative metabolism via several enzymes, yielding catechol oestrogens (CE) and hydroxyestrone, some of which are even carcinogenic. Therefore the genes encoding enzymes involved in oestrogen and other steroid hormone biosynthesis, signalling and metabolism are strong candidates for low-penetrance genetic risk factors for breast cancer, as allelic variants in these enzymes may affect the hormone levels.

#### 2.4.1.1 Steroid hormone biosynthesis genes

Oestrogen biosynthesis involves series of enzymatic steps from cholesterol to C-19 androgens and C-18 oestrogens (Figure 3, Table 8). Table 8 shows the genes involved in steroid hormone biosynthesis and their observed association with breast cancer risk.



**Figure 3.** Oestrogen biosynthesis. CYP11A, CYP17 and CYP19 are particularly important. See Table 8 for details.

#### 2.4.1.2 Steroid hormone signalling genes

The sex hormones bind to specific receptors and form complexes that in turn bind to sequences in the promoters of downstream, hormone-responsive genes and thus the sex hormones control the activation of responsive genes. Therefore the steroid hormone receptor genes *ESR1* (oestrogen receptor alpha) (Walter et al. 1985, Menasce et al.

Table 8. Genes involved in steroid hormone biosynthesis and signalling.

| Gene, locus   | Encoded Protein  | Function   | Studied polymorphisms    | Association with breast cancer   | References   |
|---|--|--|--------------------------|--|--|
| <b>Steroid hormone biosynthesis</b> (Payne 1990, Omura and Morohashi 1995, Kristensen and Borresen-Dale 2000, Thompson and Ambrosone 2000). |  |  |                          |  |  |
| <i>CYP11A</i> , 15q23-q24   | <i>CYP11A</i>  | catalyses the rate-limiting step in the biosynthesis of all steroids, the cleavage of the side chain of cholesterol to form pregnenolone and progesterone              |                          |  |  |
| <i>CYP17</i> , 10q24.3  | <i>CYP17</i> , 17 $\alpha$ hydroxylase/C17-20 lyase            | catalyzes the rate-limiting step in ovarian and adrenal biosynthesis pathways for androstenedione, immediate precursor of testosterone                                 | 5'UTR T>C                | Controversial results  | Feigelson et al. 1997, Dunning et al. 1998, Mitrunen et al. 2000, Einarsdottir et al. 2006 |
| <i>CYP19</i> , 15q21.1  | <i>CYP19</i> , aromatase/estrogen synthetase                   | aromatizes testosterone and androstenedione to E2 (estradiol) and E1 (estrone)   | a tetranucleotide repeat | seems to be a risk factor  | Kristensen et al. 1998, Healey et al. 2000b, Dunning et al. 1999a                          |
| <i>HSD3B2</i> , 1p13.1  | 3 $\beta$ HSD, 3 $\beta$ hydroxysteroid dehydrogenase          | converts pregnenolone to progesterone  |                          |  |  |
| <i>EDH17B2</i> , 16q24.1  | 17 $\beta$ HSD, 17 $\beta$ hydroxysteroid dehydrogenase type 2 | catalyzes the interconversions of the less biologically active hormones androstenedione and estrone to their more active analogues testosterone and estradiol          | several                  | no association   | Mannermaa et al. 1994  |
| <b>Steroid hormone signalling</b>   |  |  |                          |  |  |
| <i>ESR1</i> , <i>ESR2</i> , 6q25.2, 14q   | ER, oestrogen receptor alpha and oestrogen receptor beta       | binds oestrogen  | some                     | no overall effect shown  | Southey et al. 1998, Cai et al. 2003, Gold et al. 2004, Zheng et al. 2003                  |
| <i>PGR</i> , 11q22  | PR, progesterone receptor                                      | binds progesterone   | Val660Leu                | haplotype may be associated with increased risk                                | Pooley et al. 2006, Breast Cancer Association Consortium 2006                              |
| <i>AR</i> , Xq11-q12  | AR, androgen receptor  | binds androgen   | CAG repeat               | controversial results  | Lillie et al. 2003, Dunning et al. 1999b   |
| <i>SHBG</i> , 17p13-p12   | SHBG, sex hormone-binding globulin                             | binds the circulating sex hormones and so modulates the bioavailability of these hormones to target tissues, regulates the steroid-signalling system in target tissues | Asp327Asn                | associates with increased SHBG levels and reduced risk in postmenopausal women | Dunning et al. 2004, Cui et al. 2005   |

1993), *ESR2* (oestrogen receptor beta) (Enmark et al. 1997), *PGR* (progesterone receptor) (Law et al. 1987, Misrahi et al. 1987) and *AR* (androgen receptor) (Chang et al. 1988, Lubahn et al. 1988), as well as *SHBG* (sex hormone-binding globulin) are candidates for breast cancer susceptibility genes. (Table 8). Women with higher circulating levels of SHBG have been detected to have a lower risk of breast cancer (Endogenous Hormones and Breast Cancer Collaborative Group 2002). Also association between higher testosterone levels and increase in breast cancer risk in postmenopausal women has been observed (Endogenous Hormones and Breast Cancer Collaborative Group 2002).


#### **2.4.1.3 Steroid hormone metabolism genes**

Oestrogens are extensively metabolised by a number of oxidative and conjugative reactions that can lead to their deactivation and subsequent elimination. Alternatively, oxidation and conjugation reactions of oestrogens may generate metabolites that have distinct biologic activities, including altered hormonal properties; genotoxicity; and/or chemotherapeutic properties. In the metabolism of oestrogen are several enzymes/genes involved (Figure 4). (Zhu and Conney 1998, Jefcoate et al. 2000, Thompson and Ambrosone 2000). (Table 9).

Oxidative metabolism of oestrogens is mediated by the same CYPs that metabolise therapeutic agents and xenobiotics. Oestrogens are metabolised via two major pathways, i.e. hydroxylation either at the A-ring or the D-ring, leading to formation of catechol oestrogens (CE) and 16 $\alpha$ -hydroxyestrone, respectively. (Martucci and Fishman 1993, Lippert et al. 2000). Oestrogen conjugation is a major route of oestrogen metabolism. Hydroxylated oestrogens can be inactivated by *O*-methylation catalysed by COMT (Zhu and Conney 1998, Cavalieri et al. 1997). Sulfate conjugation of oestrogens and CEs is catalyzed by several members of a superfamily of cytosolic sulfotransferase enzymes, SULTs (Glatt et al. 2001). Sulfation and methylation detoxifies CEs and conjugated oestrogens are not appreciable ligands for the ERs, thus, they do not promote ER-mediated activity (Zhu and Conney 1998). CEs are the major metabolites of oestrogen, and they are intermediates for the generation of the reactive semiquinones



(CE-SQ) and quinones (CE-Q), which in turn have the ability to damage DNA and protein through redox cycling and generation of reactive oxygen species (ROS), and thus have been implicated as potential initiators of tumour formation (Cavalieri et al. 1997, Zhu and Conney 1998, Liehr 1997). Superoxide dismutases (SODs) comprise a family of metalloenzymes that catalyse the conversion of superoxide anion into hydrogen peroxide and molecular oxygen and thus protect cells from damage induced by free radicals (Winterbourn 1993). CE-Q may be conjugated with glutathione (GSH) catalysed by cytosolic glutathione S-transferases, GSTs, or reduced to CE by quinone reductase NQO1 (Cavalieri et al. 1997). (Figure 4, Table 9).



**Figure 4.** Oestrogen metabolism.

In addition to oestrogen metabolism, GSTs are a family of enzymes responsible for the metabolism of a broad range of xenobiotics and carcinogens (Mannervik 1985). GSTs catalyse the conjugation of glutathione (GSH) to a variety of endogenous and exogenous electrophiles, increasing their water solubility and excretability (Ketterer 1988). Also NQO1 is involved in metabolising numerous endogenous and

**Table 9.** Genes involved in steroid hormone metabolism.

| Gene, locus                | Encoded Protein                                   | Function   | Studied polymorphisms                                | Association with breast cancer   | References   |
|----------------------------|---|--|--|--|--|
| <i>CYP1A1</i> , 15q22-q24  | CYP1A1, cytochrome P450 subfamily I polypeptide 1 | catalyses 2-hydroxylation of oestradiol to CE, also a major role in the metabolism of several potent carcinogens e.g. in activating cigarette smoke constituents                             | some   | association among smokers or post-menopausal women, no overall association | Ambrosone et al. 1995, Ishibe et al. 1998, Bartsch et al. 2000, Basham et al. 2001 |
| <i>CYP1B1</i> , 2p22-p21   | CYP1B1, cytochrome P450 subfamily I polypeptide 1 | main CYP450 enzyme responsible for the 4-hydroxylation of oestradiol to CE   | several  | no consistent association, Leu432Val may modify risk                       | Sutter et al. 1994, Tang et al. 1996, Sillanpää et al. 2007                        |
| <i>CYP1A2</i> , 15q22-qter | CYP1A2, cytochrome P450 subfamily I polypeptide 2 |  | some   | controversial results  | Le Marchand et al. 2005, Long et al. 2006  |
| <i>COMT</i> , 22q11.2      | COMT, catechol-O-methyltransferase                | conjugation and inactivation of catechol oestrogens by transfer of a methyl group  | Val>Met change associating with decreased activity   | no overall association   | Thompson et al. 1998, Huang et al. 1999, Gaudet et al. 2006                        |
| <i>SULT1A1</i> , 16p12.1   | SULT1A1, cytosolic sulfotransferase 1A1           | catalyses sulfate conjugation of oestrogens and CEs, also involved in detoxification of xenobiotic compounds and in the metabolism and activation of pro-carcinogens from e.g. tobacco smoke | Arg213His, associates with decreased enzyme activity | may modify risk, no overall association                                    | Seth et al. 2000, Zheng et al. 2001, Sillanpää et al. 2005b                        |
| <i>SULT1E1</i> , 4q13.1    | SULT1E1, cytosolic sulfotransferase 1E1           | catalyses sulfate conjugation of oestrogens and CEs, exhibits the highest affinity for oestrogens among SULTs  | one study, 959G>A                                    | decreased risk among Koreans   | Choi et al. 2005   |
| <i>GSTM1</i> , 1p31        | GSTM1, mu-class glutathione S-transferase (GST) 1 | catalyses glutathione (GSH) conjugation of catechol oestrogen quinones (CE-Q)  | gene deletion  | null allele associated with increased risk among postmenopausal women      | Helzlsouer et al. 1998, Dunning et al. 1999a, Mirunen et al. 2001a                 |
| <i>GSTM3</i> , 1p13        | GSTM3, mu-class GST 3                             | catalyses GSH conjugation of CE-Q  | 3bp del in intron 6, only one study                  | no overall association   | Mirunen et al. 2001a   |
| <i>GSTP1</i> , 11q13       | GSTP1, pi family GST 1                            | catalyses GSH conjugation of CE-Q, predominant GST in the breast tissue and is expressed consistently in both normal and tumour breast tissue  | Ile105Val, Ala114Val                                 | overall risk not conclusive  | Dunning et al. 1999a, Vogl et al. 2004, Egan et al. 2004, Mirunen et al. 2001a     |
| <i>GSTT1</i> , 22q11       | GSTT1, theta family GST 1                         | catalyses GSH conjugation of CE-Q  | gene deletion  | no overall association   | Dunning et al. 1999a, Vogl et al. 2004, Egan et al. 2004, Mirunen et al. 2001a     |

|                          |   |   |  |  |  |
|--------------------------|---|---|--|--|--|
| <i>SOD2</i> ,<br>6q25    | MnSOD, mitochondrial manganese superoxide dismutase | catalyses the conversion of superoxide anion into hydrogen peroxide and molecular oxygen, thus protects cells from damage induced by free radicals, proposed to be a tumour suppressor                                  | Val16Ala                                   | overall association not conclusive                               | Ambrosone et al. 1999, Mitrinen et al. 2001b, Bergman et al. 2005, Egan et al. 2003, Tamimi et al. 2004b, Cai et al. 2004, Cebrian et al. 2006, Breast Cancer Association Consortium 2006, Cox et al. 2007 |
| <i>NQO1</i> ,<br>16q22.1 | NQO1, NAD(P)H: quinone oxidoreductase               | reduces reactive CE-Q back to hydroxy-CE, plays an important role in protecting tissues against the mutagenic, carcinogenic and cytotoxic effects, activates e.g. environmental procarcinogens present in tobacco smoke | Pro187Ser, leading to nonfunctional enzyme | associated with increased risk in Czech and Austrian populations | Menzel et al. 2004   |

**Table 10.** Carcinogen and xenobiotics metabolism genes studied for breast cancer association.

| Gene, locus                              | Encoded Protein  | Function   | Studied polymorphisms   | Association with breast cancer          | References  |
|--|--|--|---|---|---|
| <i>NAT1</i> ,<br><i>NAT2</i> ,<br>8p22   | NAT1, N-acetyltransferase 1<br>NAT2, N-acetyltransferase 2         | participates in the metabolism of numerous pharmaceutical drugs and carcinogens (heterocyclic amines) found in tobacco smoke and diet  | slow/intermediate/fast acetylator status vs. breast cancer risk | no overall association, may modify risk | Ambrosone et al. 1996, Alberg et al. 2004, Zheng et al. 1999, Hunter et al. 1997, Millikan et al. 1998, Dunning et al. 1999a, Lee et al. 2003, Sillanpää et al. 2005a |
| <i>MPO</i> ,<br>17q23                    | MPO, Myeloperoxidase   | antimicrobial enzyme catalyzing hydrogen peroxide-dependent oxidation of chloride to generate hypochlorous acid, transforms xenobiotics, such as PAHs and aromatic amines to highly reactive intermediates | -463 G>A, associates with reduced expression                    | reduced risk observed                   | Ahn et al. 2004, Lin et al. 2005  |
| <i>ADH1B</i> ,<br><i>ADH1C</i> ,<br>4q22 | ADH1B, alcohol dehydrogenase 1B<br>ADH1C, alcohol dehydrogenase 1C | ADH plays a rate-limiting role in the metabolic pathway for most human ethanol oxidation   | 3'UTR A>G in <i>ADH1B</i><br>Ile350Val in <i>ADH1C</i>          | no overall association                  | Freudenheim et al. 1999, Hines et al. 2000, Terry et al. 2006, Breast Cancer Association Consortium 2006, Cox et al. 2007   |

environmental quinones (Ernster and Navazio 1958, Grzeschik 1980, Povey et al. 1980, Jaiswal et al. 1988, Chen et al. 1991). It is a 2-electron reductase that detoxifies quinones derived from the oxidation of phenolic metabolites of benzene (Ernster 1967, Lind et al. 1982). In the metabolism of xenobiotics NQO1 may act as either a detoxification or activation enzyme, depending on the substrate.

#### **2.4.2 Carcinogen and xenobiotics metabolism**

Several enzymes are involved in the detoxification of xenobiotic compounds found in e.g. tobacco smoke and cooked meat. Some of these enzymes are involved in the steroid hormone metabolism also (introduced above, Table 9). The detoxifying enzymes convert carcinogenic compounds in more soluble and more readily excreted form and therefore reduce cancer risk. Some of the more soluble compounds, however, are even more carcinogenic. Thus, genetic changes that increase or decrease the activity of the detoxifying enzymes may increase the amount of reactive carcinogens and increase the risk of cancer. Such enzymes are candidates for breast cancer susceptibility and objects for genetic case-control association studies.

*NAT1* and *NAT2* genes are highly polymorphic and several different allelic variants exist that determine the NAT1/NAT2-related biotransformation capacity resulting in slow, intermediate or fast/rapid acetylator phenotype (Evans et al. 1960, Deguchi et al. 1990, Vatsis et al. 1991, Blum et al. 1991, Hickman et al. 1992, Lin et al. 1993, Deguchi 1992, Butcher et al. 1998, Hughes et al. 1998). A difference in the acetylation rate can alter the proportion of specific metabolites that are formed and genotoxicity by substrates of NAT can be affected by the amount of acetylation. Relationship between acetylator phenotype and DNA damage by chemicals that undergo N-acetylation has been found and the acetylation polymorphism may be a factor in susceptibility to toxicity and perhaps carcinogenicity of these chemicals (McQueen et al. 1982). The association of NAT1 and NAT2 acetylator phenotypes and breast cancer has been studied but the findings have been very inconsistent. (Table 10).

MPO generates ROS endogenously and is linked to cancers through DNA damage and the production of carcinogens (Klebanoff 1980, Van Rensburg et al. 1992, Petruska

et al. 1992). MPO is present in breast secretions and is thought to be involved in the chemical/metabolic induction of breast cancer (Josephy 1996, Williams et al. 2000). Also, there are indications that steroid hormones regulate *MPO* expression (Bekesi et al. 2001). Thus, it is plausible to hypothesize that MPO could play a role in breast carcinogenesis, and that factors that would increase lifetime exposures to estrogens, such as age at menarche, could modify this relationship (Ahn et al. 2004). (Table 10).

Alcohol consumption may increase the risk of breast cancer by increasing the exposure to carcinogenic metabolites (Feron et al. 1991) or by influencing the levels of reproductive steroid hormones that play a critical role in breast carcinogenesis (Ginsburg et al. 1995). Alcohol dehydrogenases (ADHs) are enzymes involved in the metabolism of ethanol to acetaldehyde. ADH plays a rate-limiting role in the metabolic pathway for most human ethanol oxidation. (Bosron and Li 1986). Polymorphisms in ADH genes have been studied for breast cancer association (Table 10).

### **2.4.3 Other hormone- and growth factor-related genes**

Breast cancer risk is known to be strongly related to endogenous hormone exposure and the genes responsive to such hormones are plausible candidates for being susceptibility genes. Also genes encoding for many growth factors involved in cellular growth, cell proliferation and differentiation, and angiogenesis are potent candidates for breast cancer risk associated factors, as all these pathways are affected in tumour growth. Several of these genes have been studied and some of them have shown association with breast cancer risk.

#### **2.4.3.1 Insulin-like growth factor family**

Insulin-like growth factor family includes the insulin-like growth factors (IGFs) I and II (IGF-I, IGF-II), six different IGF binding proteins (IGFBPs), and the receptors IGF-IR and IGF-IIR. In addition, several IGFBP proteases hydrolyze IGFBPs, resulting in release of the bound IGFs, and therefore indirectly regulate the action of IGFs. IGF family members interact with several molecules that are involved in cancer

development and progression, including the sex steroid hormones, tumour suppressors, and other growth factors. (Yu and Rohan 2000).

The IGF axis has been shown to play a role in mammary carcinogenesis (Hadsell and Bonnette 2000). IGF-I and IGFBP3 concentrations vary greatly between individuals, part of which is attributable to genetic effects (Juul et al. 1994, Juul et al. 1995, Harrela et al. 1996), and might affect the distribution of cancer risk in a population. High serum levels of IGF-I have been shown to be associated with breast cancer risk among premenopausal women (Hankinson et al. 1998, Renehan et al. 2004). The results on IGFBP3 levels and association with breast cancer risk are uncertain, possibly due to methodological reasons. A meta-analysis suggests an increase in risk with higher IGFBP3 levels (Renehan et al. 2004). However, the biological role of IGFBP3 supports the observation that higher circulating levels are protective (Al-Zahrani et al. 2006). Polymorphisms in both genes associate with the circulating levels of their respective proteins (Jernström et al. 2001, Schernhammer et al. 2003, Canzian et al. 2006, Al-Zahrani et al. 2006). (Table 11).

Growth hormone (GH) is the main stimulator of IGF-I and IGFBP3 production and genetic variants affecting the release or biological action of GH might have an effect on cancer risk. (Table 11).

#### **2.4.3.2 Vitamin D receptor, *VDR***

Vitamin D receptor, *VDR*, is a ligand-activated transcription factor interacting with many pathways, including p21, fibronectin and retinoic signaling, and its actions are involved in e.g. calcium metabolism, cell proliferation and differentiation, and immune function (Uitterlinden et al. 2004). Several polymorphisms (TaqI, ApaI, BsmI and poly-A which are all in LD with each other, and the only known protein polymorphism in the *VDR* gene, FokI) have been studied in the vitamin D receptor *VDR* gene but conclusive evidence on the association with increased risk of breast cancer has not been shown (Arai et al. 1997, Dunning et al. 1999b, Colin et al. 2000, Whitfield et al. 2001, de Jong et al. 2002, Uitterlinden et al. 2004, Chen et al. 2005). (Table 11).

**Table 11.** Other hormone- and growth factor-related genes studied for breast cancer association.

| <b>Gene, locus</b>                                       | <b>Encoded Protein</b>                               | <b>Function</b>  | <b>Studied polymorphisms</b>  | <b>Association with breast cancer</b> | <b>References</b>  |
|--|--|--|---|---------------------------------------|--|
| <i>IGF-I</i> , 12q22-q24                                 | IGF-I, insulin-like growth factor I                  | important role in regulating cell proliferation, differentiation, apoptosis and transformation, has characteristics of a circulating hormone and a tissue growth factor                                    | variants that associate with raised circulating IGF-I levels          | increased risk                        | Wen et al. 2005, Al-Zahrani et al. 2006  |
| <i>IGF-II</i> , 11p15                                    | IGF-II, insulin-like growth factor II                | important role in regulating cell proliferation, differentiation, apoptosis and transformation   |   |                                       |  |
| <i>IGFBP3</i> , 7p12-p14                                 | IGFBP3, insulin-like growth factor binding protein 3 | binds IGF-I and IGF-II in the circulation, regulates the biological activity of IGFs and thus might protect against cancer development, may have IGF-independent functions, may act as a growth stimulator | variants that associate with lower circulating IGFBP3 levels, -202C>A | decreased risk                        | Al-Zahrani et al. 2006, Breast Cancer Association Consortium 2006, Cox et al. 2007 |
| <i>GHI</i> , 17q23.3                                     | GH, growth hormone                                   | main stimulator of IGF-I and IGFBP3 production   |   |                                       |  |
| <i>GHRH</i> , 20q11.23                                   | GH releasing hormone                                 | stimulates GH release  |   |                                       |  |
| <i>GHRHR</i> , 7p15-p14                                  | GHRHR, GH releasing hormone receptor                 |  | -261C>T   | decreased risk                        | Wagner et al. 2006a  |
| <i>GHSR</i> , 3q26-31                                    | GH secretagogue receptor                             |  |   |                                       |  |
| <i>GHRL</i> , 3p26-2p5                                   | GH secretagogue receptor ligand                      | stimulates GH release  |   |                                       |  |
| <i>SST</i> , 3q28  | SST, somatostatin                                    | inhibits GH release  | several   | controversial results                 | Canzian et al. 2005, Wagner et al. 2006a   |
| <i>SSTR1-5</i> , 14q13, 17q24, 22q13.1, 20p11.2, 16p13.3 | SSTR1-5, somatostatin receptors 1-5                  | enhance or inhibit the synthesis and release of GH, and a pituitary-specific transcription factor, POU1F1  | promoter polymorphisms in <i>SSTR2</i>                                | controversial results                 | Canzian et al. 2005, Wagner et al. 2006a   |
| <i>VDR</i> , 12q11                                       | VDR, Vitamin D receptor                              | ligand-activated transcription factor interacting with many pathways   | TaqI, ApaI, BsmI, poly-A  | not associated                        | Dunning et al. 1999b, de Jong et al. 2002  |
| <i>VEGF</i> , 6p21                                       | VEGF, vascular endothelial growth factor             | regulates angiogenesis, mediates vascular permeability   | FokI  | increased risk                        | Chen et al. 2005   |
| <i>TGFBI</i> , 19q13                                     | TGFBI, transforming growth factor $\beta$            | roles in regulation of cell growth, differentiation and migration, in e.g. mammary gland development and function  | several   | controversial results                 | Krippl et al. 2003, Jin et al. 2005, Kataoka et al. 2006, Jacobs et al. 2006       |
|  |  |  | Leu10Pro  | increased risk                        | Cox et al. 2007  |

#### **2.4.3.3 Vascular endothelial growth factor, *VEGF***

Formation of new blood vessels is needed for tumour growth, invasiveness and metastasis, and vascular endothelial growth factor (VEGF) is the most important regulator of angiogenesis (Ferrara et al. 1997, Ferrara et al. 2003). VEGF is an endothelial cell-specific mitogen, an angiogenic inducer and a mediator of vascular permeability (Ferrara et al. 1997). In breast cancer tissue the mRNA expression of *VEGF* is increased compared to adjacent normal tissue (Yoshiji et al. 1996). Polymorphisms in the *VEGF* have been suggested to be associated with an increased and decreased risk of breast cancer. (Table 11).

#### **2.4.3.4 Transforming growth factor beta, *TGFBI***

TGF $\beta$  is a polypeptide cytokine that has a role in regulating cell growth, differentiation and migration. It regulates e.g. normal mammary gland development and function by activating the TGF $\beta$  signalling pathway. There is a dual-role of action in which the TGF $\beta$  signalling suppresses tumour initiation but can also promote tumour progression and metastasis when antiproliferative effect of the TGF $\beta$  signaling pathway has been overridden by other oncogenic mutations. (Bierie and Moses 2006, Derynck et al. 2001).

A Leu10Pro polymorphism resides in the signal peptide sequence of TGF $\beta$ 1 and the peptide with Pro at residue 10 causes a 2.8-fold increase in secretion compared with the Leu form (Dunning et al. 2003). In a large collaborative study some evidence for association with *TGFBI* Leu10Pro and breast cancer risk was shown (Breast Cancer Association Consortium 2006). This was further confirmed in an even larger sample material (Cox et al. 2007). The analysis of Leu10Pro variant showed a significant dose-dependent association of the Pro allele with increased risk of invasive breast cancer (per-allele OR 1.08, CI=1.04-1.11,  $P(\text{trend})=2.8 \times 10^{-5}$ ), and it was estimated to account for approximately 0.2 % of the excess familial risk of breast cancer in populations of European ancestry (Cox et al. 2007). (Table 11).



#### 2.4.4 Genes involved in double-stranded DNA break repair

Double-strand breaks are frequently induced by e.g. certain chemicals, ionizing radiation, and mechanical stress, and they are particularly difficult to repair compared with other types of DNA damage. Inability to correctly repair DSBs can cause chromosomal loss, translocations and deletions that may lead to activation of proto-oncogenes, loss of function of tumour suppressor genes or global genomic instability and thus, lead to tumorigenesis. (Khanna and Jackson 2001). The genes which normally participate in the error-free repair of breaks in double-stranded DNA predispose to breast and other cancers when inactivated and common variants of DSB-involved genes have been implicated in epidemiological studies of breast cancer risk.

Two general forms of DSB repair are found within eukaryotic cells: non-homologous end-joining (NHEJ) and homologous recombination (HR). Several types of homologous repair exist; gene conversion, break-induced replication (BIR) and single-strand annealing (SSA). Similarly, there are several alternative end-joining mechanisms. (Haber 2000). NHEJ repairs broken DNA ends by joining the broken strands together and it does not need an undamaged partner and does not rely on extensive homologies between the two recombining ends. Consequently, NHEJ is often prone to error, and small deletions often occur at the site of fusion. HR uses homologous chromosomal or sister chromatid DNA as template for synthesis of new error-free DNA. In HR, the DNA ends are first resected in the 5' to 3' direction by nucleases. The resulting 3' single-stranded tails then invade the DNA double helix of a homologous, undamaged partner molecule, and are extended by the action of DNA polymerase, which copies information from the partner. Following branch migration, the resulting DNA crossovers are resolved to yield two intact DNA molecules. In contrast to NHEJ the HR is a high fidelity process. (Khanna and Jackson 2001).

In humans, several genes are involved in the NHEJ (*KU70*, *XRCC5*, *DNA-PK*, *XRCC4*, *LIG4*) and HR (*ATM*, *BRCA1*, *BRCA2*, *RAD51*, *RAD52*, *RAD54*, *RAD51B*, *RAD51C*, *RAD51D*, *RPA*, *XRCC2*, *XRCC3*, *DMC1*, *ZNF350=ZBRK1*, *BRIP1=BACH1*) and proteins encoded by some of them (*NBS1*, *MRE11*, *RAD50*) have functions in both pathways (van Gent et al. 2001). Polymorphisms have been studied in several of these genes and some studies have reported associations, but others not (Table 12).

**Table 12.** Genes involved in double-stranded DNA break repair that have been studied for breast cancer association.

| Gene, locus                           | Contribution to breast cancer risk   | References  |
|---------------------------------------|--|---|
| <i>XRCC2</i> , 7q36.1                 | suggested to modify risk, significant main effect not confirmed  | Kuschel et al. 2002, Rafii et al. 2002, Breast Cancer Association Consortium 2006 |
| <i>LIG4</i> , 13q22-q34               | suggested to modify risk, significant main effect not confirmed  | Kuschel et al. 2002, Breast Cancer Association Consortium 2006                    |
| <i>XRCC3</i> , 14q32.3                | suggested association with weak risk, significant main effect not confirmed  | Kuschel et al. 2002, Garcia-Closas et al. 2006, Breast Cancer Association         |
| <i>XRCC4</i> , <i>Ku70</i> , 5q13-q14 | suggested association among Taiwanese  | Fu et al. 2003  |
| <i>RAD52</i> , 12p13-p12.2            | suggested association among Koreans  | Lee et al. 2005b  |
| <i>RAD51</i> , 15q15.1                | modifies risk among <i>BRCA</i> mutation carriers, no role in sporadic breast cancer   | Levy-Lahad et al. 2001, Wang et al. 2001a   |
| <i>RAD51D</i> , 17q11                 | suggested association with increased risk  | Rodriguez-Lopez et al. 2004   |
| <i>ZNF350</i>                         | suggested association in familial high-risk subgroup   | Garcia-Glosas et al. 2006   |
| <i>BRIP1</i> ( <i>BACH1</i> ), 17q22  | truncating mutations are low-penetrance breast cancer susceptibility alleles in familial breast cancer, estimated breast cancer-attributed fraction of 0.20 %, SNPs not associated | Sigurdson et al. 2004, Seal et al. 2006, Song et al. 2007                         |
| <i>FANCD2</i> , 3p25.3                | suggested association with increased risk  | Barroso et al. 2006   |
| <i>NBS1</i> , 8q21                    | suggested association with increased risk, mutations seem to contribute to risk  | Kuschel et al. 2002, Lu et al. 2006, Steffen et al. 2006                          |
| <i>RAD50</i> , 5q31                   | suggested low-penetrance risk allele among Northern Finnish breast cancer cases  | Heikkinen et al. 2006   |
| <i>BARD1</i> , 2q34                   | see text for details   |   |

#### 2.4.4.1 *BARD1*

The BRCA1-associated ring domain 1 (BARD1) protein was discovered as a binding partner of BRCA1 (Wu et al. 1996). BARD1 and BRCA1 have several features in common: similar protein structure, the embryonic lethality of their respective knockout mice and induction of genetic instability when depleted from cells (Irminger-Finger and Jefford 2006). BARD1 and BRCA1 form a functional heterodimer through the binding of their RING-finger domains (Wu et al. 1996, Meza et al. 1998). The interaction is required for several of the cellular and tumour suppressor functions of BRCA1. BRCA1-BARD1 heterodimer has been described as the functional unit in DSB repair (Jasin 2002). BARD1 and BRCA1 also associate with other repair proteins, such as the MSH2-MSH6 complex, and this interaction indicates a role in DNA mismatch repair (Wang et al. 2001b). Several reports have also demonstrated BRCA1-independent functions of BARD1, primarily in apoptosis, and BRCA1-independent increased

expression of BARD1 during mitosis, indicating that BARD1 on its own might have crucial functions. Its cancer-associated activities seem to centre on two major pathways: as an essential component of a ubiquitin ligase involved in DNA repair and cell-cycle regulation, and p53-mediated apoptosis. (Irminger-Finger and Jefford 2006).

*BARD1* gene locates on chromosome 2q34 (Wu et al. 1996, Thai et al. 1998). Germline *BARD1* mutations have been found in few cases of sporadic and familial, non-*BRCA1/2*, breast and ovarian cancer patients and some tumours, but they account for only a small fraction of cases of familial breast cancer overall (Thai et al. 1998, Ghimenti et al. 2002, Ishitobi et al. 2003, Karppinen et al. 2004, Sauer and Andrulis 2005). One missense alteration, Cys557Ser has been suggested to associate with a slightly increased risk of familial breast cancer (Ghimenti et al. 2002, Karppinen et al. 2006). Increase in risk was also observed in Icelandic case-control study, and the risk among *BRCA2* 999del5 mutation carriers was further increased if they also carried the *BARD1* 557Ser (Stacey et al. 2006). In the Chinese population other *BARD1* variant alleles have been suggested to be protective (Huo et al. 2006).

## 2.4.5 Genes involved in single-stranded DNA damage repair mechanisms

### 2.4.5.1 Base-excision repair

Base-excision repair (BER) is an important DNA repair pathway responsible for the repair of single-stranded base damage resulting from X-rays, oxygen radicals, methylation, deamination, and hydroxylation. BER acts on small lesions and involves release of the damaged base and removal of up to a few neighboring nucleotides. BER is particularly relevant for preventing mutagenesis. BER pathway includes several proteins and the genes encoding them have numerous genetic variants. These genes include *XRCC1*, *ADPRT* (PARP-1), *PCNA*, *FEN1*, *OGG1*, *MUTYH*, *APEX1=APE1*, *LIG3*, *LIG1*, *NTH1*, and *POLB* (DNA polymerase  $\beta$ ). *XRCC1* protein interacts with DNA polymerase  $\beta$ , DNA ligase III, PARP-1, and APE1 in the final ligation stage of the BER DNA repair pathway, and of the BER genes *XRCC1* is the most studied for breast cancer association. (Hoeijmakers 2001, Goode et al. 2002b). (Table 13).

**Table 13.** Polymorphisms studied for breast cancer association in genes involved in base-excision repair (BER) and nucleotide-excision repair (NER).

| Gene, locus                  | Polymorphism     | Contribution to breast cancer  | References  |
|------------------------------|------------------|--|---|
| <b>BER</b>                   |                  |  |   |
| <i>XRCC1</i> , 19q13.2       | Arg399Gln        | suggested association among African-Americans and Asian populations, suggested to modify risk among smokers or with family history, no overall association confirmed | Duell et al. 2001, Kim et al. 2002, Figueiredo et al. 2004, Metsola et al. 2005, Chacko et al. 2005, Zhang et al. 2006, Breast Cancer Association Consortium 2006 |
| <b>NER</b>                   |                  |  |   |
| <i>XPD (ERCC2)</i> , 19q13.2 | Asp312Asn        | association observed in some but not confirmed in larger studies   | Justenhoven et al. 2004, Försti et al. 2004, Kuschel et al. 2005, Breast Cancer Association Consortium 2006   |
|                              | Lys751Gln        | suggested to associate with risk and modify risk among smokers (Finns)   | Justenhoven et al. 2004, Terry et al. 2004, Metsola et al. 2005   |
|                              | Asp1140His       | marginal association among Finns   | Kumar et al. 2003   |
| <i>ERCC1</i> , 19q13.2       | 8092A>C          | suggested increased risk among Koreans   | Lee et al. 2005b  |
| <i>ERCC4 (XPF)</i> , 16p13.3 | intronic variant | suggested protective   | Milne et al. 2006   |

#### 2.4.5.2 Nucleotide-excision repair

Nucleotide-excision repair (NER) deals with the wide class of bulky helix-distorting lesions that interfere with base pairing and generally obstruct transcription and normal replication. Most NER lesions arise from exogenous sources and affect only one of the DNA strands (single-stranded). Two NER subpathways exist: global genome NER (GG-NER) and transcription-coupled repair (TCR). (Hoeijmakers 2001). Numerous proteins are involved in NER and genes encoding them include *XPA-XPG*, *ERCC1*, *RAD23A*, *RAD23B*, *CETN2*, *RPA1-3*, *GTF2H1-4*, *DDB1*, *DDB2*, *CSA*, *CSB*, *XAB2*, *BRCA1*, and *BRCA2* (Friedberg 2001). Some of these genes are candidates and have been studied for breast cancer risk association. *XPD* (= *ERCC2*) encodes a protein that is an integral member of the basal transcription factor BTF2-TFIIH complex, and it is the most studied of the NER genes for breast cancer association. (Table 13).

#### 2.4.5.3 Mismatch repair

Mismatch repair (MMR) removes nucleotides mispaired by DNA polymerases and insertion/deletion loops (ranging from one to ten or more bases) that result from slippage during replication of repetitive sequences or during recombination. Defects in this system dramatically increase mutation rates, fuelling the process of oncogenesis.

Genes involved in MMR are *hMLH1*, *hMSH2*, *hMSH3*, *hMSH6*, *hPMS1* and *hPMS2*. (Hoeijmakers 2001). So far polymorphism only in *hMLH1* has been associated with increased risk of breast cancer in Korean population (Lee et al. 2005b).

#### 2.4.6 Genes involved in cell cycle control

Cell cycle control is critical for normal growth and differentiation. Defects in checkpoints contribute to genetic instability and cancer. Genes encoding proteins that are involved in controlling cell cycle progression, include e.g. *TP53*, *CHEK2*, *ATM*, *RB1*, *CDKN1A*, *CDKN1B*, *CDKN2A*, *CCND1* and *STK15*.

The *STK15* (*AURORA2/AURKA/BTAK/ARK1*) gene on chromosome 20q13 encodes a serine/threonine kinase that acts as a key regulator of mitotic chromosome segregation, and has a role in the etiology of cancer (Shindo et al. 1998, Hirota et al. 2003, Meraldi et al. 2004). A Phe31Ile polymorphism in *STK15* has been reported to be a low-penetrance susceptibility allele affecting multiple cancer types (Ewart-Toland et al. 2005) and it has been studied for breast cancer association but the results are inconsistent. (Table 14).

**Table 14.** Breast cancer association of genes involved in cell cycle control.

| Gene, locus             | Variation           | Contribution to breast cancer  | References  |
|-------------------------|---------------------|--|---|
| <i>STK15</i> ,<br>20q13 | Phe31Ile            | differing results, suggested increased risk in women < 40 years of age   | Sun et al. 2004, Ewart-Toland et al. 2005, Cox et al. 2006, Fletcher et al. 2006, Breast Cancer Association Consortium 2006 |
| <i>CDKN2A</i> ,<br>9p21 | inherited mutations | predispose to breast cancer in the context of a syndrome of melanoma, pancreatic cancer, and breast cancer in Swedish families | Borg et al. 2000  |
|                         | Ala148Thr           | suggested association in Polish population   | Debniak et al. 2005   |

*CDKN2A* gene on chromosome 9p21 is a tumour suppressor gene that is involved in susceptibility of malignant melanoma and familial pancreatic carcinoma (Kamb et al. 1994, Monzon et al. 1998, Whelan et al. 1995, Sherr 2001). The p16 protein encoded by *CDKN2A* is a cyclin dependent kinase inhibitor that suppresses cell proliferation (Serrano et al. 1993). p16 is a cell-cycle regulator that inhibits the activities of cdk4 and cdk6, two protein kinases that in turn phosphorylate the retinoblastoma protein which is

necessary for progression through the G1 cell cycle checkpoint (Serrano et al. 1993, Sherr 2001 ). (Table 14).

#### 2.4.7 Proto-oncogenes

Proto-oncogenes are involved in the regulation of normal cell growth and differentiation. Mutations in proto-oncogenes lead to disturbances in the cell cycle and can result in abnormal growth of proliferation. Germ-line mutations in proto-oncogenes have not been found in syndromes conferring an increased susceptibility to breast cancer. However, polymorphisms have been examined which would give a modest elevation in risk. (Weber and Nathanson 2000).

The proto-oncogene *HRAS1* locates on chromosome 11p15 and encodes a protein involved in mitogenic signalling, the process by which an extracellular growth factor signal is transmitted to the nucleus (Junien et al. 1984, Barbacid 1987). A minisatellite composed of 30-100 units of a 28-base pair repeat is tightly linked to *HRAS1* (Capon et al. 1983). The carriers of a certain class of rare alleles of the minisatellite locus are proposed to be at increased risk of breast cancer, the relative risk being approximately twofold (Krontiris et al. 1993). However, two most recent studies using a more accurate method for allele sizing found no overall association and it seems that the earlier results are false due to misclassification of alleles resulting from methodological reasons (Firgaira et al. 1999, Tamimi et al. 2003). These findings suggested that the putative associations of *HRAS1* minisatellite alleles and breast cancer need to be re-evaluated with the new method of sizing alleles.

In one study an increased risk of breast cancer was observed to be associated with an Asn11Ser polymorphism of the *c-MYC* gene (Wirtenberger et al. 2005). *c-MYC* encodes for a transcription factor that is involved in cell proliferation and differentiation, cell cycle progression, malignant transformation, genomic instability/DNA-damage sensitization, and apoptosis (Patel et al. 2004).

*HER2 (ERBB2, NEU)* proto-oncogene gene locates on chromosome 17q21 and encodes for a transmembrane glycoprotein p185 with a tyrosine kinase activity (Schechter et al. 1984, Yang-Feng et al. 1985, Coussens et al. 1985, Akiyama et al.

1986). p185 is a member of the epidermal growth factor receptor family that controls a variety of cellular functions, including cell division, migration, adhesion, differentiation, and apoptosis (Yarden and Sliwkowski 2001). *HER2* amplification or overexpression is seen in about 25 % of breast cancers and has been associated with metastatic phenotype, endocrine therapy unresponsiveness, and poor prognosis (Menard et al. 2004). A common variant in *HER2*, a SNP Ile655Val in the transmembrane region could result in increased protein tyrosine kinase activity (Papewalis et al. 1991, Fleishman et al. 2002). Several studies have evaluated the association of this *HER2* polymorphism with breast cancer risk. Xie et al. (2000) first reported an increased risk of breast cancer among Val allele carriers in Chinese population. Several larger studies found no association (Millikan et al. 2003, Han et al. 2005, Benusiglio et al. 2005) but others suggested even a protective effect (Nelson et al. 2005, Cox et al. 2005c), leaving the question of *HER2* and genetic breast cancer association still controversial.

#### **2.4.8 Other genes studied for breast cancer association**

Several other genes from different pathways have been studied for association with breast cancer and many of them have not shown any significant associations. Genes with significant association with breast cancer risk are introduced here.

The *CASP8* gene on chromosome 2q33 encodes caspase-8, one of the initiator caspases that transduce apoptotic signals from the death receptors at the cell surface (Kischkel et al. 1998, Hengartner 2000). Caspase-8 is activated by external death signals and in response to DNA damage (Hengartner 2000). *CASP8* Asp302His has been estimated to account for approximately 0.3 % of the excess familial risk of breast cancer in populations of European ancestry, and this site was not polymorphic in Korean, Han Chinese or Japanese women. The functional consequences of the Asp to His substitution are not yet known, and further experiments are needed to establish whether Asp302His itself, or another variant in LD with it, is causative. (Cox et al. 2007). (Table 15).

*AIB1* (*NCOA3*) gene locates on chromosome 20q11 and encodes the AIB1/SRC-3 (amplified in breast cancer 1/steroid receptor coactivator 3) protein (Anzick et al. 1997).

*AIB1* is amplified and overexpressed in breast and ovarian tumours and it enhances ER-dependent transcription (Anzick et al. 1997). (Table 15).

**Table 15.** Other genes associating with breast cancer risk.

| Gene, locus                            | Polymorphism                             | Contribution to breast cancer  | References   |
|--|--|--|--|
| <i>CASP8</i> , 2q33                    | Asp302His                                | reduced risk allele dose-dependent, per allele OR 0.88, CI=0.84-0.92, $P_{trend}=1.1 \times 10^{-7}$ | MacPherson et al. 2004, Frank et al. 2005, Breast Cancer Association Consortium 2006, Cox et al. 2007                        |
| <i>AIB1</i> ( <i>NCOA3</i> ), 20q11    | 3'UTR CAG/CAA repeat length polymorphism | proposed to modify risk among <i>BRCA</i> mutation carriers but not confirmed in all studies         | Rebbeck et al. 2001, Kadouri et al. 2004, Haiman et al. 2000, Wilkening et al. 2005, Hughes et al. 2005, Spurdle et al. 2006 |
|  | Gln586His Thr960Thr                      | suggested protective effect among <i>BRCA</i> mutation negative                                      | Burwinkel et al. 2005  |
| <i>TNF<math>\alpha</math></i> , 6p21.3 | -417G>A                                  | increased risk suggested   | Gaudet et al. 2007   |
| <i>TNF<math>\beta</math></i> , 6p21.3  | -308                                     | conflicting results  | de Jong et al. 2003  |
|  | A252G                                    | suggested association in Korean population   | Lee et al. 2005c   |
| <i>FGFR2</i> , 10q26                   | intron 2 C>T (rs2981582)                 | increased risk allele dose-dependent   | Easton et al. 2007   |

The HLA class III subregion has been suggested to have a role in susceptibility to breast cancer (de Jong et al. 2003). Candidate genes in this region include *TNF $\alpha$*  (tumour necrosis factor) and *TNF $\beta$* . Association with breast cancer risk and polymorphisms in these genes have not been confirmed. (Table 15).

Recently, two large genome-wide searches identified the gene encoding fibroblast growth factor receptor 2 (*FGFR2*) on chromosome 10q26 to associate with increased risk of breast cancer (Easton et al. 2007, Hunter et al. 2007). *FGFR2* is a receptor tyrosine kinase that is amplified and overexpressed in 5-10 % of breast tumours (Moffa et al. 2004, Adnane et al. 1991, Jang et al. 2001). Somatic missense mutations of *FGFR2* that are likely to be implicated in cancer development have also been demonstrated in primary tumours and cell lines of multiple tumour types (Jang et al. 2001, Greenman et al. 2007). The minor allele of SNP rs2981582 in *FGFR2* gene in Europeans was associated with an increased risk in a dose-dependent manner, with a higher risk in homozygous (10.5 %) than in heterozygous carriers (6.7 %) or common homozygotes (5.5 %) (Easton et al. 2007). Haplotype analysis of 7 SNPs indicated that multiple haplotypes carrying the minor allele of rs2981582 were associated with an increased risk of breast cancer, implying that the association was being driven by




rs2981582 itself or a variant strongly correlated with it. However, the variant causally related to breast cancer risk could not be identified and functional studies will be required to determine which variant it is. (Easton et al. 2007). (Table 15).

## 2.5 Strategies for finding new breast cancer susceptibility genes

### 2.5.1 Linkage disequilibrium

Linkage disequilibrium (LD) is the non-random association of alleles at adjacent loci. The loci are in LD when a particular allele at one locus is found together on the same chromosome with a specific allele at a second locus, more often than is mathematically expected based on the allele frequencies. The extent and strength of LD in a population depends on time and recombinational distance between markers. LD gradually decays as the function of the number of generations, and the time required for disappearance of LD depends on the distance between the loci (Figure 5). (Jorde 1995).




**Figure 5.** Linkage disequilibrium around an ancestral mutation (a black star). Recombination causes the LD to gradually decay and limits the extent of the region of association over time but markers that are physically close tend to remain associated with the ancestral mutation. Chromosomal regions derived from the ancestral chromosome are shown in grey and new regions introduced by recombination in white. G = generations during which recombination occurs. M1-M8 = marker alleles.

Power to detect LD tends to be greatest when a single disease-causing mutation that accounts for a large proportion of the phenotypic variance of a trait has arisen recently on a relatively uncommon haplotype background (Jorde 2000). Mutations responsible for complex diseases will often persist for long periods of time because they are typically less subject to the effects of natural selection than are mutations responsible for Mendelian diseases. This can further diminish LD and the resultant power to detect it (Jorde 2000). In addition to recombination and natural selection LD is also influenced by other factors complicating the relationship between LD and physical distance, e.g. mutation, chromosomal location (variable recombination rates), demographic history (population structure, admixture or migration (gene flow), population growth, genetic drift), and gene conversion (Laan and Pääbo 1997, Zavattari et al. 2000, Abecasis et al. 2001, Daly et al. 2001, Pritchard and Przeworski 2001, Ardlie et al. 2001). The strength of LD and the distance over which it extends varies from one region of the genome to another. LD also varies among populations, with European populations typically showing lower nucleotide diversity and greater LD than African populations (Reich et al. 2001). In young populations LD is thought to extend further than in older populations (Mohlke et al. 2001, Ophoff et al. 2002, Varilo et al. 2003). Also microsatellite markers have been observed to detect LD over longer distances than SNP markers (Ophoff et al. 2002, Varilo et al. 2003). LD can be measured in a number of ways but the most important pairwise measures of LD are  $r^2$  and  $|D'|$ . Both range from 0 (no disequilibrium) to 1 (complete disequilibrium), but their interpretation is different.  $|D'|$  is a better measure for describing the patterns of LD and  $r^2$  is of more direct relevance to the power of association studies. (Pharoah et al. 2004).

The pattern of LD is structured in stretches of consistently high LD interspersed with short intervals of rapid breakdown (Daly et al. 2001) (Figure 6). These LD blocks result from the localization of recombination to irregularly spaced hot spots (Daly et al. 2001, Jeffreys et al. 2001). In the regions of high recombination rate LD is lower and in regions of low recombination LD is higher. The common haplotypes underlying blocks of LD can be defined by testing a sufficiently large collection of SNPs. Focusing on haplotype blocks greatly clarifies LD analyses. Once the haplotype blocks are identified they can be treated as alleles and tested for LD, thereby reflecting the underlying

population variation more accurately than any individual SNP (Daly et al. 2001, Johnson et al. 2001). By determining the extended haplotypes at any given locus in a population it is possible to identify exactly which SNPs will be redundant and which will be essential to capture the haplotypes of a gene or a region of LD, i.e. the haplotype-tagging SNPs, htSNPs (Johnson et al. 2001). Haplotype maps have been constructed and information on htSNPs can be found in databases (Patil et al. 2001, Dawson et al. 2002, Gabriel et al. 2002, Phillips et al. 2003, International HapMap Consortium 2003, International HapMap Consortium 2005, Conrad et al. 2006).

Linkage disequilibrium can be utilized in identification of disease loci e.g. in initial locus positioning, restricting and fine mapping the critical chromosomal region and searching for shared haplotypes, using different strategies (de la Chapelle and Wright 1998, Kruglyak 1999, Peltonen 2000).



**Figure 6.** The block-like structure of LD, with regions of low haplotype diversity separated by recombinational hot spots. Lines below the blocks represent examples of the number of common haplotypes that might be present for such blocks. The graphs plot LD as a function of distance. The plots show that within a block LD decays gradually with distance, or not at all. Within hot-spot areas LD falls away much more rapidly with distance. (Adapted from Goldstein 2001).

### 2.5.2 Positional cloning of Mendelian genes

Positional cloning begins with linkage analysis. In pedigrees with a hereditary (monogenic) disease the affected members share the disease phenotype and with an increased probability they also share the genomic region that carries the disease gene. Linkage analysis identifies the chromosomal region shared by the affected members in a family. This region is estimated by genotyping genetic markers (e.g. STRs), typically spaced by 5-10 cM, and then using the marker genotypes and the data on their possibilities of transmission in the family in the linkage analysis. A calculated LOD (logarithm of odds) score indicates the probability of the chromosomal region to be the one with the disease gene. The LOD scores of studied families can be combined and so strengthen the result and improve the finding of a true linkage. In addition to family or pedigree-based linkage analysis, sib-pairs or trios can be used. After the initial locus positioning LD-based fine-mapping is needed to identify the candidate-genes. In an ideal case only a few genes in a region of 1 cM are restricted as candidates. Mutation analysis of the candidate-genes finally indicates the disease gene, mutations in which associate with the disease phenotype. (Peltonen et al. 2000, Botstein and Risch 2003). Functional studies further confirm the gene-disease association.

Family-based linkage studies have been the foundation for the many successes in mapping of genes associated with Mendelian disorders, including many rare family cancer syndromes such as multiple endocrine neoplasia type 2 and adenomatous polyposis coli (APC). Linkage studies have also been successful in mapping *BRCA1* and *BRCA2*, and genes for other common cancers, such as *MSH2* and *MLH1* in colorectal cancer and *CDKN2A* in melanoma. Success of this approach is however, limited to genes with rare, highly penetrant alleles. Linkage studies lack power to detect the alleles conferring moderate risks that are likely to be the norm in complex disease. Genetic heterogeneity limits the usefulness of combining data from multiple families, as different genes may be responsible for disease clustering in different families. Within families, phenocopies and incomplete penetrance are a problem, as the carrier status of a putative disease allele cannot be definitively inferred from disease status. (Pharoah et al. 2004).

### **2.5.3 Strategies for finding genes for complex diseases**

#### **2.5.3.1 Case-control association study**

Although linkage studies have been highly successful in mapping the genes that underlie monogenic disorders, these studies are of limited use for investigating predisposition to polygenic disease, such as cancer (Pharoah et al. 2004). The main alternative to linkage studies for disease gene mapping is the association study, in which the frequency of a genetic variant in diseased individuals (cases) is compared to the frequency of the variant in individuals without the disease (controls) (Risch 2000, Cardon and Bell 2001). Allelic association is present when the distribution of genotypes differs in cases and controls. Linkage disequilibrium is utilized in association studies as it is based on the assumption that the affected share a genetic variant which is so close to the marker locus that the probability of a recombination event to occur between them is minimal. The observed marker-disease association in a region of high marker-marker LD in a case-control set indicates that the locus related to disease susceptibility is nearby. (Pharoah et al. 2004). Association studies for disease genes are generally based on the "common variant, common disease" hypothesis (Chakravarti 1999). Genetic variants arising a long time ago might have become common in the present population at frequencies ranging from a few percent upwards. Some of these variants might predispose to common diseases, and combinations of these variants are proposed to underlie differences in disease susceptibility. One possibility to enhance the detection of an association is haplotype analysis which combines the information of individual markers (Baynes et al. 2007). Haplotypes mark recognizable chromosomal segments that can be tracked through pedigrees and through populations when not broken up by recombination and be treated for mapping purposes as alleles. (Strachan and Read 2003). Genetic association studies provide an efficient design for identifying common variants that confer modest disease risks. (Pharoah et al. 2004). The power of association studies can be increased by selecting cases with a family history of the studied disease (Antoniou and Easton 2003).

#### **2.5.3.1.1 Genome-wide approach**

The strategies of association studies are the genome-wide approach and a candidate-gene (or pathway) approach. In the genome-wide approach the markers are genotyped across the whole genome, instead of only a few candidate-genes. One starting point is to use microsatellite (STR) markers for screening the genome to find associated chromosomal regions. This application utilizes linkage disequilibrium and sets of STR markers that are traditionally used for linkage studies and spaced by 5-10 cM. Marker allele distributions are compared between cases and controls, and statistically significant differences show markers that are in LD with the disease (breast cancer). LD regions are then covered with additional markers spaced more closely, to refine the association. Especially in younger populations the linkage disequilibrium is thought to extend further and in general STR markers detect LD over longer distances than SNP markers.

At present, mostly due to technical ease, the use of SNP markers is predominant and there are two general strategies for genome-wide approach. The first uses quasi-random or anonymous SNPs that are spaced accross the genome. The second strategy uses sets of LD-based htSNPs that are specifically chosen. (Jorgenson and Witte 2006). Current understanding is that 200,000-500,000 SNPs would be needed to adequately tag all SNPs with a minor allele frequency of 5 % or more (Kruglyak 1999). It has been suggested that concentrating on SNPs on genic regions in genome-wide association studies can be more efficient for detecting causal variants than the existing indirect approaches, which attempt to capture information on all variants (Jorgenson and Witte 2006).

#### **2.5.3.1.2 Candidate-gene approach**

The alternative to genome-wide association studies is the candidate-gene approach, in which the studies are focused on candidate-genes that encode proteins thought to be involved in the process of interest, e.g. in carcinogenesis, such as those involved in apoptosis, cell-cycle control, carcinogen metabolism or DNA repair. It is also possible to select the candidate-genes based on initial linkage or microsatellite marker association studies or animal models. Ideally, single nucleotide polymorphisms of

known functional significance are studied. The rationale for the candidate-gene approach is that by maximising the biological plausibility, the chances of success are increased. However, this approach is limited by its reliance on existing knowledge to identify candidate-genes based on function. (Pharoah et al. 2004).

In the ideal candidate-gene association study the common polymorphisms in the studied population and the extent of LD are first identified in order to estimate the minimum set of polymorphisms that need to be genotyped to be able to report on all other polymorphisms (htSNPs). When an association study is completed there might be residual uncertainty about the risks that are associated with the susceptibility allele. Also, the observed associated marker might not be the true functional variant and it has to be determined separately, possibly requiring functional assays. (Pharoah et al. 2004). Difficulties in assessing the actual risk-associated variant are also due to the fact that in most Western populations the linkage disequilibrium within genes is so uniform that differentiating the associated variant is impossible. At this point using another ancestral population, e.g. African or Asian, might provide further support due to weaker LD. (Risch et al. 2001, Easton et al. 2007).

Candidate-gene approach can be improved by sample selection. By choosing cases with a family history of the studied disease, e.g. breast cancer, or bilateral cases, improves the power and the sample size needed to detect an association is reduced, relative to a standard case-control association study (Antoniou and Easton 2003). The relative efficiency obtained by using familial cases is greater for rarer alleles (Antoniou and Easton 2003). Enrichment for genetic factors may also be sought by selecting cases with an early onset, although it may have a limited effect on the power to detect association in breast cancer (Antoniou and Easton 2003). Using a founder population has been extremely powerful for mapping Mendelian disorders because genetic heterogeneity is reduced and the disease might be associated with a single founder haplotype. In the search for genes for complex disorders the advantages are less clear but in the study of rarer variants of more recent origin a founder population might be useful. (Pharoah et al. 2004, Dunning et al. 2000). (Table 16).

Current efforts to identify genetic variation have concentrated on common polymorphisms. However, it is possible that much of the variation in cancer risk is due

to rarer alleles. In fact, virtually all susceptibility alleles identified so far have frequencies less than 1 %. The rarity of these variants is likely to be due to the fact that they are of recent origin and have not had time to spread through the population. It is possible that rare alleles turn out to be the more important component in cancer susceptibility. (Pharoah et al. 2004).

**Table 16.** Benefits of isolated versus outbred populations in genetic studies. (Adapted from Peltonen et al. 2000).

| <b>Population isolate</b>                                  | <b>Outbred population</b>               |
|--|---|
| Higher prevalence for some diseases                        | More affected people                    |
| More inbreeding and the opportunity to map recessive genes | More opportunity for replication        |
| More uniform genetic background                            | Markers more polymorphic                |
| Good genealogical records                                  | Genes mapped pertinent to more humanity |
| Easier to standardise phenotypic definitions               |   |
| Wider intervals of LD                                      |   |
| Closer to HWE  |   |
| Less migration and more intact families                    |   |
| More uniform environment                                   |   |


#### 2.5.4 The Finnish population

Finland has been inhabited continuously at least since the last glacial period, for some 10 000 years (Norio 2003a). It is not known exactly who these original arctic people were but most likely they were not the only immigrants who arrived in Finland, thus the country was inhabited by small immigrant groups mainly from the South and East but also from the West (Peltonen et al. 1999, Norio 2003a). Both paternally inherited Y-chromosomal haplotypes and maternally inherited mitochondrial sequences show an exceptional decrease in the genetic diversity of the Finns when compared with other European populations (Sajantila et al. 1996). The establishment of agriculture in Finland may have happened about 3 500 years ago, although it is not certain at which time and by which people it was started (Norio 2003a). In the historical era, the expansion of agriculture to "uninhabited" hunting grounds has had a decisive role in the generation of the Finnish disease heritage (Norio 2003b).

The size of the Finnish population has been estimated to have been about 2 500-10 000 until 3 300 years ago. In 1100 the population is estimated to have been less than 50 000 and in 1500-1700 around 300 000. (Norio 2003a). The first inhabited areas were the



Southern, Southeast, Southwest and coastal regions, the so called early-settlement region (Figure 7). The population expansion, populating the late-settlement region, i.e. the Eastern and Northern parts of Finland, started only after 1500 from the region of South Savo (Figure 7). Small groups of founders settled in uninhabited areas resulting in formation of small, rural sub-isolates (multiple founder effects). The population of the late-settlement region, including Northern Savo, has been expanding for 20-25 generations in isolation (mainly because of distance) and is still characterized by sub-isolates with 50 000 or less inhabitants (Nevanlinna 1972, Norio et al. 1973, Norio 2003b).



**Figure 7.** The early and late-settlement regions in Finland. The region of South Savo is shaded in grey. An example of a regional (internal) isolate of Kuusamo is also shown (shaded in black). (Adapted from Varilo and Peltonen 2004).

The main expansion of the Finnish population occurred in remarkable isolation due to linguistic, religious, cultural and geographical barriers. There has been very little immigration in Finland in the last 80-100 generations of expansion and population bottlenecks have occurred. Epidemics and particularly famines in a Northern country

sensitive to frost have probably been more important than wars in respect of the bottlenecks. The famine with its accompanying epidemics in 1696-1697 killed a quarter to a third of the population of fewer than 450 000. The latest bottleneck, due to consecutive years of crop failure, occurred as late as in 1866-1868 killing 15 % of the population. Official population statistics were founded as early as 1749. Thus, the population was about 480 000 in 1750, 1.6 millions in 1850 and 4 millions in 1950. Only during the most recent 10-12 generations, the number of people has grown rapidly to the present day 5.2 millions. The influence of the bottleneck phenomenon is similar to that of the founder effect. In both cases, genetic drift favours some genes and removes others. (Nevanlinna 1972, Norio et al. 1973, Norio 2003a).

#### **2.5.4.1 The Finnish Disease Heritage**

Genetic drift worked in the population sub-isolates modifying the gene pool resulting in enrichment of mainly recessive diseases, denoted as The Finnish Disease Heritage (FDH). This concept was first introduced in 1973 and it refers to monogenic diseases that are more prevalent in the Finnish population than in other populations (Norio et al. 1973). On the other hand, some diseases that are common elsewhere are extremely rare in Finland. Currently 36 rare hereditary diseases belong to the FDH, 32 of which are autosomal recessive, two are autosomal dominant and two are X-linked disorders (Norio 2003c). The gene has been mapped in 33 and characterized in 27 of the disorders belonging to the FDH (Norio 2003c). In the three non-mapped disorders, the number of known families is still small. Diseases of the FDH are typically caused by one major mutation originating from a common ancestor, i.e. a founder mutation, but for some diseases also other minor mutations are known (Norio 2003c). Regional clusters of some of the FDH diseases still exist, reflecting the regional sub-isolates characteristic for a founder population (Peltonen et al. 1999). Founder mutations are characterized by linkage disequilibrium (LD) and significant LD extends over large genomic regions around the disease alleles of the FDH, varying between 2 and 13 cM (Peltonen et al. 1999).

Outside of the FDH, dominant founder mutations in cancer predisposition syndromes, e.g. in hereditary nonpolyposis colorectal cancer, have spread and become highly enriched and specific for certain geographical areas (Nyström-Lahti et al. 1994, Moisio et al. 1996). Similar effect can be seen with *BRCA1* and *BRCA2* mutations, although these mutations are more various (Sarantaus et al. 2000). Thus, low-penetrance susceptibility genes and alleles can be expected to be enriched in a population like the Finns. Genes for several diseases of the FDH have been discovered in analyses exploiting the extent of LD (Järvelä et al. 1991, Hästbacka et al. 1992, Kere et al. 1993, Höglund et al. 1995, Varilo et al. 1996, Visapää et al. 1999). The Eastern Finnish population being a regional subpopulation is an especially suitable material for LD analysis (Soininen 1981, Pirinen 1982).

## **2.6 Methods for mutation detection**

However promising a candidate-gene for a hereditary predisposition is it must be shown to be mutated in affected people. Autosomal dominant disease-causing mutations are distinguished from polymorphisms by their functional consequences, their cosegregation with the disease in a family, and their absence in unaffected and unrelated individuals. In some circumstances the identification and interpretation of the mutations may, however, be more difficult. These are unsuspected locus heterogeneity, mutational homogeneity, mutations may not be unambiguously pathogenic or mutations are difficult to find, e.g. *BRCA1* and *BRCA2* genes are large and screening them for mutations is laborious. Mutation testing methods can be divided into two groups: 1) Mutation detection methods for testing samples for known mutations and 2) Mutation screening methods for screening samples for (unknown) mutations in an entire candidate-gene. (Strachan and Read 2003).

### **2.6.1 Heteroduplex analysis, HA**

Heteroduplex analysis is suitable for general mutation screening. It is based on heteroduplex formation when complementary single DNA strands of a wild-type allele

and a mutant allele are allowed to hybridize. Heteroduplexes are formed when PCR products are denatured and then renatured by cooling down slowly. The detection of heteroduplexes is based on their different mobility on nondenaturing polyacrylamide electrophoresis gels compared to wild-type (or mutant-mutant) homoduplexes. Heteroduplexes have lower electrophoretic mobility and they can be detected as extra slow-moving bands. After electrophoresis the bands are made visible by e.g. ethidium bromide staining. Insertions, deletions and most single base substitutions are detectable in fragments of  $\leq 200$  bp. HA method does not detect homozygous mutations and it does not reveal the precise position of the mutation. (Keen et al. 1991, White et al. 1992).

#### **2.6.2 Single-strand conformation polymorphism, SSCP/Conformation-sensitive gel electrophoresis, CSGE**

SSCP/CSGE can be used to detect known mutations or to screen for unknown mutations. Like HA, SSCP is based on different electrophoretic mobility between mutant and wild-type alleles. In SSCP the gene-specific PCR products are denatured to be single-stranded and run on a nondenaturing polyacrylamide gel. Single-strand molecules adopt conformations that are dependent upon their sequence. In 80-100 % of cases a single-base change is sufficient to cause altered electrophoretic mobility. Radiolabelled primers or silver staining can be used for detecting the bands on gels. Control samples must be run so that differences from the wild-type pattern can be noticed. SSCP is simple and adequately sensitive, but is inefficient for fragments longer than 200 bp, and does not reveal the nature or position of any mutation detected. The precise pattern of bands seen is very dependent on details of the electrophoresis conditions. (Orita et al. 1989, Spinardi et al. 1991, Sheffield et al. 1993).

#### **2.6.3 Protein truncation test, PTT**

PTT is a specific test for frameshift, splice-site or nonsense mutations that cause a truncated protein product (Roest et al. 1993, van der Lijst et al. 1994). In certain genes, e.g. *BRCA1* and *BRCA2*, missense mutations are seldom found and are probably usually

nonpathogenic. For such genes the PTT has several advantages. It conveniently ignores the nonpathogenic (nontruncating) silent or missense base substitutions, and it reveals the approximate location of any mutation. Also, in PTT analysis longer fragments can be analysed compared to HA and SSCP.

In the procedure coding sequence without introns (cDNA made by RT-PCR or large exons in genomic DNA) is PCR amplified using a special forward primer that includes (in addition to the gene specific sequence) a T7 promoter sequence and a eukaryotic translation initiator with an ATG start codon, designed so that the sequence amplified reads in-frame from the ATG. The PCR product is put into a coupled *in vitro* transcription-translation system, which uses the T7 promoter to make mRNA and the translation initiator to translate it. The protein product is run out on a SDS-polyacrylamide gel (PAGE). The protein products can be visualised by using autoradiography e.g. incorporating a radioactive labeled ( $S^{35}$ ) amino acid in the translation reaction. Truncating mutations result in shorter products than the wild-type protein and the size of the fragment reveals the position of the mutation. (Roest et al. 1993, van der Luijt et al. 1994).

#### **2.6.4 Restriction fragment length polymorphism, RFLP**

Restriction fragment length polymorphism analysis, RFLP, is based on the variation on the length of DNA fragments cut by different restriction enzymes. A mutation may create a cleavage site for a restriction enzyme which can be used in detecting the mutation. By this technique mutated alleles may be differentiated from wild-type alleles by analysis of patterns derived from cleavage of their DNA. If two alleles differ in the distance between sites of cleavage of a particular restriction endonuclease, the length of the fragments produced will differ when the DNA is digested with a restriction enzyme. (Strachan and Read 2003).

#### **2.6.5 Sequencing**

Sequencing can be used to further characterize and verify a mutation found in analyses done by other mutation screening methods. Sequencing can also be used to screen for

mutations but it can be more laborious than usual mutation screening methods if the mutations searched for are not known in advance. The two best-known DNA-sequencing techniques are the enzymatic (dideoxy-mediated chain-termination) method of Sanger et al. (1977) and the chemical degradation method of Maxam and Gilbert (1977). Of these two Sanger's method is more widely used.

### 3. AIMS OF THE STUDY

The general aim of this study was to investigate the genetic background of familial breast/ovarian cancer and sporadic breast cancer in Eastern Finnish population. Since the population in this area has been relatively stable and genetically isolated, founder effect in the hereditary forms of breast/ovarian cancer could be expected. The frequency and type of germ-line mutations in known high-risk breast/ovarian cancer susceptibility genes, *BRCA1* and *BRCA2*, were evaluated in Eastern Finnish breast/ovarian cancer families. In addition, the autosomes were screened to identify new chromosomal regions and genes on them as genetic risk factors for sporadic breast cancer in the Eastern Finnish population. Due to the character of a founder population, linkage disequilibrium could be utilized.

The specific aims of the study were:

- I To screen 36 breast/ovarian cancer families for *BRCA1* and *BRCA2* germ-line mutations for evaluation of the contribution of these genes in Eastern Finland.
- II To analyse the novel *BRCA2* exon 11 4088insA mutation in an Eastern Finnish family with multiple breast/ovarian cancers and favourable clinical course.
- III To study the effect of *BRCA1* 4216-2nt A→G splice acceptor site mutation on *BRCA1* protein product.
- IV To find new low-penetrance genetic risk factors for Eastern Finnish sporadic breast cancer by using autosome-wide microsatellite scan for linkage disequilibrium-based association in a case-control material.
- V To identify the putative breast cancer risk-associated gene on 22q12-q13 region by single nucleotide polymorphism (SNP) association analysis.

## 4. MATERIAL AND METHODS

### 4.1 Study subjects

#### 4.1.1 Breast and ovarian cancer families (I-III)

The pedigrees in the study are from the Kuopio University Hospital district and of Eastern Finnish origin. Of the below described 516 new breast cancer cases diagnosed between April 1990 and December 1995, 102 patients were selected on the basis of some positive family history of cancer, i.e. breast or ovarian cancer in one first degree relative and in one other relative, according to the interview data collected by the Kuopio Breast Cancer Project (Männistö et al. 1996). The criteria were not strict at this point. The 102 selected patients were invited for an additional interview, gynaecologic ultrasound examination and were offered preliminary genetic counseling. They were also asked to fill in a questionnaire to further determine their family history of cancer. A filled questionnaire was returned by 85/102 patients and 82/102 patients participated in the ultrasound examination and agreed to the interview. On the basis of the data arising from the pedigrees, 15/82 families were selected for the germ-line *BRCA1* and *BRCA2* gene mutation screening study. The selection criteria were a minimum of three first-degree relatives with breast or ovarian cancer. However, 7/15 families with less than three first-degree relatives with breast or ovarian cancer were included in the study, due to history of cancer at early age, bilateral breast cancer or several cases of breast/ovarian cancer in the second or third-degree relatives.

The questionnaire was also sent to ovarian cancer patients treated in Kuopio University Hospital between January 1993 and October 1997. Out of over 90 Eastern Finnish pedigrees, 14 were selected to be included in the study following discussions with the index patients. Criteria for the ovarian cancer families were at least two cases of ovarian cancer in the family. Seven other families fulfilling the criteria were also included in this study. Six of these seven families were identified through genetic counseling at the local Cancer Society polyclinic, and one was identified by the clinical oncologist of the research group in routine clinical consultations. Thus, altogether 36 breast and ovarian cancer families were included in the germ-line mutation screening



study (Tables 17 and 18). In addition to breast and ovarian cancers, other cancers were also reported among first degree relatives in 2/3 of the families.

**Table 17.** Characteristics of the studied breast/ovarian cancer families.

| Family  | No. of cancer in family |              | Other cancers in family   |
|---------|-------------------------|--------------|---|
|         | Breast                  | Ovarian      |   |
| BR10    | 7                       |              | gastric, colon x3, lung x2  |
| BR14/16 | 6                       |              | gastric, unknown cancer   |
| BR17    | 6 (1 bilat.)            |              | skin melanoma   |
| BR13    | 5                       |              | unknown cancer, lung  |
| PS20    | 3 (1 bilat.)            |              | Wilms tumour, brain, colon  |
| BR02    | 3                       |              |   |
| BJ01    | 3 (1 bilat.)            |              | leukaemia, unknown cancer x3  |
| BR12    | 4 (1 bilat.)            |              | lung  |
| BR08    | 3 (1 bilat.)            |              | lung  |
| BR05    | 3                       |              |   |
| PS04    | 2                       |              | skin melanoma, liver, thyroid, oesophageal, lymph node, endometrial |
| PS19    | 2 (1 bilat.)            |              | endometrial x2, unknown cancer x2                                   |
| BR01    | 2                       |              | endometrial   |
| BR04    | 2                       |              | melanoma  |
| BR11    | 2                       |              | lung, colon   |
| BR15    | 2                       |              | unknown cancer x2   |
| OV03    | 5                       | 2            | brain   |
| BR09    | 4                       | 3 (1 bilat.) | rectal, prostate  |
| PS02    | 4                       | 1            | gastric x2  |
| OV30    | 2                       | 3            | pancreas, gastric, lip  |
| OV01    | 2                       | 2            |   |
| OV85    | 3                       | 1            | lymphoma, leukaemia, colon  |
| BR06    | 4                       | 1            | cholelodal, gastric, lung, kidney                                   |
| BR18    | 2                       | 2            | prostate  |
| BR03    | 2                       | 1            | prostate, bladder, oesophageal or laryngeal, liver, unknown cancer  |
| BR07    | 1                       | 1            | gastric x2, endometrial x3  |
| OV28    | 2                       | 1            | lung, colon   |
| OV73    | 2                       | 1            | melanoma, gastric x2, cervix, thyroid, oesophageal                  |
| OV37    | 1                       | 1            | colon, lung   |
| OV84    | 1                       | 1            |   |
| OV20    |                         | 3            | unknown cancer x3   |
| OV22    |                         | 3            | lip, lung x3, gastric, lymphoma, laryngeal                          |
| OV36    |                         | 2            | pancreas, lung x3, endometrial, lip, cervix, gastric, colon         |
| OV38    |                         | 2            | endometrial x2, bone, prostate, neck, leukaemia, lymphoma           |
| OV77    |                         | 1            | neuroblastoma   |
| OV33    |                         | 1            | bladder, gastric  |

EDTA blood samples were collected from the breast and ovarian cancer patients participating in the study and from their relatives who were willing to participate and

who gave their consent. Genomic DNA was extracted from peripheral blood lymphocytes as described below.

#### **4.1.2 Population-based breast cancer case-control material (IV-V)**

The population-based sample material is carefully selected from the province of Northern Savo in the late-settlement area of Eastern Finland. This study is an extension of the Kuopio Breast Cancer Project, through which the cases for this study were ascertained (Männistö 1999). The material was collected from 1 919 women who came to Kuopio University Hospital (hospital district of 750 000 people) between April 1990 and December 1995 with any suspected breast disease for breast examination. Of these women 516 were diagnosed and histologically confirmed to have breast cancer. These women with breast cancer were asked to participate in the study and only twelve refused to participate. Thus, the participation/cooperation rate was 98 % (504/516). The recruitment protocol missed 88 women diagnosed with breast cancer in Kuopio University Hospital catchment area during the collection period, 51 of which were private patients, 11 were missed during the nurses' one-month strike in 1995 and 26 were treated elsewhere. Therefore the total contact rate was 85 % (516/604) and overall response rate 83 % (504/604). Thus, the sample is a representative collection of unselected typical breast cancer cases from Kuopio area. A proper DNA sample was available from 497 breast cancer cases for the association study.

In addition to breast cancer cases, the Kuopio Breast Cancer Project material includes randomly selected and individually matched age (within  $\pm 5$  years) and area-of-residence (urban/rural) controls from the National Population Register for each case of breast cancer (Männistö 1999). Seventy-two percent (359/497) of the cases and 73 % (334/458) of the population controls included in study V were born in the Province of Northern Savo. (Table 18). Each participant gave informed written consent. EDTA blood samples were collected from the breast cancer cases and controls participating in the study. Genomic DNA was extracted from peripheral blood lymphocytes as described below.

#### 4.1.2.1 Cases and controls in the autosome-wide scan for LD (IV)

For the autosome-wide scan for linkage disequilibrium with breast cancer 49 cases and 50 controls selected from the above described population-based material were used. Forty-nine breast cancer cases who did not have a strong family history of cancer according to the initial interview and who were born around Kuopio in four rural communities whereto moving has been minimal were selected for the study IV (Table 18). The included 50 controls are matched for age and long-term area-of-residence with the cases. Age of onset of breast cancer ranged between 30 and 82 years, median age of onset being 54.0 years. Seventy-three percent of the cases had breast cancer at or after the age of 50 years. The median age of the controls was 54.5 years (range 30-76 years).

#### 4.1.2.2 Cases and controls in refining of the breast cancer-associated region on 22q12-q13 (V)

For the further study of chromosome 22q12-q13 breast cancer association we had DNA available from 497 breast cancer cases and 458 age and long-term area-of-residence matched controls. The selected cases represent 96 % of the total of 516 breast cancers diagnosed in Kuopio University Hospital between 04/90 and 12/95.

**Table 18.** Summary of study subjects used in each study.

|                                   | Study I               | Study II              | Study III               | Study IV     | Study V                               |
|-----------------------------------|-----------------------|-----------------------|-------------------------|--------------|---------------------------------------|
| <b>Study design</b>               | Familial              | Familial              | Familial                | Case-control | Case-control                          |
| <b>Cancer</b>                     | Breast/ovarian        | Breast/ovarian        | Breast                  | Breast       | Breast                                |
| <b>No. of studied individuals</b> | 89,<br>in 36 families | 11,<br>in 1/36 family | 20, in 2/36<br>families | 49 + 50      | 497 + 458<br>(subsample<br>280 + 257) |

#### 4.2 Extraction of DNA from blood lymphocytes (I-V)

EDTA blood samples were collected from the familial breast and ovarian cancer patients participating in the study and from their relatives who were willing to participate and who gave their consent (I-III), and from the sporadic, population-based breast cancer patients and controls participating in the study (IV, V). Genomic DNA

was extracted from peripheral blood lymphocytes using standard chloroform-phenol extraction method (Vandenplas et al. 1984).

### **4.3 *BRCA* gene mutation detection (I-III)**

In study I the DNA samples were screened for germ-line mutations in the coding region and exon-intron boundaries of the *BRCA1* and *BRCA2* genes by heteroduplex analysis (Ganguly et al. 1993), protein truncation test (PTT) (Roest et al. 1993) and direct sequencing. Variants found in PTT and HA were confirmed by sequencing. In study II the *BRCA2* 4088insA mutation was detected in PTT analysis and confirmed by sequencing. In study III the *BRCA1* 4216-2nt A→G mutation was confirmed, after sequencing, with restriction fragment length polymorphism analysis (RFLP) using the restriction enzyme MseI (New England BioLabs, Beverly, MA, USA). Additionally, in study III the effect of the mutation on the intron splice-site was confirmed by cDNA analysis.

#### **4.3.1 PCR amplification and protein truncation test (PTT) of *BRCA1* and *BRCA2***



##### **exon 11 (I, II)**

The exon 11 of *BRCA1* and *BRCA2* gene were screened for germ-line mutations by protein truncation test (PTT) (Roest et al. 1993, Hogervorst et al. 1995). First, exon 11 of both genes was PCR-amplified from genomic DNA in three overlapping segments with a T7 promoter sequence attached to the 5' end of the forward primers (Plummer et al. 1995, Johannsson et al. 1996, Tavtigian et al. 1996) (Table 19, Figure 8). Annealing temperature for each primer was 56-58°C and Taq DNA polymerase (Promega, Madison, WI, USA) was used. The PCR products were used as a template for in vitro transcription translation reaction. For PTT the TNT T7 Coupled Reticulocyte Lysate System (Promega) was used according to manufacturer's instructions. The protein products were detected by autoradiography. First they were separated on a 12 % SDS-PAGE minigel (BioRad, Hercules, CA, USA), then gels were fixed with 1 M sodium acetate and dried. X-ray films were exposed in -70°C for 1-5 days and developed.

**Table 19.** Sequences of the primers used in PTT analysis of *BRCA1* and *BRCA2* gene exon 11.

| Primer name | Sequence 5' → 3'           | Location (nt) in cDNA   |
|-------------|----------------------------|-------------------------|
| BR1-AF      | T7-CTGCTTGTGAATTTTCTG      | 790 - 807               |
| BR1-AR      | CATGAGATCTTTGGGGTC         | 2355 - 2372             |
| BR1-BF      | T7-AGCAGAATGGTCAAGTG       | 1738 - 1754             |
| BR1-BR      | GTCCAGTTTCGTTGCCTC         | 2995 - 3012             |
| BR1-CF      | T7-TCAAGGTTTCAAAGCGCCAG    | 2701 - 2720             |
| BR1-CR      | GACGTCCTAGCTGTGTGAAG       | 4215+185nt in intron 11 |
| BR2-AF      | T7-GTTTATTGCATTCTTCTGTG    | 2138 - 2157             |
| BR2-AR      | TGACTTCCTGATTCTTCTAA       | 3580 - 3599             |
| BR2-BF      | T7-GTGTAAAGCAGCATATAAAAAAT | 3101 - 3122             |
| BR2-BR      | TTCGGAGAGATGATTTTGTG       | 5410 - 5430             |
| BR2-CF      | T7-GCCTTAGCTTTTACACAA      | 5230 - 5248             |
| BR2-CR      | CCCCCAAAGTACTACACAA        | 7069+135nt in intron 11 |

T7 promoter sequence: 5'-GGATCCTAATACGACTCACTATAGGAACAGACCACCATGG-3'

**A****B****Figure 8.** Exon 11 of *BRCA1* and *BRCA2* genes were analysed in three overlapping fragments using PTT. Size of each fragment (BR1-A - BR2-C) is shown in base pairs (bp). A) Schematic presentation of *BRCA1* exon 11. B) Schematic presentation of *BRCA2* exon 11.

#### 4.3.2 Heteroduplex analysis of *BRCA1* and *BRCA2* genes (I, III)

In study I and III exons 2, 3, 5-10 and 12-24 of the *BRCA1* gene were amplified from genomic DNA using the primers described by Friedman et al. (1994), Dynazyme DNA polymerase (Finnzymes, Espoo, Finland) and annealing temperatures 56-63°C (Table 20). In study I exons 2 and 12 of the *BRCA2* gene were amplified using the primers described by Couch et al. (1996) and exons 3-9 and 13-27B using primers described by Friedman et al. (1997) (Table 21). Annealing temperatures were 47-59°C and Dynazyme DNA polymerase was used. An aliquot of 3-8 µl of the PCR product was used for the DNA pairing reaction of the heteroduplex (HA) analysis. The sample was first denatured in 98°C for 5 minutes and then it was renatured by cooling the heatblock slowly to 37°C. The hetero- and homoduplexes are formed during the slow renaturation. After cooling, 6 µl of loading buffer (30 % glycerol, 0.25 % bromophenol blue, 0.25 % xylencyanol) was added into sample tube and then briefly centrifuged. The samples were analyzed on 10 % CSGE gel. Electrophoresis was carried out with 150V voltage for 12-15 hours depending on the size of the exon analysed. After electrophoresis the gels were either silver stained and dried or ethidiumbromide (EtBr) stained and photographed.

#### 4.3.3 Sequencing (I-III, V)

In study I the exon 10 of the *BRCA2* gene was screened for mutations by direct sequencing. The exon 10 was PCR amplified in two overlapping fragments using Phusion High-Fidelity DNA Polymerase (Finnzymes, Espoo, Finland) according to manufacturer's instructions. The forward and reverse primers were 5'- TGTTCCT ATGAGAAAGGTTGTGAGA-3' and 5'- TTGCCTGCTTTACTGCAAGA-3' for the first fragment, and 5'- TAAATGGAGCCCAGATGGAG-3' and 5'- AAAACACA GAAGGAATCGTCATC-3' for the second fragment. A 1.0-1.5-µl aliquot of the PCR product was included in the sequencing reactions using Big Dye Terminator v1.1 Cycle Sequencing Kit (Applied Biosystems) according to manufacturer's instructions. Sequences were run on an ABIPRISM 3100 Genetic Analyzer (Applied Biosystems)

**Table 20.** Primers used in the heteroduplex analysis (HA) of *BRCA1* gene.

| Primer pair | Exon size (bp) | Forward primer sequence 5' → 3' | Reverse primer sequence 5' → 3' | Fragment size(bp) |
|-------------|----------------|---------------------------------|---------------------------------|-------------------|
| Exon 2      | 100            | GAAGTTGTCAATTTATAAACCTTT        | TGCTTTTCTTCCCTAGTATGT           | 254               |
| Exon 3      | 54             | TCCTGACACAGCAGACATTTA           | TTGGAATTTTCGTCTCACTTA           | 349               |
| Exon 5      | 78             | CTCTTAAGGGCAGTTGTGAG            | TTCTACTGTGGTTGCTTCC             | 238               |
| Exon 6      | 89             | CTTATTTTGTGTCTTAAAAGG           | TTTCATGGACAGCACTTGAGTG          | 166               |
| Exon 7      | 140            | CACAACAAAGAGCATACATAGGG         | TCGGGTTCACTCTGTAGAAG            | 228               |
| Exon 8      | 105            | TGTTAGCTGACTGATGATGGT           | ATCCAGCAATTATTAAATAC            | 267               |
| Exon 9      | 47             | CCACAGTAGATGCTCAGTAAATA         | TAGGAAAATACCAGTTTCATAGA         | 211               |
| Exon 10     | 76             | TGGTCAGCTTTCTGTAATCG            | GTATCTACCCACTCTCTTCTCAG         | 242               |
| Exon 12     | 87             | GTCTGCCAATGAGAAGAAA             | TGTCAGCAAACCTAAGAATGT           | 265               |
| Exon 13     | 174            | AAATGGAAAGCTTCTCAAAGTA          | ATGTTGGAGCTAGGTCCTTAC           | 320               |
| Exon 14     | 127            | CTAACCTGAATTTACACTTACA          | GTGTATAAATGCCTGTATGCA           | 312               |
| Exon 15     | 191            | TGGCTGCCCAGGAAGTATG             | AACCAGAAATATCTTTATGTAGGA        | 338               |
| Exon 16     | 311            | AATTCTTAACAGAGACCAGAAC          | AATCTTAAACAGAGACCAGAAC          | 450               |
| Exon 17     | 88             | GTGTAGAACGTGCAGGATTG            | TCGCCTCATGTGGTTTA               | 264               |
| Exon 18     | 80             | GGCTCTTTAGCTTCTTAGGAC           | GAGACCAATTTCCAGCATC             | 351               |
| Exon 19     | 37             | CTGTCATTCTTCCTGTGCTC            | CATTGTTAAGGAAAAGTGGTC           | 251               |
| Exon 20     | 86             | ATATGACGTGTCTGTCTCCAC           | GGGAATCCAAATTACACAGC            | 401               |
| Exon 21     | 55             | AAGCTCTTCCTTTTGAAGTC            | GATGAGAAATAGAATAGCCTCT          | 298               |
| Exon 22     | 75             | TCCCATTTGAGAGTCTTGCT            | GAGAAGACTTCTGAGGCTAC            | 297               |
| Exon 23     | 60             | CAGAGCAAGACCCTGTCTC             | ACTGTGCTACTCAAGCACCA            | 255               |
| Exon 24     | 125            | ATGAATTGACACTAATCTCTGC          | GTAGCCAGGACAGTAGAAGGA           | 280               |

**Table 21.** Primers used in the heteroduplex analysis (HA) of *BRCA2* gene.

| Primer pair | Exon size (bp) | Forward primer sequence 5' → 3' | Reverse primer sequence 5' → 3' | Fragment size (bp) |
|-------------|----------------|---------------------------------|---------------------------------|--------------------|
| Exon 2      | 106            | TTCCAGGAGATGGGACTGAATTAG        | TGGGTTTTAGCAAGCATTTTTAG         | 302                |
| Exon 3      | 249            | GATCTTTAACTGTTCTGGGTCA          | CCCAGCATGACACAATTAATGA          | 423                |
| Exon 4      | 109            | AGAATGCAAATTTATATCCAGAGTA       | AATCAGATTATCTTTATAGAACAAA       | 249                |
| Exon 5&6    | 51&41          | TGTGTGGCATTTTAAACATCA           | CAGGGCAAAGGTATAACGCT            | 453                |
| Exon 7      | 115            | CCTTAATGATCAGGGCATTTC           | CAACCTCATCTGCTCTTCTTG           | 214                |
| Exon 8      | 50             | GGCATATCTTACCACCTTGTGA          | AGGTTTAGAGACTTCTCAAAGGC         | 406                |
| Exon 9      | 112            | CTAGTGATTTTAACTATAATTTTG        | GTTCAACTAAACAGAGGACT            | 164                |
| Exon 12     | 99             | AATTGACATTGAAGACTGACTTACTC      | AGCACTTTGGAGAGGCAGG             | 374                |
| Exon 13     | 73             | GCATCCGTTACATTCACCTGAAA         | ACGGGAAGTGTAACTTCTTAACG         | 310                |
| Exon 14A    | 428            | ACCATGTAGCAAATGAGGGTCT          | GCTTTTGTCTGTTTTCTCCAA           | 391                |
| Exon 14B    | 428            | CACAGAGTTGAACAGTGTGTAGG         | GGGCTTTAAAATTACCACCACC          | 297                |
| Exon 15     | 182            | GGCCAGGGGTGTGCTTTT              | AGGATACTAGTTAATGAAATA           | 314                |
| Exon 16     | 188            | TTTGGTAAATTTCAGTTTGGTTTG        | AGCCAACTTTTTGTTCGAGAG           | 395                |
| Exon 17     | 171            | CAGAGAATAGTTGTAGTTGTGAA         | AGAAACCTTAACCCATACTGC           | 306                |
| Exon 18A    | 355            | GATCCACTATTGGGGATTGC            | GATCTAACTGGGCCTTAACAGC          | 457                |
| Exon 18B    | 355            | GCAGATACCCAAAAGTGGC             | TCTGGACCTCCCAAAAAGTGC           | 384                |
| Exon 19     | 156            | AAGTGAATATTTTAAAGGCAGTT         | TATATGGTAAGTTTCAAGAAT           | 296                |
| Exon 20     | 145            | CACTGTGCCTGGCCTGATAC            | ATGTTAAATTCAAAGTCTCTA           | 296                |
| Exon 21     | 122            | GGGTGTTTTATGCTTGGTCT            | CATTTCAACATATTCCTTCTTG          | 304                |
| Exon 22     | 199            | AACCACACCCCTTAAGATGAGC          | GGGCATTAGTAGTGGATTTTGC          | 455                |
| Exon 23     | 164            | ACTTCTTCCATTGCATCTTCTCA         | AAAACAAAACAAAAATTCACATA         | 290                |
| Exon 24     | 139            | GCAGCGACAAAAAAACTCA             | ATTTGCCAACTGGTAGCTCC            | 365                |
| Exon 25     | 305            | GCTTTCGCCAAATTCAGCTA            | TACCAAAATGTGTGGTGATGC           | 426                |
| Exon 26     | 147            | GTCCCAAACTTTTTCATTTCTGC         | GGAGCCACATAACAACACCA            | 379                |
| Exon 27A    | 609            | CTGTGTGTAATATTTCGCTGCT          | GCAAGTCTTCGTGAGCTATTG           | 495                |
| Exon 27B    | 609            | GAATTCTCCTCAGATGACTCCA          | TCATTGCTCATTGTGCAACA            | 417                |



and analysed using Sequencing Analysis and Sequence Scanner v1.0 softwares (Applied Biosystems).

When aberrant mobility on heteroduplex gels or PTT protein product differing from the wild-type protein was detected in studies I-III, the variants were reamplified from genomic DNA and sequenced and analysed using dRhodamine Terminator Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) and AbiPrism 310 Genetic Analyzer (Applied Biosystems) according to manufacturer's instructions. The primers used for HA in study I and III were used also for sequencing the fragments. For sequencing the variant in PTT in study I and II new primers were designed (BR2-05F 5'- ATGAAGTGGGGTTTAGGGGC-3' and BR2-05R 5'-ATTTTGCTCCGTTT TAGT AG-3'). In study III primers SPL1F 5'-ACCGTTGCTACCGAGTGTCT-3' and SPL1R 5'- AGTTCAGCCATTTCTGCTG-3' and the above technique and equipment were used for sequencing of the purified cDNA.


In study V the variants from DHPLC analysis were re-amplified and sequenced using the primers used in DHPLC analysis and AbiPrism Big Dye Terminator Cycle Sequencing Ready Reaction kit with Ampli Taq DNA Polymerase (Applied Biosystems, Foster City, CA) according to manufacturer's instructions. Samples were analysed on AbiPrism 377 DNA sequencer and Sequencing Analysis 3.3 software (Applied Biosystems).

#### **4.3.4 RFLP analysis, extraction of RNA, and cDNA analysis of the *BRCA1* 4216-2nt A to G mutation (III)**

In addition to HA and sequencing, the 4216-2nt A→G splice-site mutation was confirmed by restriction fragment length polymorphism (RFLP) analysis from genomic DNA. A 265-bp-fragment was PCR amplified using the primers (12F and 12R) used in HA and the fragment was digested using 5 units of the restriction enzyme MseI (New England BioLabs, Beverly, MA, USA) in conditions recommended by the manufacturer. The recognition sequence for MseI is T/TAA. Two cleaving sites for MseI locate in the wild-type allele, but only one in the 4216-2nt A→G mutated allele. Thus, digested fragments are 82, 83 and 100 bp in non-carriers and 100 and 165 bp in

4216-2nt A→G mutation carriers. After digestion the fragments were separated on a 2.5 % LE agarose gel (FMC Bio Products), visualised by EtBr staining and photographed.

For study III also a fresh EDTA blood sample was drawn from *BRCA1* exon 12 4216-2nt A→G splice-site mutation carrier in one of the families. Total RNA was extracted from the blood lymphocytes using a Trizol LS Reagent (GibcoBRL Life Technologies, Rockville, MD, USA) according to the manufacturer's instructions. Reverse transcriptase PCR (RT-PCR) was performed from approximately 0.5 µg of total RNA using Titan™ One Tube RT-PCR System (Boehringer Mannheim, Mannheim, Germany) and primers SPL1F and SPL1R designed with the Primer3 program available at [http://www-genome.wi.mit.edu/cgi-bin/primer/primer3\\_www.cgi](http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi). SPL1F is located in *BRCA1* exon 11 at the nucleotides from 3855 to 3874 and SPL1R in exon 13 at the nucleotides from 4341 to 4360. The size of the normal cDNA fragment amplified with SPL1F and SPL1R primers is 506 bp and mutated fragment 507 bp. (Figure 9). The RT-PCR conditions were 50° for 30 min, 94° for 2 min, 46 cycles of 94° for 30 sec, 55° for 30 sec and 68° for 2 min and the final step 68° for 5 min. After amplification the size of the cDNA product was checked on a 2 % LE agarose (FMC Bio Products, Rockland, ME, USA) gel and purified with QIAquick PCR Purification kit (QIAGEN, Hilden, Germany). The purified cDNA product was used for sequencing as described above in 4.3.3.



**Figure 9.** Schematic presentation of reverse transcriptase (RT) PCR amplification for *BRCA1* 4216-2nt A→G splice-site mutation analysis. Length of *BRCA1* exons 11-13 and location of primers is shown in nucleotides (nt). The size of amplified cDNA fragment is 506 bp and in mutation carriers 507 bp.

#### 4.3.5 Analysis of mutation-associated haplotypes (I-III)

Microsatellite markers *D13S260*, *D13S1701* and *D13S267* were genotyped for the identification of the *BRCA2* gene mutation-associated haplotypes (Table 22). The microsatellites were PCR amplified with fluorescent-labelled primers using standard procedures (Sheffield et al. 1995) and analysed using an automated ABIPrism 310 Genetic Analyzer (Applied Biosystems), GeneScan 2.1 software and internal standard (see below at 4.4). Haplotypes were constructed manually utilizing the pedigree data. The location and distance between markers (kb and cM) was obtained from the human Genome Browser Gateway at the University of California at Santa Cruz (UCSC) (<http://genome.ucsc.edu>).

**Table 22.** Microsatellites used for *BRCA2* mutation-associated haplotypes.

| Marker              | Location on Chr.13  | Distance in kb | Genetic location (cM) |
|---------------------|---------------------|----------------|-----------------------|
| cen                 | 0                   |                |                       |
| <i>D13S260</i>      | 31334758 - 31335105 |                | 23.65                 |
| <i>BRCA2</i> (gene) | 31787617 - 31871805 | 452.512        |                       |
| <i>D13S1701</i>     | 32042530 - 32042819 | 170.725        |                       |
| <i>D13S267</i>      | 33162119 - 33162387 | 1119.300       | 26.87                 |
| tel                 | 114142980           |                |                       |

#### 4.4 Genotyping of microsatellites in the autosome-wide scan for LD (IV)

In study IV, 366 microsatellite markers from the 6th version of the Weber lab marker set (<http://research.marshfieldclinic.org/genetics/GeneticResearch/sets/scrset6.txt>) were genotyped in the first-stage autosome-wide scan for LD with breast cancer. The distance between markers span approximately 10 cM (range 1-20 cM) in the genome and the average heterozygosity of the markers was 0.76. The microsatellite repeats were PCR amplified from genomic DNA in single or multiplex reactions with fluorescent-labelled primers using standard procedures (Sheffield et al. 1995). PCR products from several microsatellites were pooled individually for each case and control and an internal GeneScan-500 [TAMRA] size standard (Applied Biosystems) was added into each sample. The size of the microsatellites was determined with an automated AbiPrism 310 Genetic Analyser (Applied Biosystems) and analysed using the GeneScan 2.1 software (Applied Biosystems) according to the size standard.

The number of samples was designed so that it was possible to run all controls, 45 of the cases and a blank control in one run on 96-well plate using AbiPrism 310. This was done to minimize possible variations in the sample running conditions (e.g. by using the same capillary for all samples). The remaining four of the cases were run first on another run to re-check that the reaction conditions for the marker are fine.

When linkage disequilibrium with breast cancer was observed, additional markers close to those with significant (or near significant) *P*-values were selected from the Genome Database (<http://www.gdb.org/>) for the second-stage scan (Table 23). Altogether 69 additional markers were genotyped. Intermarker distances and order were obtained from the databases at the Sanger Institute (<http://www.sanger.ac.uk>), the Human Genome Browser Gateway at the UCSC (<http://genome.ucsc.edu>), Cedar Genetics at the University of Southampton (<http://cedar.genetics.soton.ac.uk>) and Marshfield Center for Medical Genetics (<http://research.marshfieldclinic.org/genetics>). In the second-stage scan an AbiPrism 377 DNA Sequencer (Applied Biosystems) and GeneScan 2.1 (Applied Biosystems) software were used for size determination.

**Table 23.** Additional markers used in the second-stage of the genome-wide scan.

| <b>Chromosome 1</b>    | <b>Chromosome 7</b>  | <b>Chromosome 13</b> | <b>Chromosome 20</b>  |
|------------------------|----------------------|----------------------|-----------------------|
| <i>D1S2800</i>         | <i>D7S519</i>        | <i>D13S280</i>       | <i>D20S445=D20S79</i> |
| <b>Chromosome 2</b>    | <i>D7S2422</i>       | <i>D13S1265</i>      | <i>D20S887</i>        |
| <i>D2S363</i>          | <i>D7S2423</i>       | <b>Chromosome 14</b> | <i>D20S196</i>        |
| <i>D2S1277</i>         | <b>Chromosome 8</b>  | <i>D14S1016</i>      | <b>Chromosome 21</b>  |
| <i>D2S2289</i>         | <i>D8S1008</i>       | <i>D14S1054</i>      | <i>D21S1255</i>       |
| <i>D2S2358</i>         | <i>D8S1694</i>       | <b>Chromosome 16</b> | <b>Chromosome 22</b>  |
| <i>D2S2248</i>         | <b>Chromosome 9</b>  | <i>D16S3091</i>      | <i>D22S277</i>        |
| <i>D2S163</i>          | <i>D9S157</i>        | <i>D16S488</i>       | <i>D22S1142</i>       |
| <i>D2S2202</i>         | <i>D9S1684</i>       | <b>Chromosome 17</b> | <i>D22S924</i>        |
| <i>D2S2253</i>         | <b>Chromosome 10</b> | <i>D17S1810</i>      | <i>D22S283</i>        |
| <b>Chromosome 3</b>    | <i>D10S580</i>       | <i>D17S1289</i>      | <i>D22S426</i>        |
| <i>D3S3630</i>         | <i>D10S607</i>       | <i>D17S799</i>       | <i>D22S1177</i>       |
| <i>D3S1229=D3S1246</i> | <i>D10S201</i>       | <i>D17S1785</i>      | <i>IL2RB</i>          |
| <i>D3S3725</i>         | <i>D10S523</i>       | <i>D17S1820</i>      | <i>D22S1156</i>       |
| <i>D3S3520</i>         | <b>Chromosome 11</b> | <i>D17S941</i>       | <i>D22S272</i>        |
| <b>Chromosome 4</b>    | <i>D11S1302</i>      | <i>D17S1302</i>      | <i>D22S284</i>        |
| <i>D4S2975</i>         | <i>D11S1893</i>      | <b>Chromosome 19</b> | <i>D22S279</i>        |
| <i>D4S1615</i>         | <i>D11S1347</i>      | <i>D19S198</i>       | <i>D22S270</i>        |
| <i>D4S1575</i>         | <i>D11S897</i>       | <i>D19S420</i>       |                       |
| <b>Chromosome 6</b>    | <b>Chromosome 12</b> | <i>D19S559</i>       |                       |
| <i>D6S1660</i>         | <i>D12S1659</i>      | <i>D19S219</i>       |                       |
| <i>D6S980</i>          |                      | <i>D19S545</i>       |                       |
| <i>D6S1599</i>         |                      |                      |                       |

#### **4.5 Validating and selecting of SNPs for the refining of the breast cancer-associated region on 22q12-q13 (V)**

SNPs in genes in the 550-kb region between microsatellites *D22S426* and *D22S445* on chromosome 22q12-q13 were searched from databases (<http://www.sanger.ac.uk>, <http://www.ncbi.nlm.nih.gov/SNP> and <http://snp.cshl.org/db/snp>) and assays were designed for confirming the frequencies of 26 suggested SNPs and a small deletion on denaturing high-pressure liquid chromatography (DHPLC). Also the coding, 5'UTR and 3'UTR sequences of one of the genes in the associated region (*RABL4*) were screened for sequence variations using DHPLC. (Table 24). For these validations genomic DNA from 48 anonymous British samples and 48 of the 49 Eastern Finnish breast cancer cases used in the genome-wide LD scan were used. PCR amplifications for DHPLC analysis were carried out using: 1x GeneAmp PCR Gold buffer (Applied Biosystems), 0.16mM dNTP, 1.5-1.6 mM MgCl, 0.3-0.4 mM primers, 1.25-2.5 U AmpliTaq Gold DNA polymerase (Applied Biosystems) and 20 ng genomic DNA in a 60- $\mu$ l reaction and standard protocols. After amplification the samples were denatured and slowly renatured using 3 min denaturation at 95°C followed by cooling down to 25°C one degree per min. PCR amplified samples were analysed on WAVE DHPLC equipment (Transgenomics, Crewe, UK) according to manufacturer's instructions and using specific conditions for each fragment. The variants from DHPLC analysis were re-amplified and sequenced as described above in section 4.3.3.

Ten of the validated SNPs were selected for genotyping in 497 sporadic breast cancer cases and 458 matched controls (material described in 4.1.2.2). The SNP selection was based on the allele frequency (rare allele frequency > 0.1), location and distance between the SNPs (region covered as evenly as possible) and the suitability of the SNP to be detected by the 5' nuclease (Taqman®) assays. Also coding SNPs were preferred over intronic SNPs. The selected SNPs cover a 541-kb region on chromosome 22q12-q13 and adjacent SNPs are 0.9-132 kb apart. The selected SNPs are shown in Table 24 and Figure 10.

Artificial templates were prepared for both alleles in each assay for using as controls for the homozygous genotypes in the analysis. Long oligonucleotides were designed for using as primers in the artificial template synthesis (Table 25). For each SNP three


**Table 24.** Primers used in SNP validation on DHPLC. SNPs selected for genotyping are indicated with an asterisk (\*).

| GENE           | SNP ID     | Variation location<br>EXON/SNP | Location of exon<br>or SNP in<br>NT_011520.11 | MAF        | Forward primer sequence<br>5' → 3' | Reverse primer sequence 5' → 3' | Fragment<br>size (bp) |
|----------------|------------|--------------------------------|---|------------|------------------------------------|---------------------------------|-----------------------|
| <i>CACNG2</i>  | *rs738977  | intron 2 T/C                   | 16399443                                      | 0.19-0.39  | GCCAGTGGAAACCCCTTATT               | CCTTCCAGCTTGCTACCACT            | 255                   |
| <i>RAB1A</i>   |            | ex7 + 3'UTR                    | 16544771-16544969 <sup>‡</sup>                | NVF        | AGCCAAAGTGTGGAAGGAG                | AGCCAAAGTGTGGAAGGAG             | 282                   |
| <i>RAB1A</i>   | *rs3827351 | intron 7 C/A                   | 16547755                                      | 0.20-0.43  | TCCTTCTCCATCTCATTTG                | TGCTTCTGTGAGCCTCAACT            | 263                   |
| <i>RAB1A</i>   |            | ex6                            | 16549464-16549573                             | NVF        | GCTGTCTTCCAGAGTGTGA                | CGACAGGCCAAGTGAATA              | 193                   |
| <i>RAB1A</i>   |            | ex5                            | 16550479-16550596                             | NVF        | ACTATCTCTCCCTCTTGC                 | TGTGCTTGTAAACCAAGGA             | 203                   |
| <i>RAB1A</i>   |            | ex4                            | 16552698-16552757                             | NVF        | TCACCCCTAAGACAAGTCCAG              | AGGAGGTGTGACAGTCCAG             | 202                   |
| <i>RAB1A</i>   |            | ex3                            | 16553866-16553925                             | NVF        | TCCAGAACAAACCTTCCAC                | CCTGGTAAAGGCATCACCTC            | 203                   |
| <i>RAB1A</i>   |            | ex2                            | 16554343-16554419                             | NVF        | GCGGAGATCTCAAGTCAG                 | ATGGTGAACCGAGCACCTTCT           | 200                   |
| <i>RAB1A</i>   |            | ex1                            | 16562234-16562267                             | NVF        | GCTCTTGTCTCTGGGTACG                | TAACTGCCCTTTGGGAGAGA            | 200                   |
| <i>RAB1A</i>   |            | 5'UTR F&R                      | 16562268-16562636 <sup>‡</sup>                | NVF        | GGCTATGGTTGCTTGGAGAG               | GGCTGATCTCAAGGTTCAGT            | 292                   |
| <i>RAB1A</i>   |            | 5'UTR F&R2                     | 16562268-16562636 <sup>‡</sup>                | NVF        | GGCTATGGTTGCTTGGAGAG               | ACAAGAGGGGCTGCTAGAGA            | 371                   |
| <i>RAB1A</i>   |            | 5'UTR F&R3                     | 16562268-16562636 <sup>‡</sup>                | NVF        | GGCTATGGTTGCTTGGAGAG               | TTCAGAGCGCAAGTTTTCAGA           | 464                   |
| <i>PVALB</i>   | *rs3484    | 3'UTR G/A                      | 16587386                                      | 0.49       | GCTTCTCCAGATTTCACAG                | CGCTCTGACTCATCTTCTCT            | 234                   |
| <i>PVALB</i>   | rs739031   | intron 5 G/T                   | 16598664                                      | 0.048      | TGGCAGGAACAAACAACA                 | AACACGTCACCTCTGCACAC            | 224                   |
| intergenic     | rs713953   | Z82185 24057 C/T               | 16623838                                      | 0.07-0.156 | AATTCCTGCTAGCCTTCTCTG              | GCTTTCCAAAGAGAAAAGAGCA          | 218                   |
| <i>NCIF4</i>   | *rs760517  | intron 2 C/T                   | 16649501                                      | 0.32-0.48  | CCACAGACATGTTTTCAGCA               | CGGAAGTCCCTGGAAATATC            | 231                   |
| <i>NCIF4</i>   | *rs738148  | intron 8 A/G                   | 16661017                                      | 0.31-0.44  | TGAAGAGGTTCTGGGACAG                | CGGAAAACAGGAAGGCAATA            | 260                   |
| intergenic     | rs735563   | AL08637 52259 A/G              | 16687446                                      | 0.22       | GTGCTCTGCTCAGATGCAAA               | CCCTCTTGAGCTAGGTTCAGG           | 240                   |
| <i>CSF2RB</i>  | rs16845    | exon 7 Glu249Gln               | 16716958                                      | 0.04       | GTGGGACCCACTGAGAG                  | GAGGAAGCTTGGAGAGAGCA            | 266                   |
| <i>CSF2RB</i>  | *rs7285064 | exon 10 C/T Ser426             | 16720514                                      | 0.15-0.375 | AGACCGAGACCTCCAGAAC                | GGGATTTTCTGGCAGTGAAG            | 226                   |
| <i>CSF2RB</i>  | rs1801122  | exon 14 Pro603Thr              | 16724172                                      | 0.02       | CAGCTGCCTCAGATCTACCC               | CCAGAGGGACCAAGTTCAC             | 261                   |
| <i>CSF2RB</i>  | rs1801115  | exon 14 Gly647Val              | 16724305                                      | NA         | AGGGTGGGAGCCAGAAAT                 | ATAGCCACAGGGGCTGTCTT            | 244                   |
|                | rs131840   | exon 14 G/A Pro648             | 16724309                                      |            |                                    |                                 |                       |
|                | rs1801114  | exon 14 Val652Met              | 16724319                                      |            |                                    |                                 |                       |
| <i>CSF2RB</i>  | rs1801116  | exon 14 G/A Pro800             | 16724765                                      | NA         | TTTACTCCGAGCGAGAGAG                | TGTGGAGCTCCTTCCAAAT             | 217                   |
| <i>TST</i>     | not polym. | exon 4 T/C Asp268              | 16797673                                      | 0          | GCCTTGGCTGCCTACCTC                 | CAATTTCTAAAACATGTCATCTCT        | 229                   |
| <i>TST</i>     | rs4764     | exon 4 C/T His256              | 16797709                                      | 0.043      | CCCCTAAGCTGTGTCTCTG                | GCTTGCCGACAGAGGTAGG             | 234                   |
| <i>TST</i>     | rs1803405  | exon 4 Glu227Gly               | 16797797                                      | 0          | CCCTAAGCTGTGTCTCTGC                | GACCAAGGCCATCGTACAC             | 264                   |
| <i>TST</i>     | rs16363    | intron 3 tggtt ins/del         | 16800400                                      | 0.41-0.50  | AGGCCGCTCTACAGAACT                 | ACGAGGAACAAATGGATCAGC           | 229                   |
| intergenic     | *rs25095   | Z73420 37027 G/A               | 16824543                                      | 0.28-0.48  | CAGGAGCAAGGGAGGAG                  | AAATCAGAGCCAGTTCCTG             | 219                   |
| <i>KCTD17</i>  | rs710184   | intron 4 A/G                   | 16843774                                      | 0.19       | TGTCCTCTCACCTTGTAGC                | ACCACCCCGCAACATAC               | 221                   |
| <i>TMPRSS6</i> | rs736852   | intron 7 T/C                   | 16876720                                      | 0.31-0.47  | CCTTCTCAGTGGCCAGGTAA               | GGCCTGTGTGTACTTTTCTCAA          | 203                   |
| <i>TMPRSS6</i> | *rs733655  | intron 3 T/C                   | 16885566                                      | 0.20-0.475 | CATCTTCACCTCGTAGGT                 | TGAGTGTCTCTGCATCATCC            | 232                   |
| intergenic     | rs228937   | AL022314 77611 G/T             | 16911486                                      | 0.18043    | CTTCCGGCTCTGTACTGTAC               | CCTTATACCGGGTGCAGATG            | 231                   |
| <i>IL2RB</i>   | rs228940   | 3'UTR C/T                      | 16913835                                      | NA         | TTCCCTGACTCTTCAAGTGC               | GAAGCCATGTGGAAAGATG             | 265                   |
| <i>IL2RB</i>   | *rs228941  | 3'UTR G/C                      | 16914236                                      | 0.29       | CCCTGACGCTATTACCAAT                | ACGACACCGGAGGTACAAGT            | 238                   |
| <i>IL2RB</i>   | *rs228942  | exon 11 Asp391Glu              | 16915134                                      | 0.125      | CCCTAAGCAGCAACCACTCG               | AGGAGACTGGGGGAGAAGAG            | 262                   |

NVF= No variation found, NA= Not available/assay did not work properly, <sup>‡</sup>= Location of exon and UTR sequence

oligonucleotides were designed, two of them allele specific and one common for both alleles. Depending on the position of the SNP in the amplicon allele specific primers were either forward or reverse and common primer correspondingly reverse or forward. In the 100- $\mu$ l synthesis reaction 10 pmol of each primer (forward and reverse), 1X Klenow E.Coli Buffer, 0.4 mM dNTP, Klenow enzyme and 2  $\mu$ l of 0.5 M EDTA was used. After synthesis 900  $\mu$ l of dH<sub>2</sub>O was added and cooled to 4°C. For using in Taqman reactions the artificial templates were diluted to 1/100 - 1/1 000 000, depending on the assay.

#### Chr. 22



**Figure 10.** The selected SNPs cover a 541-kb region on chromosome 22q12-q13.

**Table 25.** Oligos used as primers in artificial template synthesis. Polymorphic base is in bold and underlined. Complementary sequence is indicated with an asterisk (\*).

| SNP       | Oligo sequence 5' → 3'   |   |  |  |
|-----------|--|---|--|--|
| rs738977  | <b>Forward T*</b><br>GGAGAGGATGCTTTTACCAAGTGGTTTACCGGCTGCAGGTAT<br>TTCAT   | <b>Forward C*</b><br>GGAGAGGATGCTTTTACCAAGTGGTTTACCGGCTGCAGGTAT<br>TTCAT              | <b>Reverse T/C</b><br>CACAGCATTAGCGGGAAGGATGTGTCTAATGGATGAAATAC<br>C                 |  |
| rs3827351 | <b>Forward C*</b><br>CGCTTGATAGAACTCAGAAATTAAAAACCCACCCCAACCTCCA<br>AACAGGAGTCAACAA  | <b>Forward A*</b><br>CGCTTGATAGAACTCAGAAATTAAAAACCCACCCCAACCTCCA<br>AACAGTGAAGTCAACAA | <b>Reverse C/A</b><br>CCACTTCAACACCTCAACAGAGTTTAAACCCGTAGCCCACTGG<br>CCTTTGTGACTC    |  |
| rs3484    | <b>Forward G*</b><br>CCTCTCTGCCCCTGAACACCCAAATCTCGGCCCTCTCTCGCCACCC<br>TCC   | <b>Forward A*</b><br>CCTCTCTGCCCCTGAACACCCAAATCTCGGCCCTCTCTCGCCACCC<br>TCC            | <b>Reverse G/A</b><br>AAACGAACCTGAACAGAAATGCAGGAGGGTGGC                              |  |
| rs760517  | <b>Forward C/T</b><br>GGTGGCCCTCCAAAGGGTTCTTCCAACTCGGCCTCCCCCAAGG<br>GCAGGCTCCAAAC   | <b>Reverse C*</b><br>TCAAGACACACTCGCACAGTTCACAGACTTTGGAGCCTGGCC<br>TGGGGAGTTTGGAGC    | <b>Reverse T*</b><br>TCAAGACACACTCGCACAGTTCACAGACTTTGGAGCCTGGCC<br>TGGGGAAAGTTTGGAGC |  |
| rs738148  | <b>Forward A</b><br>GGCCCTGGAGAGGAGCTAGTAGTGACCACTGGTTTAGCCAGT<br>TTAGCAGTTTCCA  | <b>Forward G</b><br>GGCCCTGGAGAGGAGCTAGTAGTGACCACTGGTTTAGCCAGT<br>TTGGCAGTTTCCA       | <b>Reverse A/G</b><br>GTAAGAGCCCTGAGTTCTCTGAGATGGAGATCTATGATTAGGG<br>TAAATGGAAACTGC  |  |
| rs7285064 | <b>Forward C/T</b><br>TTGGGGATTCTTGGCAGTGAGAACTATGAGCTGGTCTCTGG<br>CCAGAAGCTCTGGACTCCAGCCTTACACCTACCCGACTCGGTG<br>TCCAGGAGCGGCCTCACTCCACTC | <b>Reverse C</b><br>GGGTCAAGACCTCCCGCACCGGCTACAAACGGGATCTGGAGC<br>GAGTGGAGTG          | <b>Reverse T</b><br>GGGTCAAGACCTCCCGCACCGGCTACAAACGGGATCTGGAGT<br>AGTGGAGTG          |  |
| rs25095   | <b>Forward G*</b><br>AAGCGGTGACTTAGTGTAAATGATACTCAGCCTCACTTTCTAT<br>TGCAATCGTTTCCCAA   | <b>Forward A*</b><br>AAGCGGTGACTTAGTGTAAATGATACTCAGCCTCACTTTCTAT<br>TGCAATGTGTTCCCAA  | <b>Reverse G/A</b><br>CAITGACAGGAGGTGCGAATACGTTAGGCTGGCTGGCAGAT<br>TCAGCGTTGGGAAAC   |  |
| rs733655  | <b>Forward C/T</b><br>ACAGTGTGTGCTAACCACTACTACATGGGGTACGCCAGTTA<br>ACCAAGACAGATGTGCCTCCCT  | <b>Reverse T*</b><br>TGCCAGAGCCACGCCCTTCTTACCCACCACTGTCAAGCTTCACTA<br>AGGGAGGCA       | <b>Reverse C*</b><br>TGCCAGAGCCACGCCCTTCTTACCCACCACTGTCAAGCTTCACTA<br>AGGGAGGCA      |  |
| rs228941  | <b>Forward G/C</b><br>CCCTGACCTCTTCCCTGGGTTTCTGCCCCAGCCTCCCTCTCC<br>CTCCCTCCCC   | <b>Reverse G*</b><br>AGCGTGGCCATATTTGGGTTTGGAAAGCACGCTCAGGCTGC<br>CCTGTGACCGGGAGGGGA  | <b>Reverse C*</b><br>AGCGTGGCCATATTTGGGTTTGGAAAGCACGCTCAGGCTGC<br>CCTGTGACCGGGAGGGGA |  |
| rs228942  | <b>Forward C*</b><br>ACCTGTGGGTGCCCGGCCACACCCCTCATCAGGGTCTTCTCT<br>CTG   | <b>Forward A*</b><br>ACCTGTGGGTGCCCGGCCACACCCCTCATCAGGGTCTTCTCTCT<br>TG               | <b>Reverse C/A</b><br>AGGCTGCCAGGTGTACTTTACTTACGACCCCTACTCAGAGG<br>AAGA              |  |



#### **4.6 Genotyping of SNPs for the refining of the breast cancer-associated region on 22q12-q13 (V)**

For the ten selected SNPs forward and reverse primers and two fluorescent labelled probes were designed for 5' nuclease (Taqman) assays. Genotyping was carried out in 497 sporadic breast cancer cases and 458 matched controls according to assay manufacturer's instructions. The primer and probe sequences of the genotyped SNPs are shown in Table 26. In the 15- $\mu$ l Taqman reactions we used 1x Taqman universal PCR master mix (Applied Biosystems), 900 nM forward and reverse primers (Applied Biosystems), 100, 200 or 400 nM each probe (Applied Biosystems) and 20 ng of genomic DNA. All assays were carried out in 96-well plates. Each plate contained 72 test samples, 8 negative controls with no DNA and 8 positive controls for each SNP allele. The following PCR conditions were used on an MJ Tetrad thermal cycler (MJ Research, Waltham, MA, USA) for Taqman® reactions: first 2 min at 50°C, then 10 min at 95°C followed by 39 cycles of 15 s at 95°C and 1 min at 60, 62 or 64°C. Samples were post-read on an AbiPrism 7700 Sequence Detector (Applied Biosystems) and were analysed using the Allelic Discrimination Sequence Detection Analysis 1.7 software (Applied Biosystems). Genotypes were manually assigned under dye components.

#### **4.7 Allelic imbalance (AI) analysis (V)**

##### **4.7.1 Paraffin-embedded samples and extraction of DNA for AI analysis**

Paraffin-embedded samples of breast cancer tissue from 45 of the 49 cases in study IV were used for the AI analysis in study V. An experienced pathologist microdissected the samples using a 24 G needle under a light microscope in carefully matched tumour areas containing at least 70 % tumour cells (Tuhkanen et al. 2004). For each specimen DNA was extracted from five 10  $\mu$ m thick consecutive sections by the proteinase K-phenol-chloroform method according to standard protocols (Karjalainen et al. 2000).

**Table 26.** Primer and probe sequences of the genotyped SNPs.


| RefSNP ID | PCR primer sequence   | C (nM) | Probe sequence <sup>a,b</sup>  | C (nM) | Annealing temp. |
|-----------|---|--------|--|--------|-----------------|
| rs738977  | F: GAGGATGCTTTTACCAAGTGGTTT<br>R: AGCATTAGCGGGAAGGATGT            | 900    | A1: CGGCTGCGGGTATTTTCATCCAT (v) <sup>c</sup><br>A2: CCGGCTGCAGGATTTTCATCCATT (f) <sup>c</sup>      | 200    | 62°C            |
| rs3827351 | F: TTGATAGAACTCAGAAATTTAAACCCAC<br>R: CTTC AACACCTCAACAGAGTTTAACC | 900    | A1: CCAAAACAIGAGTCACCAAAGGCCA (v) <sup>c</sup><br>A2: CAAAACAGAGTCACCAAAGGCCAGT (f) <sup>c</sup>   | 200    | 62°C            |
| rs3484    | F: CTCTGCCCTGAACACCCAA<br>R: CGAACTGAACAGAAATGCAGGA               | 900    | A1: CTCGGCCCTCTTGCCACC (v) <sup>c</sup><br>A2: CTCGGCCCTCTTGCCAC (f) <sup>c</sup>                  | 200    | 62°C            |
| rs760517  | F: TGCCCTCCAAGGGTTCTTC<br>R: AGACACACTCGCACAGTTCACAG              | 900    | A1: TCCAAAACCTCCCCCAAGGCC (f)<br>A2: TCCAAAACCTCCCCCAAGGCCA (v)                                    | 400    | 60°C            |
| rs738148  | F: CCTGGAGAGGAGCTAGTAGTGACC<br>R: AGAGCCCTGAGTCTCTGAGATG          | 900    | A1: TTAGCCAGTTTAGCAGTTTCCATTACCCCT (f)<br>A2: TTAGCCAGTTTAGCAGTTTCCATTACCCCT (f)                   | 400    | 64°C            |
| rs1801117 | F: GGGATTTTCTGGCAGTGAGAAA<br>R: TCAGGACCTCCCGCACC                 | 900    | A1: CTCACTCCACTCGCTCCAGATCCC (f) <sup>c</sup><br>A2: CTCACTCCACTCGCTCCAGATCCC (v) <sup>c</sup>     | 120    | 62°C            |
| rs25095   | F: CGGGTGACTTAGTGTAATGATACTCAG<br>R: TGCACAGGAGGTGCGAATAC         | 900    | A1: ACTTCTATTGCAATGTTTCCCCAACG (v) <sup>c</sup><br>A2: TTCTATTGCAATGTTTCCCCAACG (f) <sup>c</sup>   | 200    | 62°C            |
| rs733655  | F: GTGTGTGCTAACCACTACTACATGG<br>R: CAGAGCCACGGCTTTCTTACC          | 900    | A1: TGCCTCCCTTGTGAAGCTGACA (v)<br>A2: CCTCCCTCGTGAAAGTGACAGTG (f)                                  | 200    | 62°C            |
| rs228941  | F: TGACCTCTTCCCTGGGTTTCT<br>R: GTGGCCATATTGGGTTTGG                | 900    | A1: CTCCTCCCTCCCGTCCACAGG (f)<br>A2: CCTCCCTCCCGTCCACAGGCA (v)                                     | 400    | 62°C            |
| rs228942  | F: CTGTGGGTGCCCCGG<br>R: CCTGCCAGGTGACTTTACTTACGA                 | 900    | A1: CACCTCATCAGGGTCTTCTCTGAG (f) <sup>c</sup><br>A2: ACACCTCATCAGGTTCTTCTCTGAGTAG (v) <sup>c</sup> | 100    | 62°C            |

<sup>a</sup> Probe label (v)= VIC, (f)= FAM. A1= allele 1 (major), A2= allele 2 (minor).<sup>b</sup> Polymorphic base underlined.<sup>c</sup> Complementary sequence.

#### 4.7.2 Genotyping of microsatellites for AI analysis

For the AI studies 13 microsatellite markers from study IV were used (Figure 11.). These markers span an approximate 6.8 Mb on chromosome 22. Microsatellite repeats were PCR amplified from genomic (blood sample) DNA (in study IV) and from tumour DNA with fluorescent-labelled (FAM, HEX or TET) forward primers and an ordinary reverse primer using standard procedures (Sheffield et al. 1995). Fragment sizes were analysed using internal size standard, AbiPrism 310 Genetic Analyser (Applied Biosystems) and GeneScan 2.1 software (Applied Biosystems). Some of the markers were run multiplex. Intermarker distances and order were obtained from Human Genome Browser Gateway at the University of California at Santa Cruz (<http://genome.ucsc.edu>) and at <http://www.sanger.ac.uk>.

Chr. 22



**Figure 11.** Microsatellite markers used in the AI study. Location of each marker is shown in megabases (Mb) from the centromere.

#### 4.7.3 Calculation of AI values

AI was calculated from the informative cases which were heterozygous in the normal tissue (blood) sample, using the peak heights from GeneScan analysis in formula  $AI = (T2 \times N1) / (T1 \times N2)$ , where T is the tumour allele and N is the normal allele (1 and 2 representing the two alleles). An AI value greater than 1.67 or less than 0.60 was considered AI (Canzian et al. 1996). Samples with values from 0.6 to 0.7 and from 1.55 to 1.67 were re-amplified and reanalysed at this particular locus. Only when this second result was positive tumour was assessed to have AI. A locus with AI in more than 30 % of the cases was considered significant.

### 4.8 Statistical analyses (IV, V)

#### 4.8.1 Microsatellite and SNP allele and genotype frequencies (IV, V)

Significance levels for comparisons of the microsatellite and SNP allele and genotype frequencies between cases and controls were computed using Fisher's exact test and Monte Carlo approximation implemented in SPSS.  $\chi^2$  and Fisher's exact tests were also performed to compare the frequency of the possible associated allele versus the frequency of all other alleles pooled together between cases and controls. *P*-values less than or equal to 0.05 were considered significant and *P*-values were unadjusted for multiple comparisons.

#### 4.8.2 Hardy-Weinberg equilibrium (IV, V)

Deviation of the SNP and microsatellite genotype distributions from the Hardy-Weinberg equilibrium (HWE) was assessed using the standard  $\chi^2$  -test in Associate v2.33 or HWE softwares of the LINKUTIL software package (<http://www.genemapping.cn/util.htm>, Terwilliger and Ott 1994), analysing cases and controls combined as well as cases and controls separately. *P*-values less than 0.05 were considered as significant deviation from the HW equilibrium. *P*-values in HW equilibrium tests were not adjusted for multiple correction.

#### **4.8.3 Breast cancer risk estimations (Odds ratios, OR) (IV, V)**

Breast cancer-associated risks for the SNP genotypes and rare/common allele carriers and the associated alleles of the significant microsatellites were estimated as odds ratios (OR) with 95 % confidence intervals (CI) using crosstabulation in SPSS. Odds ratio estimation was considered reliable when the 95 % confidence interval did not intersect unity.

#### **4.8.4 Haplotype estimation (IV, V)**

In study IV the haplotype frequencies were computed using the 'haplo.score' software (Schaid et al. 2002). Haplo.score uses unphased genotype data and via the EM (Expectation Maximization) algorithm (Slatkin and Excoffier 1996) it estimates the haplotype frequencies. This software implements a global score test for haplotype frequency differences between the cases and controls, and provides haplotype-specific tests, which allow evaluation of all individual haplotypes when the global score test is significant. To improve the  $\chi^2$  approximation of the score test we used a threshold level of 0.005 and omitted haplotypes with frequency less than 0.5 %. Marker combinations where the global score test for differences in haplotype frequencies between cases and controls reached significance (at  $P=0.05$  level), were investigated further by simulations (10 000). This involved simulating the case/control status of the individuals in the study and performing the test each time. The simulations were carried out using haplo.score.

In study V the haplotype frequencies for multiple SNP loci were estimated using the EM algorithm implemented in SNPAllyze v4.0 software (Dynacom, Chiba, Japan). SNPAllyze was also used for testing the association of the estimated haplotypes and breast cancer. SNPAllyze implements a global test for haplotype frequency differences between cases and controls and provides haplotype-specific tests which allow evaluation of all individual haplotypes. It also provides permuted (empirical)  $P$ -values for the global test and individual haplotypes. Permutation tests were done using FASTEHPLUS (Zhao and Sham 2002) and SNPAllyze. All permuted  $P$ -values were reached using 10 000 replicates.  $P$ -values  $\leq 0.05$  were considered significant.

#### **4.8.5 Pairwise linkage disequilibrium (V)**

Linkage disequilibrium (LD) between two SNPs was estimated by calculating the  $D'$ -values for all 45 possible pairs of the ten SNPs using the combined set of cases and controls (altogether 891-931 samples for each SNP) (Devlin and Risch 1995).

#### **4.8.6 Power estimations (V)**

Power estimations for the SNP association studies were calculated using the Genetic Power Calculator at <http://pngu.mgh.harvard.edu/~purcell/gpc/>, case-control for discrete traits (Purcell et al. 2003). In these estimations we used  $\alpha=0.05$  and breast cancer prevalence of 0.8 % (<http://www.cancerregistry.fi/>).  $D'$  was set at 1 (assuming complete linkage disequilibrium between marker allele and disease-associated allele) and the allele frequencies were assumed equal for the risk SNP and the marker SNP. Also the risks were assumed equal (1.5 or 2) for the homozygous and heterozygous high-risk allele carrying genotypes.

### **4.9 Ethical aspects**

All study participants gave their informed written consent. The mutation screening study was approved by the joint ethics committee of Kuopio University and Kuopio University Hospital and appropriate permission from the Ministry of Social Affairs and Health in Finland was obtained. The Kuopio Breast Cancer Project was approved by the joint ethics committee of Kuopio University and Kuopio University Hospital.

## 5. RESULTS

### 5.1 *BRCA1* and *BRCA2* mutations in breast/ovarian cancer families (I-III)

Altogether 89 individuals in 36 breast and ovarian cancer families were studied and five different *BRCA1/BRCA2* gene mutations were found; two in *BRCA1* gene and three in *BRCA2* (Table 27). In addition to the truncating mutations three polymorphisms were detected in the *BRCA1* gene and seven in *BRCA2* gene (Table 28).

**Table 27.** *BRCA1* and *BRCA2* gene mutations found.

| Mutation             | Family | Index case*   | No. of brca in family <sup>†</sup> | No. of ovca in family <sup>§</sup> | Other cancers in family <sup>‡</sup>  |
|----------------------|--------|---------------|------------------------------------|------------------------------------|---|
| <b><i>BRCA1:</i></b> |        |               |                                    |                                    |   |
| ex 12<br>4216-2ntA→G | BR01   | brca (36)     | 1 (50)                             | -                                  | endometrial (50)  |
| ex 12<br>4216-2ntA→G | OV01   | ovca (46)     | 2 (51, 43)                         | 1 (38)                             | na  |
| ex 20<br>5730 C→T    | PS04   | -             | 2 (32, 52)                         | -                                  | melanoma (77),<br>liver (43),<br>thyroid (60-70),<br>oesophageal (65),<br>lymph node (29)<br>endometrial (73) |
| <b><i>BRCA2:</i></b> |        |               |                                    |                                    |   |
| ex 9<br>999del5      | BR17   | brca (34, 45) | 4 (25, 40 bilat., 43, 58)          | -                                  | melanoma (35)   |
| ex 9<br>999del5      | OV30   | ovca (62)     | 2 (27, na)                         | 2 (na, na)                         | gastric (na),<br>lip (na),<br>lymphoma (4),<br>pancreas (na)  |
| ex 11<br>4088insA    | BR09   | brca (74, 77) | 2 (86, 46)                         | 3 (61, 58, 58 bilat.)              | rectal (57),<br>prostate (80)   |
| ex 11<br>6503delTT   | OV03   | ovca (51)     | 5 (37, na, na, na, na)             | 1 (na)                             | brain (na)  |

\*Carcinoma and age at diagnosis of the index patient

<sup>†</sup>Number of breast cancer cases in the family, index case not included, age at diagnosis in parenthesis.

<sup>§</sup>Number of ovarian cancer cases in the family, index case not included, age at diagnosis in parenthesis.

<sup>‡</sup>Other cancers in family, age at diagnosis in parenthesis.

na=not available

#### 5.1.1 *BRCA1* mutations found (I, III)

The detected *BRCA1* gene mutations were 4216-2nt A→G in exon 12, and 5370 C→T in exon 20. The 5370 C→T mutation in exon 20 was found in sequencing analysis and

exon 12 4216-2nt A→G mutation was detected in heteroduplex analysis and confirmed by RFLP and sequencing from genomic DNA and cDNA.

**Table 28.** Polymorphisms found in the *BRCA1* and *BRCA2* genes.

| Gene/<br>Exon/Intron | Alteration* | Effect              | rs Number <sup>†</sup> | Observed in |                   |
|----------------------|-------------|---------------------|------------------------|-------------|-------------------|
|                      |             |                     |                        | Cases       | Controls          |
| <b><u>BRCA1:</u></b> |             |                     |                        |             |                   |
| Intron 7             | -34ntT→C    | IVS7 -34T→C         | -                      | 1/24        | na                |
| Exon 13              | T4427C      | Ser1436Ser (S1436S) | rs1060915              | 12/24       | na                |
| Exon 16              | A4956G      | Ser1613Gly (S1613G) | rs1799966              | 11/24       |                   |
| <b><u>BRCA2:</u></b> |             |                     |                        |             |                   |
| 5'UTR                | 203G→A      | 5'UTR -26nt G→A     | rs1799943              | 14/33       | 6/31 <sup>‡</sup> |
| Intron 8             | +56ntC→T    | IVS8 +56C→T         | rs2126042              | 7/33        | 9/31 <sup>‡</sup> |
| Exon 10              | A1093C      | Asn289His           | rs766173               | 1/29        | na                |
| Exon 10              | A1342C      | Asn372His           | rs144848               | 17/30       | na                |
| Exon 10              | A1593G      | Ser455Ser           | rs1801439              | 1/34        | na                |
| Intron 14B           | +53ntC→T    | IVS14 +53C→T        | rs11147489             | 1/33        | 0/31 <sup>‡</sup> |
| Intron 24            | -16ntT→C    | IVS24 -16T→C        | rs11571818             | 1/33        | 1/31 <sup>‡</sup> |

\* nt corresponding to mRNA sequence GenBank U43746; <sup>†</sup> RefSNP number, obtained from the NCBI dbSNP at <http://www.ncbi.nih.gov/SNP/>; <sup>‡</sup> Anonymous population controls were used. na= not analysed


#### 5.1.1.1 *BRCA1* exon 12 4216-2nt A→G (III)

The *BRCA1* exon 12 4216-2nt A→G mutation was found in 3/8 and 3/12 individuals tested in two families. This mutation substitutes the adenine (A) nucleotide at position 4216-2nt to guanine (G) and disrupts the splice acceptor AG dinucleotide. The substituted guanine creates a new splice acceptor AG with an adenine at position 4216-3nt causing splicing error which leads to the inclusion of an extra nucleotide G in the mRNA. Aberrant splicing causes a frame-shift and changes the codon GGU for Glycine at position 1366 to GGG and codon GAA for Glutamine at 1367 to translation termination codon UGA (Figure 12). This leads to a truncated *BRCA1* protein of 1366 amino acid instead of the 1863-amino-acid wild-type protein.

In family A (BR01) the 4216-2nt A→G mutation was found in one breast cancer patient (female) and in two healthy individuals (male and female). Other five tested family members (male and female) were non-carriers. In family OV01 one breast cancer patient, one ovarian cancer patient and one healthy male were 4216-2nt A→G mutation



carriers. Nine other tested family members were non-carriers, one of them being affected with breast cancer at the age of 51 years.



**Figure 12.** The 4612-2nt A to G splice-site mutation in *BRCA1* gene leads to aberrant splicing and truncated protein. Wild-type sequence is shown in the upper part and mutated sequence in the lower part. The splice acceptor *ag* dinucleotide is shown in italics.

#### 5.1.1.2 *BRCA1* exon 20 5370 C→T (I)

The 5370 C→T mutation in exon 20 was found in one family (PS04). This mutation changes the codon for Arginine (AGA) at position 1751 to a translation termination codon UGA. This leads to a truncated *BRCA1* protein of 1751 amino acid. The mutation was found in the only individual (female melanoma patient) tested in the family.

#### 5.1.2 *BRCA2* mutations found (I, II)

Three mutations were detected in the *BRCA2* gene; 999del5 in exon 9, 4088insA in exon 11 and 6503delTT in exon 11. The 4088insA and 6503delTT mutations were detected in PTT analysis and confirmed by sequencing. Exon 9 999del5 was detected in heteroduplex analysis and confirmed by sequencing.

#### **5.1.2.1 *BRCA2* exon 9 999del5 (I)**

The exon 9 999del5 mutation was found in two families. This mutation is a five base-pair deletion TCAAA, starting at nucleotide 999 in codon 257. The deletion causes a frameshift leading to early translation termination in codon 273 and a truncated protein of 273 amino acids instead of the 3418-amino-acid wild-type protein.

In family BR30 the mutation was found in a breast cancer patient who was the only individual whom a sample was available for testing.


In family OV30 the 999del5 mutation was found in three of eleven tested family members. One of the mutation carriers had ovarian cancer and the other two were healthy females.

#### **5.1.2.2 *BRCA2* exon 11 4088insA (II)**

The 4088insA mutation was found in one family (BR09). The insertion of an adenine in a chain of 7 adenines at nucleotides 4081-4087 causes a frameshift leading to the change of Asparagine (AAT) at codon 1287 to Lysine (AAA), and the Asparagine (AAT) at 1288 to a stop codon UAA. This leads to premature translation termination at 1288 and a truncated protein of 1288 amino acids (Figure 13). In the family BR09 the mutation was found in 6/11 tested family members. One male carrying the mutation is healthy. One male carrier is affected with prostate cancer and four female carriers are affected with bilateral breast cancer, bilateral ovarian cancer, breast, ovarian and rectal cancer, and one with breast and ovarian cancer. Other five tested family members were non-carriers.

#### **5.1.2.3 *BRCA2* exon 11 6503delTT (I)**

The third *BRCA2* gene mutation detected was 6503delTT in exon 11. This mutation was found in one family (OV03). The deletion of nucleotides 6503 and 6504 causes a frameshift starting at codon 2092 and creates a stop codon at 2098 leading to a truncated protein of 2098 amino acids. In the family OV03 four 6504delTT mutation carriers were detected, three of them healthy (one male and two females) and one affected with ovarian cancer. The three other tested family members were non-carriers.



**Figure 13.** Sequencing of the *BRCA2* 4088insA mutation. (a) Wild-type sequence in the region of the mutation. The arrow indicates the position of the adenine insertion. (b) Sequence of the 4088insA mutated allele. The arrow indicates the inserted adenine. The frame-shift resulting from the adenine insertion is visible as tangled sequence.

### 5.1.3 Haplotype analysis (I, II)

Three-marker haplotypes consisting of alleles at markers *D13S260*, *D13S1701* and *D13S267*, spanning 1.83 Mb (3.2 cM), were manually constructed for the three *BRCA2* exon 9 999del5 mutation carriers and 8 non-carriers in family OV30. The mutation carriers shared haplotype 166 - 304 - 258. This haplotype was not detected in any of the non-carriers. In family BR17 a DNA sample was available only from the mutation carrier and her haplotype was 168/174 - 292/304 - 252/256. It is likely that the allele 304 of the closest marker to the 999del5 mutation, *D13S1701*, is shared by all mutation carriers in families OV30 and BR17.

In family OV03 with *BRCA2* exon 11 6503delTT mutation the haplotype using markers *D13S260*, *D13S1701* and *D13S267* was constructed for two mutation carriers and two non-carriers. The 6503delTT mutation associated haplotype was 170 - 292 - 244/256 and it was not detected among the non-carriers.

Two-marker haplotype was constructed for six mutation carriers and five non-carriers in family BR09 with the *BRCA2* 4088insA mutation using markers *D13S260* and *D13S267*. All mutation carriers shared the haplotype 166 - 250 and it was not detected in the non-carriers.

Haplotypes in families BR01, OV01 and PS04 carrying *BRCA1* 4216-2ntA→G and 5370 C→T mutations, have been assessed as part of the collaborative analysis of haplotypes and founder effects of *BRCA1* and *BRCA2* families in Finland (Sarantaus et al. 2000).

## **5.2 Autosome-wide scan for linkage disequilibrium-based association with breast cancer (IV)**

### **5.2.1 Allelic association**

In the first-stage scan 366 microsatellite markers were genotyped on 22 chromosomes in 49 breast cancer cases and 50 controls from Eastern Finnish population. In 16 chromosomes altogether 21 markers showed significant difference in allele frequencies when compared between breast cancer cases and controls, with *P*-values 0.003-0.049 in Fisher's exact test, indicating that these markers are in LD with breast cancer (Study IV, Table 1). In addition, two markers associated with breast cancer with borderline significance, with *P*-values 0.054 and 0.057 (Study IV, Table 1). More than one marker on same chromosome was detected to be in LD with breast cancer on 5 chromosomes, but none of these markers were adjacent. On six chromosomes no markers were detected to be in LD with breast cancer. For the second-stage scan 23 loci (21 with the most significant *P*-values and 2 near significant) on 16 chromosomes were selected and altogether 69 additional flanking markers were analysed. Four of the 69 analysed markers showed significant LD with breast cancer with *P*-values 0.013-0.046 (Study IV, Table 1). The breast cancer association was further analysed by single-allele tests for 25 of the 27 significant markers where the difference between cases and controls in the count of the tested allele was  $\geq 10$ . Thirty-nine alleles of 25 markers were tested and 28 alleles showed significant association, two showed borderline significance and 8 were not significant (Study IV, Table 1). The odds ratios indicated 9 alleles possibly being protective (OR 0.013-0.47) and 21 alleles associating with increased breast cancer risk (OR 1.84-12.86) (Study IV, Table 1).

### 5.2.2 Genotype association and Hardy-Weinberg equilibrium (HWE)

Among all analysed markers (366+69) only six markers showed significant difference in genotype frequencies between cases and controls. These were *D3S3053*, *D7S1818*, *D19S178*, *D19S254*, *D22S1177* and *D22S445*, with *P*-values 0.041, 0.014, 0.048, 0.007, 0.040 and 0.010, respectively. In the HWE analysis the genotypes of controls deviated significantly from equilibrium at one marker, *DIS179*, and slightly at *D22S426* (Study IV, Table 1). The cases deviated also at *DIS179* and at *D17S809* with *P*-values ~0 and 0.011, respectively. In the combined analysis of cases and controls together deviation from HWE was detected with markers *DIS179*, *D17S809* and *D22S426* with *P*-values ~0, 0.013 and 0.006, respectively. The substantial deviation at marker *DIS179* in all three tests is probably due to the large number of observed alleles, 15, at this marker.

### 5.2.3 Haplotypes

In the haplotype analysis 76 different marker combinations were tested for two-, three and four-marker haplotypes. Four marker combinations showed significant difference in haplotype frequencies between cases and controls with global test score  $\leq 0.05$  and simulated *P*-values of 0.003-0.021 (Study IV, Table 2). In addition, three other showed borderline significance with global scores 0.077, 0.095 and 0.068 (Study IV, Table 2). With the four significant marker combinations the number of observed haplotypes with frequency of  $\geq 0.5$  % varied from 22 to 45. Among these individual haplotypes four were found more frequently in cases and seven were more frequent in controls (*P*-values 0.007-0.038, simulated *P*-values 0.004-0.030; Study IV, Table 2). The simulations and haplotype frequency estimations for cases and controls separately were not done for the three marker combinations with borderline significance.

The results from allelic association and haplotype analysis suggest that in the Eastern Finnish population breast cancer associated regions locate on chromosomes 3p26, 11q23 and 22q12-q13.

### **5.3 Further refinement of chromosome 22q12-q13 breast cancer-associated region (V)**

#### **5.3.1 SNP allelic and genotype association**

Ten SNPs were genotyped in the 550-kb breast cancer-associated region on chromosome 22q12-13, detected in the autosome-wide scan in study IV. Breast cancer association of the SNPs were tested by comparing the allele and genotype frequencies between 497 breast cancer cases and 458 controls using the entire sample and a subsample that included the cases born in the province of Northern Savo (n=280) and their age and long-term area-of-residence matched controls (n=257). In the entire sample no significant difference were detected in allele frequencies between cases and controls with any of the ten SNPs but a borderline significance was detected for the rare allele carriers (genotypes TC and CC) of rs733655 with an OR of 1.28 (95% CI=0.99-1.65, Table 29). In the subsample two SNPs associated with breast cancer, rs728506 with *P*-value 0.033, and rs733655 with *P* of 0.009 (Table 29). In the subsample also the rs733655 rare allele carriers (genotypes TC and CC) had a significant risk of breast cancer compared to the homozygous common allele carriers (TT) with an OR of 1.90 (95% CI=1.34-2.69). The genotype frequencies differed significantly with rs733655 in the entire sample (*P*=0.044) and the heterozygous genotype TC was the risk-associated genotype with OR 1.39 (95% CI=1.06-1.83, Study V, Table 2). Breast cancer association with rs733655 was detected also in the subsample where the genotype frequencies differed between cases and controls significantly with *P* of 0.0003 and the TC genotype associated with breast cancer risk with an OR of 2.11 (95% CI=1.46-3.05, Study V, Table 3). In the subsample a protective effect associated with rs7285064 heterozygous genotype CT with OR 0.62 (95% CI=0.41-0.96, Study V, Table 3).

#### **5.3.2 Hardy-Weinberg equilibrium**

The deviation of the genotype frequencies from the Hardy-Weinberg equilibrium was tested separately for cases and controls in each sample (entire and subsample). In the controls the rs733655 genotypes were slightly off from HWE in the entire sample and more significantly in the subsample. A borderline deviation was detected among cases

**Table 29.** SNP allele frequencies, allele frequency comparisons and allele specific risks.

| SNP      | $P^*$                      |                        | Allele | Allele Frequency           |          |                        |          | Rare allele carrier risk   |           |                        | Common allele carrier risk |                            |           |                        |           |
|----------|----------------------------|------------------------|--------|----------------------------|----------|------------------------|----------|----------------------------|-----------|------------------------|----------------------------|----------------------------|-----------|------------------------|-----------|
|          | Entire sample <sup>†</sup> | Subsample <sup>§</sup> |        | Entire sample <sup>†</sup> |          | Subsample <sup>§</sup> |          | Entire sample <sup>†</sup> |           | Subsample <sup>§</sup> |                            | Entire sample <sup>†</sup> |           | Subsample <sup>§</sup> |           |
|          |                            |                        |        | Cases                      | Controls | Cases                  | Controls | OR                         | CI        | OR                     | CI                         | OR                         | CI        | OR                     | CI        |
| rs738977 | 0.958                      | 0.296                  | C      | 0.834                      | 0.835    | 0.864                  | 0.84     |                            |           |                        |                            | 0.93                       | 0.44-1.95 | 0.75                   | 0.26-2.13 |
|          |                            |                        | T      | 0.166                      | 0.165    | 0.136                  | 0.16     | 1.00                       | 0.75-1.32 | 0.75                   | 0.51-1.11                  |                            |           |                        |           |
| rs382735 | 0.581                      | 0.837                  | A      | 0.718                      | 0.730    | 0.710                  | 0.717    |                            |           |                        |                            | 0.87                       | 0.54-1.41 | 0.81                   | 0.44-1.47 |
|          |                            |                        | C      | 0.282                      | 0.270    | 0.290                  | 0.283    | 1.05                       | 0.81-1.37 | 0.99                   | 0.70-1.39                  |                            |           |                        |           |
| rs3484   | 0.118                      | 0.620                  | A      | 0.559                      | 0.532    | 0.552                  | 0.536    |                            |           |                        |                            | 1.13                       | 0.82-1.55 | 1.07                   | 0.71-1.63 |
|          |                            |                        | G      | 0.441                      | 0.477    | 0.448                  | 0.464    | 0.78                       | 0.58-1.03 | 0.91                   | 0.62-1.32                  |                            |           |                        |           |
| rs760517 | 0.493                      | 0.894                  | C      | 0.687                      | 0.672    | 0.681                  | 0.677    |                            |           |                        |                            | 1.00                       | 0.65-1.54 | 1.01                   | 0.59-1.73 |
|          |                            |                        | T      | 0.313                      | 0.328    | 0.319                  | 0.323    | 0.89                       | 0.68-1.15 | 0.97                   | 0.69-1.37                  |                            |           |                        |           |
| rs738148 | 0.501                      | 0.170                  | G      | 0.729                      | 0.715    | 0.733                  | 0.694    |                            |           |                        |                            | 1.29                       | 0.80-2.08 | 1.41                   | 0.77-2.57 |
|          |                            |                        | A      | 0.271                      | 0.285    | 0.267                  | 0.306    | 0.96                       | 0.74-1.25 | 0.82                   | 0.58-1.15                  |                            |           |                        |           |
| rs728506 | 0.470                      | <b>0.033</b>           | C      | 0.883                      | 0.872    | 0.904                  | 0.861    |                            |           |                        |                            | 1.33                       | 0.44-3.99 | 1.52                   | 0.34-6.84 |
|          |                            |                        | T      | 0.117                      | 0.128    | 0.096                  | 0.139    | 0.90                       | 0.66-1.23 | 0.62                   | 0.41-0.95                  |                            |           |                        |           |
| rs25095  | 0.202                      | 0.260                  | A      | 0.541                      | 0.511    | 0.558                  | 0.521    |                            |           |                        |                            | 1.23                       | 0.90-1.69 | 1.40                   | 0.91-2.14 |
|          |                            |                        | G      | 0.459                      | 0.489    | 0.442                  | 0.479    | 0.88                       | 0.66-1.18 | 0.91                   | 0.62-1.34                  |                            |           |                        |           |
| rs733655 | 0.254                      | <b>0.009</b>           | T      | 0.721                      | 0.744    | 0.687                  | 0.76     |                            |           |                        |                            | 1.22                       | 0.75-1.98 | 1.18                   | 0.63-2.20 |
|          |                            |                        | C      | 0.279                      | 0.256    | 0.313                  | 0.24     | 1.28                       | 0.99-1.65 | <b>1.90</b>            | <b>1.34-2.69</b>           |                            |           |                        |           |
| rs228941 | 0.292                      | 0.172                  | G      | 0.706                      | 0.728    | 0.698                  | 0.737    |                            |           |                        |                            | 0.92                       | 0.57-1.48 | 0.54                   | 0.26-1.10 |
|          |                            |                        | C      | 0.294                      | 0.272    | 0.302                  | 0.263    | 1.16                       | 0.90-1.50 | 1.17                   | 0.83-1.65                  |                            |           |                        |           |
| rs228942 | 0.163                      | 0.315                  | C      | 0.796                      | 0.821    | 0.791                  | 0.817    |                            |           |                        |                            | 0.88                       | 0.45-1.71 | 0.88                   | 0.34-2.27 |
|          |                            |                        | A      | 0.204                      | 0.179    | 0.209                  | 0.183    | 1.15                       | 0.88-1.50 | 1.23                   | 0.86-1.75                  |                            |           |                        |           |

\* $P$ -value from chisquared Fisher's exact test for the difference in allele frequencies cases vs. controls.

† 497 breast cancer cases and 458 controls.

§ 280 breast cancer cases and 257 controls.

in the subsample in rs738977 and rs733655 genotypes with  $P$ -values 0.052 and 0.082, respectively. (Study V, Tables 2 and 3).

### 5.3.3 Pairwise linkage disequilibrium

Linkage disequilibrium between SNPs was estimated in the entire sample by calculating pairwise  $D'$  values for all 45 possible SNP pairs. In six of the 45 pairs a  $D' \geq 0.5$  were observed (Table 30).  $D'$  was detected to decline as distance between SNPs increased, as was expected. The highest  $D'$ , 0.99, was detected between rs228941 and rs228942 which are in the same gene (*IL2RB*) and only 898 kb apart. Moderate LD with  $D'$  0.49-0.61 was observed between SNPs that are up to 236 kb apart.

**Table 30.**  $D'$  values for pairwise linkage disequilibrium in the entire sample (cases and controls combined). The distance between a SNP pair is shown in kilobases (kb) in the upper half and  $D'$  value in the lower half.

|           | rs738977 | rs3827351 | rs3484 | rs760517 | rs738148 | rs7285064 | rs25095 | rs733655 | rs228941 | rs228942 |
|-----------|----------|-----------|--------|----------|----------|-----------|---------|----------|----------|----------|
| rs738977  |          | 148.3     | 187.9  | 250.1    | 261.6    | 321.1     | 425.1   | 486.1    | 514.8    | 515.7    |
| rs3827351 | 0.56     |           | 39.6   | 101.7    | 113.3    | 172.8     | 276.8   | 337.8    | 366.5    | 367.4    |
| rs3484    | 0.05     | 0.59      |        | 62.1     | 73.6     | 133.1     | 237.2   | 298.2    | 326.9    | 327.7    |
| rs760517  | 0.08     | 0.26      | 0.17   |          | 11.5     | 71.0      | 175.0   | 236.1    | 264.7    | 265.6    |
| rs738148  | 0.12     | 0.04      | 0.10   | 0.57     |          | 59.5      | 163.5   | 224.5    | 253.2    | 254.1    |
| rs7285064 | 0.14     | 0.06      | 0.08   | 0.16     | 0.18     |           | 104.0   | 165.1    | 193.7    | 194.6    |
| rs25095   | 0.04     | 0.08      | 0.12   | 0.15     | 0.06     | 0.25      |         | 61.0     | 89.7     | 90.6     |
| rs733655  | 0.13     | 0.28      | 0.26   | 0.04     | 0.28     | 0.09      | 0.49    |          | 28.7     | 29.6     |
| rs228941  | 0.02     | 0.03      | 0.13   | 0.10     | 0.24     | 0.19      | 0.24    | 0.05     |          | 0.9      |
| rs228942  | 0.23     | 0.08      | 0.02   | 0.07     | 0.24     | 0.61      | 0.01    | 0.02     | 0.99     |          |

### 5.3.4 Haplotypes

Two-marker haplotypes were estimated for the adjacent SNPs that were in LD with  $D' \geq 0.5$ . Altogether five pairs were tested separately in the entire sample and subsample. The global chi-squared  $P$  value and empirical  $P$  were significant for only one marker pair in the subsample, rs25095-rs733655. An individual haplotype AC of this SNP pair is significantly more frequent in cases with empirical  $P$  of 0.003 (Study V, Table 5). In the entire sample this haplotype has a borderline significance with empirical  $P$  0.073, but non-significant global  $P$ . Other haplotypes were not significant.



### 5.3.5 Allelic imbalance analysis

In 37/45 (82%) of the studied breast tumours AI was detected at one or more of the 13 microsatellite marker loci analysed on chromosome 22q12-q13. Five microsatellite loci, *D22S1142*, *D22S924*, *D22S1177*, *D22S445* and *D22S279*, showed significant AI (Table 31).

**Table 31.** Allelic imbalance detected on chromosome 22q12-q13.

| Marker          | Location (Mb) | AI    |                |
|-----------------|---------------|-------|----------------|
|                 |               | n*    | % <sup>†</sup> |
| <i>D22S277</i>  | 34.596        | 5/35  | 14             |
| <i>D22S1142</i> | 34.598        | 12/36 | <b>33</b>      |
| <i>D22S924</i>  | 34.956        | 15/33 | <b>45</b>      |
| <i>D22S283</i>  | 35.075        | 11/40 | 28             |
| <i>D22S426</i>  | 35.348        | 2/12  | 17             |
| <i>D22S1177</i> | 35.588        | 12/22 | <b>55</b>      |
| <i>IL2RB</i>    | 35.871        | 10/40 | 25             |
| <i>D22S445</i>  | 35.890        | 9/26  | <b>35</b>      |
| <i>D22S1156</i> | 36.706        | 2/25  | 13             |
| <i>D22S272</i>  | 37.410        | 8/32  | 25             |
| <i>D22S284</i>  | 38.641        | 9/35  | 26             |
| <i>D22S279</i>  | 39.347        | 7/18  | <b>39</b>      |
| <i>D22S270</i>  | 41.378        | 1/30  | 3              |

\* Number of tumours showing AI/Total number of informative tumours.

<sup>†</sup> AI percentage.

## 6. DISCUSSION

### 6.1 High-risk susceptibility genes in Eastern Finnish breast and ovarian cancer families (I-III)

#### 6.1.1 Eastern Finnish *BRCA1* and *BRCA2* mutation frequency is not different from other parts of Finland (I)

Five different *BRCA1* and *BRCA2* gene mutations were found in the Eastern Finnish population; 4216-2nt A→G in exon 12, and 5370 C→T in exon 20 of *BRCA1* gene, and 999del5 in exon 9, 4088insA in exon 11 and 6503delTT in exon 11 of *BRCA2*. These mutations account for 19.4 % of the 36 families studied. *BRCA1* mutations were detected in 8.3 % (3/36) and *BRCA2* mutations in 11.1 % (4/36) of the families. In the previous studies conducted in Northern or Southern Finnish families the corresponding figures were 7 % and 10 % for *BRCA1*, and 6 % and 11% for *BRCA2* (Huusko et al. 1998, Vehmanen et al. 1997a). Thus, the frequencies that we observed do not differ significantly from those previously reported in Finland.

Some mutations in the present study may have remained undetected since the sensitivity of mutation detection methods is not complete. It is also possible to miss some mutations due to phenocopies as in some families DNA from only one breast or ovarian cancer case was available for mutation screening. However, missing many mutations is unlikely because 17 previously detected Finnish *BRCA1* and *BRCA2* mutations were tested, and the mutation frequencies are similar to those previously reported in Finnish breast/ovarian cancer families. Also, large genomic rearrangements involving *BRCA* genes have not been found in the Finnish population (Lahti-Domenici et al. 2001, Laurila et al. 2005, Karhu et al. 2006).

In addition to truncating mutations three polymorphisms were detected in the *BRCA1* gene and seven in *BRCA2* gene. All ten polymorphisms are known and have been found in other populations. *BRCA2* Asn372His has been associated with an increased risk of breast cancer although in a very large collaborative sample set the association was not confirmed (Healey et al. 2000a, Breast Cancer Association Consortium 2006). In Kuopio Breast Cancer Project sample material the association was not significant

(Healey et al. 2000a). Some evidence was however reported of the 372His homozygotes in the youngest age group having an increased risk of breast cancer (Breast Cancer Association Consortium 2006). Increased risk of ovarian cancer has also been detected among Asn372His carriers (Auranen et al. 2003). None of the other nine polymorphisms has been reported to associate with breast/ovarian cancer risk. Thus, it is unlikely that these polymorphisms contribute significantly to risk of familial breast cancer in our sample material.

In general, the mutations in the present study were found in families with several cancer cases. However, in many families with breast and ovarian cancer a mutation was not found, and two mutations were found in families with only two cases of breast cancer. The *BRCA1* 5370 C→T mutation was found in a family in which only two cases of breast cancer (second-degree relatives) were known, but additionally many cases of other cancers. The mutation in this family was found from a patient with melanoma, and DNA from the breast cancer cases was not available for testing. The *BRCA1* 4216-2nt A→G mutation was also detected in a family where the index case had breast cancer at the age of 36 and her mother had endometrial and breast cancers but no other cancer cases were known in the family. This demonstrates that very strict criteria, i.e. three first-degree relatives with breast/ovarian cancer, for *BRCA1/2* mutation testing are not applicable in all cases.

In the present study *BRCA* mutations were not detected in any of the six families with ovarian cancer only, which is consistent with the previous findings. In a study of 233 unselected Finnish ovarian cancer patients *BRCA1/2* mutations were detected in 5.6 % of the patients (Sarantaus et al. 2001b). Mutations were present in 13 families and only two of them did not have a history of breast cancer in the family, in addition to ovarian cancer (Sarantaus et al. 2001b). Even though the number of families with ovarian cancer only is low in the present study, it supports the previous conclusion that ovarian cancer patients who have also been diagnosed with breast cancer or have family history of breast or breast/ovarian cancer could benefit from referral to genetic counselling and *BRCA* mutation testing (Sarantaus et al. 2001b).


### 6.1.2 Limited *BRCA1* and *BRCA2* mutation spectrum in Eastern Finland (I)

Although the *BRCA1* and *BRCA2* mutation frequencies observed in the present study correspond to those previously found in Finland, the mutation spectrum, however, is different. We detected only two of the 15 previously reported *BRCA1* and two of the 11 reported *BRCA2* mutations (Vehmanen et al. 1997a, Vehmanen et al. 1997b, Roth et al. 1998, Huusko et al. 1998, Sarantaus et al. 2000, Syrjäkoski et al. 2000, Sarantaus et al. 2001a, Sarantaus et al. 2001b, Pääkkönen et al. 2001, Eerola et al. 2001, Vahteristo et al. 2001a, Syrjäkoski et al. 2004a). The Eastern Finnish (Northern Savo) mutation spectrum resembles that of the Northern part of the country where only a few *BRCA1/2* mutations have been found (Huusko et al. 1998). Two of the four recurrent mutations that we detected represent the most frequently observed *BRCA1/2* gene founder mutations in Finland, 4216-2nt A→G in *BRCA1* and 999del5 in *BRCA2*. The other two, *BRCA1* 5370 C→T and *BRCA2* 6503delTT, have been less frequently detected in Finland and have been observed in other countries too. In addition to *BRCA1* 4216-2nt A→G and *BRCA2* 999del5, the three other most frequent Finnish mutations in each gene were not detected in the Eastern Finnish families (*BRCA1* 3604delA, 3744delT/3745delT and 4446C→T, and *BRCA2* 7708C→T, 8555T→G and 9346-2ntA→T). Additionally, we observed the 4088insA mutation in *BRCA2* exon 11. In Finland *BRCA2* 4088insA mutation is unique to the family described in study II. It has been reported in the BIC database to have been found in one Italian ovarian cancer patient but not anywhere else.

However, it should be kept in mind that the 36 families in the present study cover only a portion of the families with hereditary breast and ovarian cancer in the Northern Savo region, and are a portion diagnosed in the 1990's. It is possible that if the study material was larger some of the most prevalent founder mutations would be found, especially *BRCA2* 7708C→T and 9346-2ntA→T, which have been described in Finnish families originating from Northern Savo (Sarantaus et al. 2000).

### 6.1.3 Geography and haplotypes in the *BRCA* mutation-carrier families (I, II)

The *BRCA1* gene 4216-2nt A→G mutation was detected in two families, which are from the Northern part of Central Finland and in Northern Savo region (Figure 14). In the collaborative analysis the origins of the Finnish families with 4216-2nt A→G mutation were observed to cluster in Central Ostrobothnia and further in Central Finland (Sarantaus et al. 2000).



**Figure 14.** Birth places of the oldest generation available in five *BRCA1/2* mutation families. Black circles represent the birth places of *BRCA2* 999del5 mutation family (grandparents of index case in family OV30), black square *BRCA2* 6503delTT (father of index case in family OV03), star *BRCA1* 4216-2nt A→G mutation families (index cases of families BR01 and OV01), diamond *BRCA2* 4088insA (index case of family BR09). Location of Kuopio is shown with a triangle.

The origin of the *BRCA1* 5370 C→T mutation families in Finland seems to be in the most Eastern parts, or in Russia (Sarantaus et al. 2000). The birthplace of the oldest generation known (father of the index case) in the family with *BRCA2* 6503delTT

mutation is also from Eastern Finland, close to Kuopio (Figure 14). The single family carrying *BRCA2* 4088insA mutation originates from Karelia (Figure 14).

The *BRCA2* 999del5 mutation was also detected in two families. In one of the families all four grandparents of the index case could be traced to have been born in a small region in Eastern Finland, communities in Northern Savo and Northern Karelia (Figure 14). In the other family with 999del5 mutation the birth places were not available. A haplotype analysis in Finnish and Icelandic breast cancer families with *BRCA2* 999del5 mutation revealed two distinct haplotypes; a rare one that shared a core-haplotype with the Icelandic haplotype, and another one more frequent in Finland (Barkardottir et al. 2001). Families with this more frequent haplotype originated from two regions in Finland, the other of them in Eastern Finland near Kuopio where the family in the present study also originates.

Our results support the idea of the birth of population islets in Finland during resettlement of the Eastern parts of Finland in the 16th century. Only two of the eight most common Finnish founder mutations were detected in our material. In addition, the novel mutation *BRCA2* 4088insA has not been found in Southern or Northern Finland. Our study agrees with the suggestion that there are only a limited number of *BRCA1/BRCA2* mutations in the Finnish population and the majority of them have been found. The relatively low total number of known *BRCA1/BRCA2* mutations in Finland, including the ones reported in this study, suggests that other possibly moderate risk inducing gene mutations may exist in the Finnish population. Indeed, recently a founder mutation in *PALB2* gene has been found in Finnish breast cancer families, including a family from Kuopio region (Erkko et al. 2007). The knowledge of founder mutations representative of certain geographical regions facilitates the laborious mutation screening of large *BRCA1* and *BRCA2* genes.

#### **6.1.4 A novel Eastern Finnish *BRCA2* mutation and favourable prognosis (II)**

In the present study we have identified a rare 4088insA mutation in the *BRCA2* gene. In addition to the Eastern Finnish family in the present study it has not been detected in other families in Finland. The mutation is associated with high penetrance: all four

female mutation carriers have presented with breast/ovarian cancer and all of them were either bilateral breast cancer or ovarian cancer, or breast and ovarian cancer in the same patient. One of the two male mutation carriers has prostate cancer. The 4088insA mutation seems to be of an auspicious nature as despite of even an advanced stage and poor prognosis assessed by classical factors all affected family members had an excellent response to treatment and a better than expected clinical course. Still, the mutation is persistent as breast cancer occurred in a patient even 25 years after the first (ovarian) cancer. On the other hand, the patient diagnosed with bilateral ovarian cancer of the size of two grape fruits and metastatic spreads, was also cured and is still disease free and alive after nearly 26 years.

An interesting question is whether the mutation itself is associated with good prognosis/outcome or is there another gene that modifies the outcome of the mutation-associated cancers in the family. Also, as *BRCA2* gene functions in double-stranded DNA break (DSDB) repair, cells deficient in functional *BRCA2* are more sensitive to agents that cause DSDB (Patel et al. 1998, Sharan et al. 1997, Abbott et al. 1998). Therefore certain chemotherapeutic drugs and radiation may be especially efficient in *BRCA2*-deficient cancer patients but on the other hand the radiation therapy in mutation carriers may initiate tumorigenesis by causing DSDB that are not subsequently repaired. Indeed, some evidence for *BRCA2*-deficient cancer cells being more sensitive to ionizing radiation, mitoxantrone, amsacrine and etoposide have been presented (Sharan et al. 1997). Also *BRCA1/2*-related breast cancer cases had a better clinical response rate to neoadjuvant chemotherapy than non-carriers (Chappuis et al. 2002). In the family described here the patient with bilateral breast cancer did not receive radiation therapy for the first breast cancer and patient who received radiation therapy for breast cancer had subsequent rectal and ovarian cancers which are unlikely due to scatter dose of radiation to breast. The third breast cancer patient in the family has also received radiation therapy, but only recently, and has not developed bilateral breast cancer. Better response to platinum chemotherapy has been shown among *BRCA1* or *BRCA1/2* mutation heterozygotes but not among *BRCA2* mutation carriers only, compared to non-carriers, and it may have contribution to the improved prognosis (Husain et al. 1998, Cass et al. 2003). Patients in the present study received cisplatinum-based

chemotherapy for ovarian cancer, and one patient received also chemotherapy including mitoxantrone (CMF/CNF) for breast cancer prior to the ovarian cancer. Thus, it can be speculated that the mutation enhances the response to such chemotherapy, improving prognosis.

There is clearly need for collection of data on the effect of individual mutations, as well as of response to treatment among mutation carriers. The 4088insA mutation provides an example of how the knowledge of the effect of an individual mutation may be important in estimating prognosis for mutation carriers. The mutation in a breast/ovarian cancer family may be of auspicious nature and the knowledge of this may provide some relief in the distress for the family members carrying the mutation. In the near future the genetic information of the germ-line of the cancer patient may be important in assessing prognosis of the disease on a wider scale.

## **6.2 Sporadic breast cancer and new low-penetrance susceptibility genes in Eastern Finland (IV, V)**

### **6.2.1 Autosomal-wide scan for LD-based breast cancer association (IV)**

Our scan was conducted in order to gain knowledge of possible genetic breast cancer risk factors on population level. We selected the cases that did not have a strong family history of breast cancer from a small geographical region for optimal homogeneous gene pool and as small effective number of unrelated founders as possible (Kruglyak 1999). Also when the cases are a homogeneous group from small geographical region the subset of genetic risk factors they carry could be also homogeneous and the number of them reduced. We aimed at identifying low-penetrance risk factors for breast cancer which does not show up clearly in families. The genetic diversity of this population is believed to be lower than that of the Finnish population in general and this feature has been the basis for mapping several genes and even genes for complex diseases (Ekelund et al. 2000). Due to the nature of our population this study falls in between traditional LD study and family linkage study. Being a young isolated population this group of Eastern Finns could be seen as a sample set in between a family set and a group of non-relatives. When this work was initiated (1996) it was not recognised as it is according to



the current knowledge that it could be useful to have family material even in association studies (Antoniou and Easton 2003). Also, at the time of the sample selection the contribution of high-penetrance genes, e.g. *BRCA1* and *BRCA2*, was not known and we therefore chose non-familial breast cancer cases in order to identify low-penetrance susceptibility genes. Very recently though, it has been shown that the same low-penetrance risk factors could be identified in a sample set of sporadic breast cancer and among cases with a strong family history of breast cancer (Hunter et al. 2007, Easton et al. 2007).

Of the 366 markers in the first-stage scan 59 % were spaced by  $\leq 10$  cM, 30 % by 11-15 cM and 11 % were spaced by 16-20 cM. Less than half of the screened chromosomes harboured one or two gaps of 18-20 cM. However, in this approximately 20 generations old founder population relatively few meioses have occurred and linkage disequilibrium should extend beyond the distances between adjacent markers in the first and second-stage scans in general (Ophoff et al. 2002). In a sub-isolate of Kuusamo in the late-settlement region up to 50 % of microsatellites showed LD at distances 3.5-7.5 cM and 30 % at distances  $>7.5$  cM (Varilo et al. 2003). Our study population was selected from a small rural area in the late-settlement region and could possess the genetic characteristics more similar to the Kuusamo population than the late-settlement population in general. It is also worth noting that with a 10-cM map LD between marker and risk allele needs to extend 5 cM, as our scan is not for LD between markers but between marker and risk allele that lies somewhere between two markers spaced by 10 cM in average. LD between SNP markers extends a much shorter distance. A study comparing the extent of LD between SNP markers with minor allele frequency  $>0.1$  in four populations of European ancestry but different demographic histories, Ashkenazim, Africaners, East Anglian British and Eastern Finns, found that the average LD was similar in all populations (Dunning et al. 2000). However, there are regions in the genome where the LD varies between populations, e.g. the Eastern Finnish population showed somewhat higher LD in regions on chromosome 22 (Dunning et al. 2000). Also, in the Eastern Finnish population LD was detected at marker pairs 200-500 kb apart while the other three studied populations did not show LD different from the baseline at these distances (Dunning et al. 2000).

#### **6.2.1.1 Candidate-genes for further studies on breast cancer association (IV)**

In the present study altogether 27 markers in 16 chromosomes showed association with breast cancer in the two-staged autosome-wide scan using 435 microsatellite markers. In the haplotype analysis three chromosomal regions, 3p26, 11q23 and 22q12-q13 were further suggested as candidate locations for breast cancer-associated genes.

Testing 366 markers at the 0.05 significance level is predicted to give 18 positive associations by chance alone ( $366 \times 0.05$ ). Since we found 21 positives in the first stage, it is reasonable to propose that some of these represent true associations. Haplotype analysis using global-score method and markers in and around associated chromosomal sites revealed four positive regions, which fits well with this assumption. Estimating the number of possible false positive results in the second-stage scan is more complicated because the markers were selected next to the markers that are in LD with breast cancer and therefore are not independent. The Bonferroni correction was not used here as it is very conservative and we view these results as a guide for further investigation and not statistically definite significant associations. Thus we wanted to follow up all possible associations in the second-stage scan. Also, it is unclear how the adjusting for multiple comparisons should be best done in genome-wide screenings (Ophoff et al. 2002) and a method that identifies false positives has not been introduced. When studying complex diseases it is expected that there will be multiple loci affecting the trait and a very strict control of the number of false positives (such as the Bonferroni method) will result in a considerable loss of power to identify secondary signals (Ophoff et al. 2002).

We are aware that all possible breast cancer-associated loci may not have been recognised in our screen due to the longer distance between markers in some chromosomal regions and a limited sample set. Also, at the time when this scan was initiated the exact physical locations of some markers were not available. As a consequence some markers used in the second-stage scan are not as close to markers that were in LD in the first-stage scan as initially expected. This, in turn, could lead to false negative results. Therefore, a denser marker map and a bigger sample set could probably have led to an improved result of the scan. Considering all that, it may not be best to limit the search in the three chromosomal regions, 3p26, 11q23 and 22q12-q13, identified in the haplotype analysis but to investigate possible candidate-genes near all

individual markers that showed breast cancer association. Some of the genes are proposed to be cancer-related and some of them could be candidates for breast cancer-associated genes based on the proposed function of the gene product. In the 560-kb haplotype region on 3p26 there are three genes, two of which are not very well characterised or properly named. Evidence from loss of heterozygosity (LOH) studies suggests a few candidate tumour suppressor genes lie on chromosome 3p (Yang et al. 2002c) but none of them locates in the 560-kb region of the significant haplotype found in our scan. Interestingly though, a linkage analysis conducted on Finnish prostate cancer families identified 3p26 as a locus for hereditary prostate cancer gene (Schleutker et al. 2003, Rökman et al. 2005). On 11q23 the 900-kb haplotype region includes 13 named genes and four hypothetical genes. Chromosome 11q21-24 region is also known for LOH in several different cancers including breast cancer. In the 550-kb region of the two significant haplotypes on 22q13.1 locates seven named genes, two non-characterised and two hypothetical genes. Another 16 interesting genes locate near the markers with breast cancer association in other chromosomes.

Recently three genome-wide association studies for breast cancer risk alleles have been conducted (Easton et al. 2007, Stacey et al. 2007, Hunter et al. 2007). Two of them used sporadic case-control material (Hunter et al. 2007, Stacey et al. 2007), whereas in the third GWS case material selected for a strong family history of breast cancer was used in the initial scan (Easton et al. 2007). Regardless, these GWSs identified the strongest associations to *FGFR2* (Easton et al. 2007, Hunter et al. 2007) and *TNRC9* (Easton et al. 2007, Stacey et al. 2007) genes, although each of them identified also additional associations that were not detected in other studies. This indicates that low-penetrance risk factors for breast cancer can be detected using GWS. On the other hand even the new-generation GWSs using SNPs do not find all risk factors and the identification of the causal variant can be troublesome. In our microsatellite scan we detected 27 positive associations. One of these (*D2S434*) locates near to (670 kb) the chromosome 2q35 association found in the GWS conducted by Stacey and coworkers (2007). A borderline significant ( $P=0.95$ ) marker (*D10S1213*) and an associated allele with it was detected 2 Mb from the described *FGFR2* association, and only 400 kb from the other associated SNP on 10q (rs10510126). Clearly though, it can not be argued to

be the same association without further studies. Other associations detected in our scan were not in the vicinity of the associations described in the more recent GWSs. However, more associations were detected than expected by chance alone and some of these can represent true risk factors. Thus, our scan is also a feasible method for initial screen for identification of potential candidate regions for follow-up studies.

## **6.2.2 Breast cancer association on chromosome 22q12-q13 (V)**

### **6.2.2.1 *TMPRSS6* as a candidate-gene (V)**

In the present study, the most promising breast cancer association was detected with a SNP rs733655 in *TMPRSS6* gene. We observed an increased risk associating with the heterozygous genotype TC of rs733655 and with the rare allele C carriers but not at the homozygous genotype CC carriers. This may be partly explained by too small number of observed rare homozygotes alone to show significant increase in risk and a larger sample set might solve this problem.

In the controls the rs733655 genotypes deviated slightly from HWE whereas the cases were in equilibrium. However, in the RefSNP European test population sample the HWE probability is 0.05 indicating that in the normal situation the genotype frequencies deviate from those expected under HWE ([http://www.ncbi.nlm.nih.gov/SNP/snp\\_ref.cgi?rs=733655](http://www.ncbi.nlm.nih.gov/SNP/snp_ref.cgi?rs=733655)). The rs733655 allele frequencies that we observed among controls are also similar to those expected in the NCBI RefSNP database. The allele frequencies among our cases differ from those expected (RefSNP) as genotypes that we observed do not deviate from HWE. If the number of controls by genotype is replaced by the numbers expected under HWE, the difference in genotype frequencies between cases and controls still remains significant in the subsample (which includes only cases born in the province of Northern Savo and their age and long-term area-of-residence matched controls) ( $P=0.013$ ). Also, the risk for the heterozygous genotype TC remains significant in the subsample, although reducing from 2.11 to 1.67 (CI 1.17-2.39). When validating the SNPs for this study we sequenced 20 samples for rs733655 and no sequence variation other than rs733655 was observed in the region of the designed TaqMan primer and probe sequences.

Genotyping assays were optimized and allele calling was unambiguous as allele controls were used. Also 141 samples from altogether six plates were re-genotyped, confirming the original genotype and thus, a genotyping error is also excluded. Therefore, we consider that the detected difference between the case and control populations is real. Although it is common to exclude markers which deviate from HWE from association studies it is possible to miss true risk factors by doing so. When genotyping artefacts are excluded it leaves open the possibility that common variants that show departure from HWE are subject to selection, acting either directly or indirectly via selective sweep across other variants in LD with them (Baynes et al. 2007).

*TMPRSS6* gene encodes a membrane-bound serine proteinase 6 called matriptase-2. Matriptase-2 is a member of a family of type II transmembrane serine proteinases which have possible role in cancer development. Matriptase-2 has the ability to degrade extracellular matrix components, suggesting that it may participate in some of the matrix-degrading processes occurring in both normal and pathological conditions, including cancer progression (Velasco et al. 2002, Netzel-Arnett et al. 2003). As matriptase-2 is only recently discovered, little is known about the physiological function. Matriptase-2 is expressed in normal breast tissue and the expression is elevated in breast cancer (invasive ductal carcinoma) (Overall et al. 2004). *TMPRSS6* has also been detected to be mutated in breast carcinomas (Sjöblom et al. 2006). Thus, the possibility of matriptase-2 involvement in cancer progression is supported and *TMPRSS6* is a potential novel candidate for breast cancer risk affecting gene. More specific investigation for the association between matriptase-2 and breast cancer, as well as the identification of the causal genetic variant is needed. To resolve the global importance of *TMPRSS6* in genetic risk of breast cancer the association has to be tested in other populations also.

According to the power calculations our sample set of 497 cases and 458 controls had >95 % power to detect a risk allele that is in perfect LD with the marker allele ( $D'=1$ ) and has a relative risk of 2. The power to detect a risk allele that is not in perfect LD with the marker allele ( $D'=0.5$ ) and has a relative risk of 2 varies between 44-74 % with the ten SNPs. In the stratified sample the power for the detected risk (2.11 for the

TC and 1.16 for TT genotype) with SNP rs733655 is 83 %. Within a short distance (1 kb) we detected highly significant LD ( $D'=1$ ) and even at distances longer than 200 kb  $D'$  is  $\geq 0.5$  at some instances. It is presumable that LD does not remain at the constant level across the whole 516-kb region and some associations may therefore have been missed.

Our negative results do not rule out association involving other nearby SNPs and positive results do not necessarily indicate the discovery of the causal SNP but a marker in LD with a true causal SNP located some distance away.

#### **6.2.2.2 Other genes on chromosome region 22q12-q13 (V)**

Another SNP, rs7285064 in *CSF2RB* gene, showed moderate association with breast cancer. In the subsample the heterozygous genotype associated with a reduced risk of breast cancer indicating a protective effect. Further evidence concerning the importance of this gene in breast cancer is necessary. Other genes in the studied region have not been reported to associate with (breast) cancer.

#### **6.2.2.3 Allelic imbalance on chromosome region 22q12-q13 (V)**

Matriptase-2 is expressed in normal breast tissue and the expression is elevated in breast cancer (invasive ductal carcinoma) (Overall et al. 2004), which points out that matriptase-2 is expected not to be a tumor suppressor. Our AI analysis provided further support to this since AI was not detected in the vicinity of *TMPRSS6* gene. This also indicates that the detected AI seems to be a separate event from the original breast cancer association, i.e. hereditary risk factor, while AI may indicate the involvement of a nonhereditary factor. However, in line with other studies (Allione et al. 1998, Iida et al. 1998, Bryan et al. 2000, Castells et al. 2000, Hirano et al. 2001) the observed AI on the 22q12-q13 region does not exclude existence of a tumor suppressor, other than *TMPRSS6*.

## 7. SUMMARY AND CONCLUSIONS

Breast cancer is the most common cancer in women and ovarian cancer is also among the most common cancers. The objective of this study was to study the contribution of known high-risk breast and ovarian cancer susceptibility genes, *BRCA1* and *BRCA2*, in Eastern Finnish families. The second objective was to find chromosomal regions and genes in them that associate with the risk of (sporadic) breast cancer.

This study demonstrated that:

1. Germ-line mutations in *BRCA1* and *BRCA2* genes account for 19.4 % of the studied Eastern Finnish breast/ovarian cancer families. The observed Eastern Finnish mutation spectrum differs from those observed in the Northern and Southern parts of the country. Only two of the eight most common Finnish founder mutations were detected in the Eastern Finnish families. In addition, one of the mutations has not been found elsewhere in Finland (*BRCA2* 4088insA). The results add to the evidence of the geographical differences in distribution of Finnish *BRCA1* and *BRCA2* mutations. This screen also provides further evidence for the presumption that the majority of Finnish *BRCA1/BRCA2* founder mutations have been found and that the proportion of *BRCA1/BRCA2* mutations in Finnish breast/ovarian cancer families is around 20 %.
2. A *BRCA2* mutation 4088insA, novel to Finland, was found in one family of Eastern Finnish origin. The mutation is highly penetrant; four of the six carriers are affected with breast and ovarian, bilateral breast or bilateral ovarian cancer and one with prostate cancer. Despite of the high penetrance this mutation seems to be associated with better than expected outcome. The mutation positive breast/ovarian cancer patients in this family had an excellent response to treatment even when prognosis assessed by classical factors was poor.
3. The *BRCA1* exon 12 4216-2nt A→G substitution mutation was confirmed to disrupt the splice acceptor dinucleotide AG and to form another splice acceptor

dinucleotide. Wrong splicing site leads to inclusion of an extra nucleotide and a frameshift in translation that causes a premature translation termination and a truncated protein product.

4. In the first stage of the autosome-wide linkage disequilibrium-based screen 21 STR markers on 16 chromosomes showed initial association with breast cancer. A further analysis with additional markers and haplotypes restricted breast cancer associations to three chromosomal regions, 3p26, 11q23 and 22q13.
5. A 516-kb region on chromosome 22q12-q13 was further refined using 10 SNP markers. Significant association was detected with one SNP located in the intronic sequence of *TMPRSS6* gene encoding matriptase-2. The heterozygous genotype and the minor allele were associated with increased risk of breast cancer. In the AI analysis of 22q12-q13 abundant AI was detected more centromeric than previously reported, further supporting the existence of a tumour suppressor gene or genes in this region.

In conclusion, mutations detected in the known high-risk and high-penetrance breast and ovarian cancer susceptibility genes, *BRCA1* and *BRCA2*, account for approximately 20 % of the familial breast/ovarian cancer in Eastern Finland. This corresponds to the observed contribution of these genes in families in other parts of our country. In our search for additional breast cancer risk associated genes we have found an interesting new candidate, *TMPRSS6*, which warrants additional association and functional studies. In addition, chromosomal regions 3p26 and 11q23 may harbour breast cancer-associated genes.



## 8. REFERENCES

- Abbott DW, Freeman ML, Holt JT. Double-strand break repair deficiency and radiation sensitivity in BRCA2 mutant cancer cells. *J Natl Cancer Inst* 1998;90:978-85.
- Abecasis GR, Noguchi E, Heinzmann A, et al. Extent and distribution of linkage disequilibrium in three genomic regions. *Am J Hum Genet* 2001;68:191-7.
- Adjei AA, Weinshilboum RM. Catecholesterol sulfation: Possible role in carcinogenesis. *Biochem Biophys Res Commun* 2002;292:402-8.
- Adnane J, Gaudray P, Dionne CA, et al. BEK and FLG, two receptors to members of the FGF family, are amplified in subsets of human breast cancers. *Oncogene* 1991;6:659-63.
- Ahn J, Gammon MD, Santella RM, et al. Myeloperoxidase genotype, fruit and vegetable consumption, and breast cancer risk. *Cancer Res* 2004;64:7634-9.
- Akiyama T, Sudo C, Ogawara H, Toyoshima K, Yamamoto T. The product of the human c-erbB-2 gene: a 185-kilodalton glycoprotein with tyrosine kinase activity. *Science* 1986;232:1644-6.
- Alberg AJ, Daudt A, Huang HY, et al. N-acetyltransferase 2 (NAT2) genotypes, cigarette smoking, and the risk of breast cancer. *Cancer Detection Prev* 2004;28:187-93.
- Allinen M, Huusko P, Mäntyniemi S, Launonen V, Winqvist R. Mutation analysis of the CHK2 gene in families with hereditary breast cancer. *Br J Cancer* 2001;85:209-12.
- Allione F, Eisinger F, Parc P, Noguchi T, Sobol H, Birnbaum D. Loss of heterozygosity at loci from chromosome arm 22q in human sporadic breast carcinomas. *Int J Cancer* 1998;75:181-6.
- Al-Zahrani A, Sandhu MS, Luben RN, et al. IGF1 and IGFBP3 tagging polymorphisms are associated with circulating levels of IGF1, IGFBP3 and risk of breast cancer. *Hum Mole Genet* 2006;15:1-10.
- Ambrosone CB, Freudenheim JL, Graham S, et al. Cytochrome P4501A1 and glutathione S-transferase (M1) genetic polymorphisms and postmenopausal breast cancer risk. *Cancer Res* 1995;55:3483-5.
- Ambrosone CB, Freudenheim JL, Graham S, et al. Cigarette smoking, N-acetyltransferase 2 genetic polymorphisms, and breast cancer risk. *J Am Med Assoc* 1996;276:1494-501.
- Ambrosone CB, Freudenheim JL, Thompson PA, et al. Manganese superoxide dismutase (MnSOD) genetic polymorphisms, dietary antioxidants, and risk of breast cancer. *Cancer Res* 1999;59:602-6.
- Angele S, Romestaing P, Moullan N, et al. ATM haplotypes and cellular response to DNA damage: association with breast cancer risk and clinical radiosensitivity. *Cancer Res* 2003;63:8717-25.

Anglian Breast Cancer Study Group. Prevalence and penetrance of BRCA1 and BRCA2 mutations in a population-based series of breast cancer cases. *Br J Cancer* 2000;83:1031-8.

Antoniou AC, Gayther SA, Stratton JF, Ponder BA, Easton DF. Risk models for familial ovarian and breast cancer. *Genet Epidemiol* 2000;18:173-90.

Antoniou A, Pharoah PDP, McMullan G, et al. A comprehensive model for familial breast cancer incorporating BRCA1, BRCA2 and other genes. *Br J Cancer* 2002;86:76-83.

Antoniou AC, Easton DF. Polygenic inheritance of breast cancer: implications for design of association studies. *Genet Epidemiol* 2003;25:190-202.

Antoniou A, Pharoah PDP, Narod S, et al. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected case series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet* 2003;72:1117-30.

Antoniou AC, Pharoah PDP, Narod S, et al. Breast and ovarian cancer risks to carriers of the BRCA1 5382insC and 185delAG and BRCA2 6174delT mutations: a combined analysis of 22 population based studies. *J Med Genet* 2005;42:602-3.

Antoniou AC, Easton DF. Models of genetic susceptibility to breast cancer. *Oncogene* 2006;25:5898-905.

Anzick SL, Kononen J, Walker RL, et al. AIB1, a steroid receptor coactivator amplified in breast and ovarian cancer. *Science* 1997;277:965-8.

Arai H, Miyamoto K, Taketani Y, et al. a vitamin D receptor gene polymorphism in the translation initiation codon: effect on protein activity and relation to bone mineral density in Japanese women. *J Bone Miner Res* 1997;12:915-21.

Ardlie K, Liu-Cordero SN, Eberle MA, et al. Lower-than-expected linkage disequilibrium between tightly linked markers in humans suggests a role for gene conversion. *Am J Hum Genet* 2001;69:582-9.

Athma P, Rappaport R, Swift M. Molecular genotyping shows that ataxia-telangiectasia heterozygotes are predisposed to breast cancer. *Cancer Genet Cytogenet* 1996;92:130-4.

Auranen A, Spurdle AB, Chen X, et al. BRCA2 Arg372His polymorphism and epithelial ovarian cancer risk. *Int J Cancer* 2003;103:427-30.

Auranen A, Song H, Waterfall C, et al. Polymorphisms in DNA repair genes and epithelial ovarian cancer risk. *Int J Cancer* 2005;117:611-8.

Bakkenist CJ, Kastan MB. DNA damage activates ATM through intermolecular autophosphorylation and dimer dissociation. *Nature* 2003;421:499-506.

Baldwin RL, Nemeth E, Tran H, et al. BRCA1 promoter region hypermethylation in ovarian carcinoma: A population-based study. *Cancer Res* 2000;60:5329-33.

Balz V, Prissack HB, Bier H, Bojar H. Analysis of BRCA1, TP53, and TSG101 germline mutations in German breast and/or ovarian cancer families. *Cancer Genet Cytogenet* 2002;138:120-7.

Banin S, Moyal L, Shieh SY, et al. Enhanced phosphorylation of p53 by ATM in response to DNA damage. *Science* 1998;281:1674-7.

Barbacid M. ras Genes. *Am Rev Biochem* 1987;56:779-827.

Barkardottir RB, Sarantaus L, Arason A, et al. Haplotype analysis in Icelandic and Finnish BRCA2 999del5 breast cancer families. *Eur J Hum Genet* 2001;9:773-9.

Barroso E, Milne RL, Fernandez LP, et al. FANCD2 associated with sporadic breast cancer risk. *Carcinogenesis* 2006;27:1930-7.

Bartek J, Falck J, Lucas J. CHK2 kinase - a busy messenger. *Nature Reviews Mol Cell Biol* 2001;2:877-86.

Bartkova J, Horejsi Z, Koed K, et al. DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature* 2005;434:864-70.

Bartsch H, Nair J, Owen RW. Dietary polyunsaturated fatty acids and cancers of the breast and colorectum: emerging evidence for their role as risk modifiers. *Carcinogenesis* 1999;20:2209-18.

Bartsch H, Nair U, Risch A, Rojas M, Wikman H, Alexandrov K. Genetic polymorphism of CYP genes, alone or in combination, as a risk modifier of tobacco-related cancers. *Cancer Epidemiol Biomarkers Prev* 2000;9:3-28.

Basham VM, Pharoah PDP, Healey CS, et al. Polymorphisms in CYP1A1 and smoking: no association with breast cancer risk. *Carcinogenesis* 2001;22:1797-1800.

Baynes C, Healey CS, Pooley KA, et al. Common variants in the ATM, BRCA1, BRCA2, CHEK2 and TP53 cancer susceptibility genes are unlikely to increase breast cancer risk. *Breast Cancer Res* 2007;9:R27.

Baysal BE, DeLoia JA, Willett-Brozick JE, et al. Analysis of CHEK2 gene for ovarian cancer susceptibility. *Gyn Oncol* 2004;95:62-9.

Beckmann MW, Picard F, An HX, et al. Clinical impact of detection of loss of heterozygosity of BRCA1 and BRCA2 markers in sporadic breast cancer. *Br J Cancer* 1996;73:1220-6.

Bekesi G, Kakucs R, Varbiro S, et al. Induced myeloperoxidase activity and related superoxide inhibition during hormone replacement therapy. *BJOG* 2001;108:474-81.

Bell DW, Varley JM, Szydlo TE, et al. Heterozygous germ line hCHK2 mutations in Li-Fraumeni syndrome. *Science* 1999;286:2528-31.

Ben David Y, Cherit A, Hirsch-Yechezkel G, et al. Effect of BRCA mutations on the length of survival in epithelial ovarian tumors. *J Clin Oncol* 2002;20:463-6.

- Benusiglio PR, Lesueur F, Luccarini C, et al. Common ERBB2 polymorphisms and risk of breast cancer in a white population: a case-control study. *Breast Cancer Res* 2005;7:R204-R209.
- Bergman M, Ahnström M, Palmebäck Wegman P, Wingren S. Polymorphism in the manganese superoxide dismutase (MnSOD) gene and risk of breast cancer in young women. *J Cancer Res Clin Oncol* 2005;131:439-44.
- Bergthorsson JT, Jonasdottir A, Johannesdottir G, et al. Identification of a novel splice-site mutation of the BRCA1 gene in two breast cancer families: screening reveals low frequency in Icelandic breast cancer patients. *Hum Mutat* 1998;suppl 1:S195-7.
- Bernstein JL, Thompson WD, Risch N, Holford TR. Risk factors predicting the incidence of second primary breast cancer among women diagnosed with a first primary breast cancer. *Am J Epidemiol* 1992;136:925-36.
- Bernstein L, Henderson BE, Hanisch R, Sullivan-Halley J, Ross RK. Physical exercise and reduced risk of breast cancer in young women. *J Natl Cancer Inst* 1994;86:1403-8.
- Bernstein JL, Teraoka S, Southey MC, et al. Population-based estimates of breast cancer risks associated with ATM gene variants c.7271T>G, and c.1066-6T>G (IVS10-6T>G) from the Breast Cancer Family Registry. *Hum Mut* 2006; 27:1122-8.
- Bertone-Johnson E. Epidemiology of ovarian cancer: a status report. *Lancet* 2005;365:101-2.
- Bianco T, Chenevix-Trench G, Walsh DCA, Cooper JE, Dobrovic A. Tumour-specific distribution of BRCA1 promoter region methylation supports a pathogenetic role in breast and ovarian cancer. *Carcinogenesis* 2000;21:147-51.
- BIC. Breast Cancer Information Core. <http://research.nhgri.nih.gov/bic/>
- Bierie B, Moses HL. TGF $\beta$ : the molecular Jekyll and Hyde of cancer. *Nat Rev Cancer* 2006;6:506-20.
- Bignell G, Micklem G, Stratton MR, Ashworth A, Wooster R. The BRC repeats are conserved in mammalian BRCA2 proteins. *Hum Mol Genet* 1997;6:53-8.
- Blasina A, Van de Weyer I, Laus MC, Luyten WHML, Parker AE, McGowan CH. A human homologue of the checkpoint kinase Cds1 directly inhibits Cdc25 phosphatase. *Curr Biol* 1999;9:1-10.
- Blum M, Demierre A, Grant DM, Heim M, Meyer UA. Molecular mechanisms of slow acetylation of drugs and carcinogens in humans. *Proc Natl Acad Sci USA* 1991;88:5237-41.
- Bochar DA, Wang L, Baniya H, et al. BRCA1 is associated with a human SWI/SNF-related complex: linking chromatin remodeling to breast cancer. *Cell* 2000;102:257-65.
- Bogdanova N, Enßen-Dubrowskaja N, Feshchenko S, et al. Association of two mutations in the CHEK2 gene with breast cancer. *Int J Cancer* 2005;116:263-6.

Borg Å, Sandberg T, Nilsson K, et al. High frequency of multiple melanomas and breast and pancreas carcinomas in CDKN2A mutation positive melanoma families. *J Natl Cancer Inst* 2000;92:1260-6.

Borg Å. Molecular and pathological characterization of inherited breast cancer. *Cancer Biol* 2001;11:375-85.

Bork P, Blomberg N, Nilges M. Internal repeats in the BRCA2 protein sequence. *Nat Genet* 1996;13:22-3.

Bosron WF, Li TK. Genetic polymorphism of human liver alcohol and aldehyde dehydrogenases, and their relationship to alcohol metabolism and alcoholism. *Hepatology* 1986;6:502-10.

Botha JL, Bray F, Sankila R and Parkin DM. Breast cancer incidence and mortality trends in 16 European countries. *Eur J Cancer* 2003;39:1718-29.

Botstein D, Risch N. discovering genotypes underling human phenotypes: past successes for mendelian disease, future approaches for complex disease. *Nat Genet* 2003;33 Suppl:228-37.

Boyd M, Harris F, McFarlane R, Davidson HR, Black DM. A human BRCA1 gene knockout. *Science* 1995;375:541-2.

Boyd J, Sonoda Y, Federici MG, et al. Clinicopathologic features of BRCA-linked and sporadic ovarian cancer. *JAMA* 2000;283:2260-5.

Boyd J. Specific keynote: Hereditary ovarian cancer: what we know. *Gynecol Oncol* 2003;88:S8-S10.

Breast Cancer Association Consortium. Commonly studied single-nucleotide polymorphisms and breast cancer: Results from the Breast Cancer Association Consortium. *J Natl Cancer Inst* 2006;98:1382-96.

Breast Cancer Linkage Consortium. Pathology of familial breast cancer: differences between breast cancers in carriers of BRCA1 or BRCA2 mutations and sporadic cases. *Lancet* 1997;349:1505-10.

Breast Cancer Linkage Consortium. Cancer risks in BRCA2 mutation carriers. *J Natl Cancer Inst* 1999;91:1310-6.

Brekelmans CTM, Seynaeve C, Menke-Pluymers M, et al. Survival and prognostic factors in BRCA1-associated breast cancer. *Ann Oncol* 2006;17:391-400.

Brown AL, Lee C-H, Schwarz JK, Mitiku N, Piwnica-Worms H, Chung JH. A human Cds1-related kinase that functions in the cellular response to DNA damage. *Proc Natl Acad Sci* 1999;96:3745-50.

Bryan EJ, Thomas NA, Palmer K, Dawson E, Englefield P, Campbell IG. Refinement of an ovarian cancer tumour suppressor gene locus on chromosome arm 22q and mutation analysis of CYP2D6, SREBP2 and NAGA. *Int J Cancer* 2000;87:798-802.

Brzovic PS, Meza J, King MC, Klevit RE. The cancer-predisposing mutation C61G disrupts homodimer formation in the NH2-terminal BRCA1 RING finger domain. *J Biol Chem* 1998;273:7795-9.

Burwinkel B, Wirtenberger M, Klaes R, et al. Association of NCOA3 polymorphisms with breast cancer risk. *Clin Cancer Res* 2005;11:2169-74.

Butcher NJ, Ilett KE, Minchin RF. Functional polymorphisms of the human arylamine JV-acetyltransferase type 1 gene caused by C190T and G560A mutations. *Pharmacogenetics* 1998;8:67-72.

Börresen AL, Andersen TI, Tretli S, Heiberg A, Møller P. Breast cancer and other cancers in Norwegian families with ataxia-telangiectasia. *Genes Chromosomes Cancer* 1990;2:339-40.

Börresen AL, Andersen TI, Garber J, et al. Screening for germ line TP53 mutations in breast cancer patients. *Cancer Res* 1992;52:3234-6.

Cai Q, Shu XO, Jin F, et al. Genetic polymorphisms in the estrogen receptor alpha gene and risk of breast cancer. Results from the Shanghai Breast Cancer Study. *Cancer Epidemiol Biomarkers Prev* 2003;12:853-9.

Cai X, Shu XO, Wen W, et al. Genetic polymorphism in the manganese superoxide dismutase gene, antioxidant intake, and breast cancer risk: results from the Shanghai Breast Cancer Study. *Breast Cancer Res* 2004;6:R647-R655.

Caligo MA, Agata S, Aceto G, et al. The CHEK2 c.1100delC mutation plays an irrelevant role in breast cancer predisposition in Italy. *Hum Mutat* 2004;24:100-1.

Cancer Committee of the International Federation of Gynaecology and Obstetrics. Staging announcement: FIGO Cancer Committee. *Gynecol Oncol* 1986;25:383-5.

Canman CE, Lim DS, Cimprich KA, et al. Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. *Science* 1998;281:1677-9.

Canzian F, Salovaara R, Hemminki A, et al. Semiautomated assessment of loss of heterozygosity and replication error in tumours. *Cancer Res* 1996;56:3331-7.

Canzian F, McKay JD, Cleveland RJ, et al. Genetic variation in the growth hormone synthesis pathway in relation to circulating insulin-like growth factor-I, insulin-like growth factor binding protein-3, and breast cancer risk: Results from the European prospective investigation into cancer and nutrition study. *Cancer Epidemiol Biomarkers Prev* 2005;14:2316-25.

Canzian F, McKay JD, Cleveland RJ, et al. Polymorphisms of genes coding for insulin-like growth factor 1 and its major binding proteins, circulating levels of IGF-1 and IGFBP-3 and breast cancer risk: results from the EPIC study. *Br J cancer* 2006;94:299-307.

Capon DJ, Chen EY, Levinson AD, Seeburg PH, Goeddel DV. Complete nucleotide sequences of the T24 human bladder carcinoma oncogene and its normal homologue. *Nature* 1983;302:33-7.

Cardon LR, Bell JI. Association study designs for complex diseases. *Nat Rev Genet* 2001;2:91-9.

Carroll BT, Couch FJ, Rebbeck TR, Weber BL. Polymorphisms in PTEN in breast cancer families. *J Med Genet* 1999;36:94-6.

Carter CL, Corle DK, Micozzi MS, Schatzkin A, Taylor PR. A prospective study of the development of breast cancer in 16,692 women with benign breast disease. *Am J Epidemiol* 1988;128:467-77.

Cass I, Baldwin RL, Varkey T, Moslehi R, Narod SA, Karlan BY: Improved survival in women with BRCA-associated ovarian carcinoma. *Cancer* 2003;97:2187-95.

Castells A, Gusella JF, Ramesh V, Rustgi AK. A region of deletion on chromosome 22q13 common to human breast and colorectal cancers. *Cancer Res* 2000;60:2836-9.

Cavalieri EL, Stack DE, Devanesan PD, et al. Molecular origin of cancer: catechol estrogen-3,4-quinones as endogenous tumor initiators. *Proc Natl Acad Sci USA* 1997;94:10937-42.

Cebrian A, Pharoah PD, Ahmed S, et al. Tagging single-nucleotide polymorphisms in antioxidant defense enzymes and susceptibility to breast cancer. *Cancer Res* 2006;66:1225-33.

Chacko P, Rajan B, Joseph T, Mathew BS, Pillai MR. Polymorphisms in DNA repair gene XRCC1 and increased genetic susceptibility to breast cancer. *Breast Cancer Res Treat* 2005;89:15-21.

Chakravarti A. population genetics-making sense out of sequence. *Nat Genet* 1999;21 suppl:56-60.

Chang CS, Kokontis J, Liao ST. Molecular cloning of human and rat complementary DNA encoding androgen receptors. *Science* 1988;240:324-6.

Chang JC, Hilsenbeck SG, Fuqua SAW. Genomic approaches in the management and treatment of breast cancer. *Br J Cancer* 2005;92:618-24.

Chapman MS, Verma IM. Transcriptional activation by BRCA1. *Nature* 1996;382:678-9.

Chappuis PO, Goffin J, Wong N, et al. A significant response to neoadjuvant chemotherapy in BRCA1/2 related breast cancer. *J Med Genet* 2002;39:608-10.

Chaturvedi P, Eng WK, Zhu Y, et al. Mammalian Chk2 is a downstream effector of the ATM-dependent DNA damage checkpoint pathway. *Oncogene* 1999;18:4047-54.

Chehab NH, Malikzay A, Appel M, Halazonetis TD. Chk2/hCds1 functions as a DNA damage checkpoint in G1 by stabilizing p53. *Genes Devel* 2000;14:278-88.

CHEK2 Breast Cancer Case-Control Consortium. CHEK2\*1100delC and susceptibility to breast cancer: A collaborative analysis involving 10,860 breast cancer cases and 9,065 controls from 10 studies. *Am J Hum Genet* 2004;74:1175-82.

CHEK2-Breast Cancer Consortium. Low-penetrance susceptibility to breast cancer due to CHEK2\*1100delC in noncarriers of BRCA1 or BRCA2 mutations. *Nat Genet* 2002;31:55-9.

Chen LZ, Harris PC, Apostolou S, et al. A refined physical map of the long arm of human chromosome 16. *Genomics* 1991;10:308-12.

Chen Y, Chen CF, Riley DJ, et al. Aberrant subcellular localization of BRCA1 in breast cancer. *Science* 1995;270:789-91.

Chen CF, Li S, Chen Y, Chen PL, Sharp ZD, Lee WH. The nuclear localization sequences of the BRCA1 protein interact with the importin- $\alpha$  subunit of the nuclear transport signal receptor. *J Biol Chem* 1996;271:32863-8.

Chen J, Silver DP, Walpita D, et al. Stable interaction between the products of the BRCA1 and BRCA2 tumor suppressor genes in mitotic and meiotic cells. *Molecular Cell* 1998a;2:317-28.

Chen PL, Chen CF, Chen Y, Xiao J, Sharp ZD, Lee WH. The BRC repeats in BRCA2 are critical for RAD51 binding and resistance to methyl methanesulfonate treatment. *Proc Natl Acad Sci USA* 1998b;95:5287-92.

Chen J, Lindblom A. Germline mutation screening of the STK11/LKB1 gene in familial breast cancer with LOH on 19p. *Clin Genet* 2000;57:394-7.

Chen WY, Bertone-Johnson, Hunter DJ, Willett WC, Hankinson SE. Associations between polymorphisms in the vitamin D receptor and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2005;14:2335-9.

Chenevix-Trench G, Spurdle AB, Gatei M, et al. Dominant negative ATM mutations in breast cancer families. *J Natl Cancer Inst* 2002;94:205-15.

Choi JY, Lee KM, Park SK, et al. Genetic polymorphisms of SULT1A1 and SULT1E1 and the risk and survival of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2005;14:1090-5.

Clarke RB, Anderson E, Howell A. Steroid receptors in human breast cancer. *Trends Endocrinol Metab* 2004;15:316-23.

Claus EB, Schildkraut JM, Thompson WD, Risch NJ. The genetic attributable risk of breast and ovarian cancer. *Cancer* 1996;77:2318-24.

Claus EB, Schildkraut J, Iversen ES Jr, et al. Effect of BRCA1 and BRCA2 on the association between breast cancer risk and family history. *J Natl Cancer Inst* 1998;90:1824-9.

Colin EM, Weel AE, Uitterlinden AG, et al. Consequences of vitamin D receptor gene polymorphisms for growth inhibition of cultures human peripheral blood mononuclear cells by 1, 25-dihydroxyvitamin D3. *Clin Endocrinol* 2000;52:211-6.

Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53 297 women with breast cancer and 100 239 women without breast cancer from 54 epidemiological studies. *Lancet* 1996;347:1713-27.



Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52 705 women with breast cancer and 108 411 women without breast cancer. *Lancet* 1997;350:1047-59.

Collaborative Group on Hormonal Factors in Breast Cancer. Familial breast cancer: Collaborative reanalysis of individual data from 52 epidemiological studies including 58,209 women with breast cancer and 101,986 women without the disease. *Lancet* 2001;358:1389-99.

Collins N, McManus R, Wooster R, et al. Consistent loss of the wild-type allele in breast cancers from a family linked to the BRCA2 gene on chromosome 13q12-13. *Oncogene* 1995;10:1673-5.

Collins N, Wooster R, Stratton MR. Absence of methylation of CpG dinucleotides within the promoter of the breast cancer susceptibility gene BRCA2 in normal tissues and in breast and ovarian cancers. *Br J Cancer* 1997;76:1150-6.

Conrad DF, Jakobsson M, Coop G, et al. A worldwide survey of haplotype variation and linkage disequilibrium in the human genome. *Nat Genet* 2006;38:1251-60.

Cornelis RS, Neuhausen SL, Johansson O, et al. High allele loss rates at 17q12-q21 in breast and ovarian tumors from BRCA1-linked families. The Breast Cancer Linkage Consortium. *Genes Chromosomes Cancer* 1995;13:203-10.

Cortez D, Wang Y, Qin J, Elledge SJ. Requirement of ATM-dependent phosphorylation of Brcal in the DNA damage response to double-strand breaks. *Science* 1999;286:1162-6.

Couch FJ, Farid LM, DeShano ML, et al. BRCA2 germline mutations in male breast cancer cases and breast cancer families. *Nat Genet* 1996;13:123-5.

Coussens L, Yang-Feng TL, Liao YC, et al. tyrosine kinase receptor with extensive homology to EGF receptor shares chromosomal location with neu oncogene. *Science* 1985;230:1132-9.

Cox DG, Kraft P, Hankinson SE, Hunter DJ. Haplotype analysis of common variants in the BRCA1 gene and risk of sporadic breast cancer. *Breast Cancer Res* 2005a;7:R171-R175.

Cox DG, Hankinson SE, Hunter DJ. No association between BRCA2 N372H and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2005b;14:1353-4.

Cox DG, Hankinson SE, Hunter DJ. The erbB2-/HER2/neu receptor polymorphism Ile655Val and breast cancer risk. *Pharmacogenet Genomics* 2005c;15:447-50.

Cox DG, Hankinson SE, Hunter DJ. Polymorphisms of the AURKA (STK15/Aurora kinase) gene and breast cancer risk (United States). *Cancer Cause Control* 2006;17:81-3.

Cox A, Dunning AM, Garcia-Closas M, et al. A common variant in CASP8 is associated with breast cancer risk. *Nat Genet* 2007;39:352-8.

Csokay B, Tihomirova L, Stengrevics A, Sinicka O, Olah E. Strong founder effects in BRCA1 mutation carrier breast cancer patients from Latvia. *Hum Mut* 1999; Mutation in Brief #258.

Cui Y, Shu XO, Cai Q, et al. Association of breast cancer risk with a common functional polymorphism (Asp327Asn) in the sex hormone-binding globulin gene. *Cancer Epidemiol Biomarkers Prev* 2005;14:1096-101.

Cybulski C, Gorski B, Huzarski T, et al. CHEK2 is a multiorgan cancer susceptibility gene. *Am J Hum Genet* 2004;75:1131-5.

Cybulski C, Gorski B, Huzarski T, et al. CHEK2-positive breast cancers in young Polish women. *Clin Cancer Res* 2006;12:4832-5.

Cybulski C, Wokolorczyk D, Huzarski T, et al. A deletion in CHEK2 of 5,395 bp predisposes to breast cancer in Poland. *Breast Cancer Res Treat* 2007;102:119-22. Epub 2006 Aug 8.

Daly MJ, Rioux JD, Schaffner SF, Hudon TJ, Lander ES. High-resolution haplotype structure in the human genome. *Nat Genet* 2001;29:229-32.

D'Andrea AD, Grompe M. The Fanconi anaemia/BRCA pathway. *Nat Rev Cancer* 2003;3:23-34.

Dasika GK, Lin S-CJ, Zhao S, Sung P, Tomkinson A, Lee E Y-HP. DNA damage-induced cell cycle checkpoints and DNA strand break repair in development and tumorigenesis. *Oncogene* 1999;18:7883-99.

Davies AA, Masson JY, McIlwraith MJ, et al. Role of BRCA2 in control of the RAD51 recombination and DNA repair protein. *Mol Cell* 2001;7:273-82.

Davis DL, Bradlow HL, Wolff M, Woodruff T, Hoel DG, Anton-Culver H. Medical hypothesis: Xenoestrogens as preventable causes of breast cancer. *Environ Health Perspect* 1993;101:372-7.

Dawson E, Abecasis GR, Bumpstead S, et al. A first-generation linkage disequilibrium map of human chromosome 22. *Nature* 2002;418:544-8.

de Bock GH, Schutte M, Krol-Warmerdam EM, et al. Tumour characteristics and prognosis of breast cancer patients carrying the germline CHEK2\*1100delC variant. *J Med Genet* 2004;10:731-5.

de Jong MM, Nolte IM, te Meerman GJ, et al. Genes other than BRCA1 and BRCA2 involved in breast cancer susceptibility. *J Med Genet* 2002;39:225-42.

de Jong MM, Nolte IM, de Vries, et al. The HLA class III subregion is responsible for an increased breast cancer risk. *Hum Mol Genet* 2003;12:2311-9.

de la Chapelle, Wright FA. Linkage disequilibrium mapping in isolated populations: The example of Finland revisited. *Proc Natl Acad Sci USA* 1998;95:12416-23.

Debniak T, Gorski B, Huzarski T, et al. a common variant of CDKN2A (p16) predisposes to breast cancer. *J Med Genet* 2005;42:763-5.

Deguchi T, Mashimo M, Suzuki T. Correlation between acetylator phenotypes and genotypes of polymorphic arylamine N-acetyltransferase in human liver. *J Biol Chem* 1990;265:12757-60.

Deguchi T. Sequences and expression of alleles of polymorphic arylamine N-acetyltransferase of human liver. *J Biol Chem* 1992;267:18140-7.

Derynck R, Akhurst RJ, Balmain A. TGF- $\beta$  signaling in tumour suppression and cancer progression. *Nat Genet* 2001;29:117-29.

Devlin B, Risch N. A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics* 1995;29:311-22.

Dobrovic A, Simpfendorfer D. Methylation of the BRCA1 gene in sporadic breast cancer. *Cancer Res* 1997;57:3347-50.

Duell EJ, Millikan RC, Pittman GS, et al. Polymorphisms in the DNA repair gene XRCC1 and breast cancer. *Cancer Epidemiol Biomarkers Prev* 2001;10:217-22.

Dufault MR, Betz B, Wappenschmidt B, et al. Limited relevance of the CHEK2 gene in hereditary breast cancer. *Int J Cancer* 2004;110:320-5.

Dunning AM, Chiano M, Smith NR, et al. Common BRCA1 variants and susceptibility to breast and ovarian cancer in the general population. *Hum Mol Genet* 1997;6:285-9.

Dunning AM, Healey CS, Pharoah PD, et al. No association between a polymorphism in the steroid metabolism gene CYP17 and risk of breast cancer. *Br J Cancer* 1998;77:2045-7.

Dunning AM, Healey CS, Pharoah PDP, Teare MD, Ponder BAJ, Easton DF. A systematic review of genetic polymorphisms and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 1999a;8:843-54.

Dunning AM, McBride S, Gregory J, et al. No association between androgen or vitamin D receptor gene polymorphisms and risk of breast cancer. *Carcinogenesis* 1999b;20:2131-5.

Dunning AM, Durocher F, Healey CS, et al. The extent of linkage disequilibrium in four populations with distinct demographic histories. *Am J Hum Genet* 2000;67:1544-54.

Dunning AM, Ellis PD, McBride S, et al. A transforming growth factor $\beta$ 1 signal peptide variant increases secretion in vitro and is associated with increased incidence of invasive breast cancer. *Cancer Res* 2003;63:2610-5.

Dunning AM, Dowsett M, Healey CS, et al. Polymorphisms associated with circulating sex hormone levels in postmenopausal women. *J Natl Cancer Inst* 2004;94:936-45.

Dupont WD, Page DL. Risk factors for breast cancer in women with proliferative breast disease. *N Engl J Med* 1985;312:146-51.

Easton DF, Bishop DT, Ford D, Crockford GP. Genetic linkage analysis in familial breast and ovarian cancer: results from 214 families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 52:678-701, 1993.

Easton DF. Cancer risks in A-T heterozygotes. *Int J Radiat Biol* 1994;66(6Suppl):S177-S182.

Easton DF, Ford D, Bishop DT. Breast and ovarian cancer incidence in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Am J Hum Genet* 1995;56:265-71.

Easton DF, Steele L, Fields P, et al. Cancer risks in two large breast cancer families linked to BRCA2 on chromosome 13q12-13. *Am J Hum Genet* 1997;61:120-8.

Easton DF. How many more breast cancer predisposition genes are there? *Breast Cancer Res* 1999;1:1-4.

Easton DF, Poley KA, Dunning AM, et al. Genome-wide association study identifies novel breast cancer susceptibility loci. *Nature* 2007;447:1087-93.

Eerola H, Vahteristo P, Sarantaus L, et al. Survival of breast cancer patients in BRCA1, BRCA2, and non-BRCA1/2 breast cancer families: a relative survival analysis from Finland. *Int J Cancer* 2001;93:368-72.

Egan KM, Cai Q, Shu XO, et al. Genetic polymorphisms in GSTM1, GSTP1, and GSTT1 and the risk for breast cancer: results from the Shanghai Breast Cancer Study and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2004;13:197-204.

Egan KM, Thompson PA, Titus-Ernstoff L, Moore JH, Ambrosone CB. MnSOD polymorphism and breast cancer in a population-based case-control study. *Cancer Letters* 2003;199:27-33.

Einarsdottir K, Rylander-Rudqvist T, Humphreys K, et al. CYP17 polymorphism in relation to breast cancer risk: a case-control study. *Breast Cancer Res* 2005;7:R890-R896.

Ekelund J, Lichtermann D, Hovatta I, et al. Genome-wide scan for schizophrenia in the Finnish population: evidence for a locus on chromosome 7q22. *Hum Mol Genet* 2000;9:1049-57.

Elledge S. Cell cycle checkpoints: preventing an identity crisis. *Science* 1996;274:1664-72.

Endogenous Hormones and Breast Cancer Collaborative Group. Endogenous sex hormones and breast cancer in postmenopausal women: Reanalysis of nine prospective studies. *J Natl Cancer Inst* 2002;94:606-16.

Eng C, Schneider K, Fraumeni JF Jr, Li FP. Third international workshop on collaborative interdisciplinary studies on p53 and other predisposing genes in Li-Fraumeni syndrome. *Cancer Epidemiol Biomarkers Prev* 1997;6:379-83.

Eng C. PTEN: One gene, many syndromes. *Hum Mut* 2003;22:183-98.

Enmark E, Peltö-Huikko M, Grandien K, et al. Human estrogen receptor  $\beta$ -gene structure, chromosomal localization, and expression pattern. *J Clin Endocrinol Metab* 1997;82:4258-65.

Erkko H, Xia B, Nikkilä J, et al. A recurrent mutation in PALB2 in Finnish cancer families. *Nature* 2007;446:316-9.

Ernster L, Navazio F. Soluble diaphorase in animal tissues. *Acta Chem Scand* 1958;12:595.

Ernster L. DT diaphorase. *Methods Enzymol* 1967;10:309-17.

Esteller M, Silva JM, Dominguez G, et al. Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J Natl Cancer Inst* 2000;92:564-9.

Evans DA, Manley KA, McKusick VA. Genetic control of isoniazid metabolism in man. *Br Med J* 1960;5197:485-91.

Ewart-Toland A, Dai Q, Gao YT, et al. Aurora-A/STK15 T+91A is a general low penetrance cancer susceptibility gene: a meta-analysis of multiple cancer types. *Carcinogenesis* 2005;26:1368-73.

Ewertz M, Duffy SW, Adami HO, et al. Age at parity and risk of breast cancer: a meta-analysis of 8 studies from the Nordic countries. *Int J Cancer* 1990;46:597-603.

Fackenthal JD, Marsh DJ, Richrdsen AL, et al. Male breast cancer in Cowden syndrome patients with germline PTEN mutations. *J Med Genet* 2001;38:159-64.

Falck J, Mailand N, Syljuåsen R, Bartek J, Lukas J. The ATM-Chk2-Cdc25A checkpoint pathway guards against radioresistant DNA synthesis. *Nature* 2001a ;410:842-7.

Falck J, Lukas C, Protopopova, Lukas J, Selivanova G, Bartek J. Functional impact of concomitant versus alternative defects in the Chk2-p53 tumour suppressor pathway. *Oncogene* 2001b;20:5503-10.

Feigelson HS, Coetzee GA, Kolonel LN, Ross RK, Henderson BE. A polymorphism in the CYP17 gene increases the risk of breast cancer. *Cancer Res* 1997;57:1063-5.

Feron VJ, Til HP, de Vrijer F, Woutersen RA, Cassee FR, van Bladeren PJ. Aldehydes: occurrence, carcinogenic potential, mechanism of action and risk assessment. *Mutat Res* 1991;259:363-85.

Ferrara N, Davis-Smyth T. The biology of vascular endothelial growth factor. *Endocr Rev* 1997;18:4-25.

Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med* 2003;9:669-76.

Figueiredo JC, Knight JA, Briollais L, Andrulis IL, Ozelik H. Polymorphisms XRCC1-R399Q and XRCC3-T241M and the risk of breast cancer at the Ontario Site of the Breast cancer Family Registry. *Cancer Epidemiol Biomarkers Prev* 2004;13:583-91.

Finnish Cancer Registry. <http://cancerregistry.fi/>.

Finnish Cancer Registry - Institute for Statistical and Epidemiological Cancer Research: Cancer Incidence in Finland 1998 and 1999. Cancer Statistics of the National Research and Development Centre for Welfare and Health. Cancer Society of Finland Publication No. 63. Helsinki 2002.

Finnish Cancer Registry. Cancer Statistics for Finland. August 30, 2004. <http://cancerregistry.fi/>.

- Firgaira FA, Seshadri R, McEvoy CRE, et al. HRAS1 rare minisatellite alleles and breast cancer in Australian women under age forty years. *J Natl Cancer Inst* 1999;91:2107-11.
- FitzGerald MG, Bean JM, Hedge SR, et al. Heterozygous ATM mutations do not contribute to early onset of breast cancer. *Nat Genet* 1997;15:207-10.
- FitzGerald MG, Marsh DJ, Wahrer D, et al. Germline mutations in PTEN are an infrequent cause of genetic predisposition to breast cancer. *Oncogene* 1998;17:727-31.
- Fleishman SJ, Schlessinger J, Ben-Tal N. A putative molecular-activation switch in the transmembrane domain of erbB2. *Proc Natl Acad Sci USA* 2002;99:15937-40.
- Fletcher O, Johnson N, Palles C, et al. Inconsistent association between the STK15 F31I genetic polymorphism and breast cancer risk. *J Natl Cancer Inst* 2006;98:1014-8.
- Fodor FH, Weston A, Bleiweiss IJ, et al. Frequency and carrier risk associated with common BRCA1 and BRCA2 mutations in Ashkenazi Jewish breast cancer patients. *Am J Hum Genet* 1998;63:45-51.
- Folsom AR, Kaye SA, Prineas RJ, Potter JD, Gapstur SM, Wallace RB. Increased incidence of carcinoma of the breast associated with abdominal adiposity in postmenopausal women. *Am J Epidemiol* 1990;131:794-803.
- Ford D, Easton DF, Bishop DT, Narod SA, Goldgar DE. Risks of cancer in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Lancet* 1994;343:692-5.
- Ford D, Easton DF, Stratton M, et al. Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families. *Am J Hum Genet* 1998;62:676-89.
- Foulkes WD, Wong N, Brunet JS, et al. Germ-line BRCA1 mutation is an adverse prognostic factor in Ashkenazi Jewish women with breast cancer. *Clin Cancer Res* 1997;3:2465-9.
- Frank TS, Deffenbaugh AM, Reid JE, et al. Clinical characteristics of individuals with germline mutations in BRCA1 and BRCA2: Analysis of 10,000 individuals. *J Clin Oncol* 2002;20:1480-90.
- Frank B, Bermejo JL, Hemminki K, et al. Re: Association of a common variant of CASP8 gene with reduced risk of breast cancer. *J Natl Cancer Inst* 2005;97:1012-3.
- Freedman ML, Penney KL, Stram DO, et al. Common variation in BRCA2 and breast cancer risk: a haplotype-based analysis in the multiethnic cohort. *Hum Mol Genet* 2004;13:2431-41.
- Freedman ML, Penney KL, Stram DO, et al. A haplotype-based case-control study of BRCA1 and sporadic breast cancer risk. *Cancer Res* 2005; 65:7516-22.
- Freudenheim JL, Ambrosone CB, Moysich KB, et al. Alcohol dehydrogenase 3 genotype modification of the association of alcohol consumption with breast cancer. *Cancer Causes Control* 1999;10:369-77.

Friedberg EC. How nucleotide excision repair protects against cancer. *Nature Rev Cancer* 2001;1:22-33.

Friedman LS, Ostermeyer EA, Szabo CI, et al. Confirmation of BRCA1 by analysis of germline mutations linked to breast and ovarian cancer in ten families. *Nat Genet* 1994;8:399-404.

Friedman LS, Gayther SA, Kurosaki T, et al. Mutation analysis of BRCA1 and BRCA2 in a male breast cancer population. *Am J Hum Genet* 1997;60:313-9.

Friedman E, Bruchim RB, Kruglikova A, et al. Double heterozygotes for the Ashkenazi founder mutations in BRCA1 and BRCA2 genes. *Am J Hum Genet* 1998;63:1224-7.

Friedrichsen DM, Malone KE, Doody DR, Daling JR, Ostrander EA. Frequency of CHEK2 mutations in a population based case-control study of breast cancer in young women. *Breast cancer Res* 2004;6:R629-R635.

Fu YP, Yu JC, Cheng TC, et al. Breast cancer risk associated with genotypic polymorphism of the nonhomologous end-joining genes: a multigenic study on cancer susceptibility. *Cancer Res* 2003;63:2440-6.

Fuks F, Milner J, Kouzarides T. BRCA2 associates with acyltransferase activity when bound to P/CAF. *Oncogene* 1998;17:2531-4.

Futreal PA, Liu Q, Shattuck-Eidens D, et al. BRCA1 mutations in primary breast and ovarian carcinomas. *Science* 1994;266:120-3.

Försti A, Angelini S, Festa F, et al. Single nucleotide polymorphisms in breast cancer. *Oncol Rep* 2004;11:917-22.

Gabriel SB, Schaffner SF, Nguyen H, et al. The structure of haplotype blocks in the human genome. *Science* 2002;296:2225-9.

Ganguly A, Rck MJ, Prockop DJ. Conformation-sensitive gel electrophoresis for rapid detection of single-base differences in double-stranded PCR products and DNA fragments: Evidence for solvent-induced bends in DNA heteroduplexes. *Proc Natl Acad Sci USA* 1993;90:10325-9.

Garber JE, Goldstein AM, Kantor AF, Dreyfus MG, Fraumeni JF Jr, Li FP. follow-up study of twenty-four families with Li-Fraumeni syndrome. *Cancer Res* 1991;51:6094-7.

Garcia-Closas M, Egan KM, Newcomb PA, et al. Polymorphisms in DNA double-strand break repair genes and risk of breast cancer: two population-based studies in USA and Poland, and meta-analyses. *Hum Genet* 2006;119:376-88.

Garcia-Higuera I, Taniguchi T, Ganesan S, et al. Interaction of the Fanconi anemia proteins and BRCA1 in a common pathway. *Molecular Cell* 2001;7:249-62.

Garfinkel L, Boffetta P, Stellman SD. Alcohol and breast cancer: a cohort study. *Prev Med* 1988;17:686-93.

- Gatti RA, Tward A, Concannon P. Cancer risk in ATM heterozygotes: a model of phenotypic and mechanistic differences between missense and truncating mutations. *Mol Genet metab* 1999;68:419-23.
- Gaudet MM, Chanock S, Lissowska J, et al. Comprehensive assessment of genetic variation of catechol-O-methyltransferase and breast cancer risk. *Cancer Res* 2006;66:9781-5.
- Gaudet MM, Egan KE, Lissowska J, et al. Genetic variation in tumor necrosis factor and lymphotoxin-alpha (TNF-LTA) and breast cancer risk. *Hum Genet* 2007;121:483-90.
- Gayther S, Warren W, Mazoyer S, et al. Germline mutations in the BRCA1 gene in breast and ovarian cancer families provide evidence for a genotype-phenotype correlation. *Nat Genet* 1995;11:428-33.
- Gayther SA, Mangion J, Russell P, et al. Variation of risks of breast and ovarian cancer associated with different germline mutations of the BRCA2 gene. *Nat Genet* 1997a;15:103-5.
- Gayther SA, Harrington P, Russell P, Kharkevich G, Garkavtseva RF, Ponder BAJ. Frequently occurring germ-line mutations of the BRCA1 gene in ovarian cancer families from Russia. *Am J Hum Genet* 1997b;60:1239-42.
- Ghimenti C, Sensi E, Presciuttini S, et al. Germline mutations of the BRCA1-associated ring domain (BARD1) gene in breast and breast/ovarian families negative for BRCA1 and BRCA2 alterations. *Gene Chromosome Canc* 2002;33:235-42.
- Giardiello FM, Welsh SB, Hamilton SR, et al. Increased risk of cancer in the Peutz-Jeghers syndrome. *N Engl J Med* 1987;316:1511-4.
- Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. *Gastroenterology* 2000;119:1447-53.
- Ginsburg ES, Walsh BW, Shea BF, Gao X, Gleason RE, Barbieri RL. The effects of ethanol on the clearance of estradiol in postmenopausal women. *Fertil Steril* 1995;63:1227-30.
- Glatt H, Boeing H, Engelke CEH, et al. Human cytosolic sulphotransferases: genetics, characteristics, toxicological aspects. *Mutat Res* 2001;482:27-40.
- Globocan 2002, <http://www-dep.iarc.fr/>
- Gold B, Kalush F, Bergeron J, et al. Estrogen receptor genotypes and haplotypes associated with breast cancer risk. *Cancer Res* 2004;64:8891-900.
- Goldstein DB. Islands of linkage disequilibrium. *Nat Genet* 2001;29:109-11.
- Goode EL, Dunning AM, Kuschel B, et al. Effect of germ-line genetic variation on breast cancer survival in a population-based study. *Cancer Res* 2002a;62:3052-7.
- Goode EL, Ulrich CM, Potter JD. Polymorphisms in DNA repair genes and associations with cancer risk. *Cancer Epidemiol Biomarkers Prev* 2002b;11:1513-30.



Gorski B, Byrski T, Huzarski T, et al. Founder mutations in the BRCA1 gene in Polish families with breast-ovarian cancer. *Am J Hum Genet* 2000;66:1963-8.

Gram IT, Braaten T, Terry PD, et al. Breast cancer risk among women who start smoking as teenagers. *Cancer Epidemiol Biomarkers Prev* 2005;14:61-6.

Gras E, Cortes J, Diez O, et al. Loss of heterozygosity on chromosome 13q12-q14, BRCA-2 mutations and lack of BRCA-2 promoter hypermethylation in sporadic epithelial ovarian tumors. *Cancer* 2001;92:787-95.

Greenman C, Stephens P, Smith R, et al. Patterns of somatic mutation in human cancer genomes. *Nature* 2007;446:153-8.

Gretarsdottir S, Thorlacius S, Valgarsdottir R, et al. BRCA2 and p53 mutations in primary breast cancer in relation to genetic instability. *Cancer Res* 1998;58:859-62.

Grzeschik KH. Assignment of a structural gene for a fourth human diaphorase (DIA4) to chromosome 16 in man-mouse somatic cell hybrids. *Hum Genet* 1980;53:189-93.

Gudmundsson J, Johannesdottir G, Arason A, et al. Frequent occurrence of BRCA2 linkage in Icelandic breast cancer families and segregation of a common BRCA2 haplotype. *Am J Hum Genet* 1996;58:749-56.

Haber JE. Partners and pathways repairing a double-strand break. *Trend Genet* 2000;16:259-64.

Hadsell DL, Bonnette SG. IGF and insulin action in the mammary gland: Lessons from transgenic and knockout models. *J Mammary Gland Biol Neoplasia* 2000;5:19-30.

Haiman CA, Hankinson SE, Spiegelman D, et al. Polymorphic repeat in AIB1 does not alter breast cancer risk. *Breast Cancer Res* 2000;2:378-85.

Haiman CA, Stram DO, Cheng I, et al. Common genetic variation at PTEN and risk of sporadic breast and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2006;15:1021-5.

Hall JM, Lee MK, Newman B, et al. Linkage of early-onset familial breast cancer to chromosome 17q21. *Science* 1990;250:1684-9.

Han W, Kang D, Lee JE, et al. A haplotype analysis of HER-2 gene polymorphisms: Association with breast cancer risk, HER-2 protein expression in the tumor, and disease recurrence in Korea. *Clin Cancer Res* 2005;11:4775-82.

Hankinson SE, Willett WC, Colditz GA, et al. Circulating concentrations of insulin-like growth factor I and risk of breast cancer. *Lancet* 1998;351:1393-6.

Harrela M, Koistinen H, Kaprio J, et al. Genetic and environmental components of interindividual variation in circulating levels of IGF-I, IGF-II, IGFBP-1, and IGFBP-3. *J Clin Invest* 1996;98:2612-5.

Hartwell LH, Weinert T. Checkpoints: controls that ensure the order of cell cycle events. *Science* 1989;246:629-34.

Hartwell L. Defects in a cell cycle checkpoint may be responsible for the genomic instability of cancer cells. *Cell* 1992;71:543-6.

Hartwell, Kastan. Cell cycle control and cancer. *Science* 1994;266:1821-8.

Healey CS, Dunning AM, Teare DM, et al. A common variant in BRCA2 is associated with both breast cancer risk and prenatal viability. *Nat Genet* 2000a;26:362-4.

Healey CS, Dunning AM, Durocher F, et al. Polymorphisms in the human aromatase cytochrome P450 gene (CYP19) and breast cancer risk. *Carcinogenesis* 2000b;21:189-93.

Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res* 2006;12:3209-15.

Heck KE, Pamuk ER. Explaining the relation between education and postmenopausal breast cancer. *Am J Epidemiol* 1997;145:366-72.

Hedenfalk I, Duggan D, Chen Y, et al. Gene-expression profiles in hereditary breast cancer. *New Engl J Med* 2001;344:539-48.

Heikkinen K, Rapakko K, Karppinen SM, et al. RAD50 and NBS1 are breast cancer susceptibility genes associated with genomic instability. *Carcinogenesis* 2006;27:1593-9.

Helzlsouer KJ, Selmin O, Huang HY, et al. Association between glutathione S-transferase M1, P1, and T1 genetic polymorphisms and development of breast cancer. *JNCI* 1998;90:512-8.

Hemminki A, Markie D, Tomlinson I, et al. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* 1998;391:184-7.

Hengartner MO. The biochemistry of apoptosis. *Nature* 2000;407:770-76.

Hickman D, Risch A, Camilleri JP, Sim E. Genotyping human polymorphic arylamine N-acetyltransferase: identification of new slow allotypic variants. *Pharmacogenetics* 1992;2:217-26.

Hilton JL, Geisler JP, Rathe JA, Hattermann-Zogg MA, DeYoung B, Buller RE. Inactivation of BRCA1 and BRCA2 in ovarian cancer. *J Natl Cancer Inst* 2002;94:1396-1406.

Hines LM, Hankinson SE, Smith-Warner SA, et al. A prospective study of the effect of alcohol consumption and ADH3 genotype on plasma steroid hormone levels and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2000;9:1099-105.

Hirano A, Emi M, Tsuneizumi M, et al. Allelic losses of loci at 3p25.1, 8p22, 13q12, 17p13.3, and 22q13 correlate with postoperative recurrence in breast cancer. *Clin Cancer Res* 2001;7:876-82.

Hirao A, Cheung A, Duncan G, et al. Chk2 is a tumor suppressor that regulates apoptosis in both an ataxia telangiectasia mutated (ATM)-dependent and an ATM-independent manner. *Molecular and Cellular Biology* 2002;22:6521-32.

Hirota T, Kunitoku N, Sasayama T, et al. Aurora-A and an interacting activator, the LIM protein Ajuba, are required for mitotic commitment in human cells. *Cell* 2003;114:585-98.

Hoeijmakers JHJ. Genome maintenance mechanisms for preventing cancer. *Nature* 2001;411:366-74.

Hogervorst FBL, Cornelis RS, Bout M, et al. Rapid detection of BRCA1 mutations by the protein truncation test. *Nat Genet* 1995;10:208-12.

Holt JT, Thompson ME, Szabo C, et al. Growth retardation and tumour inhibition by BRCA1. *Nat Genet* 1996;12:298-302.

Howe GR, McLaughlin J. Breast cancer mortality between 1950 and 1987 after exposure to fractionated moderate-dose-rate ionizing radiation in the Canadian fluoroscopy cohort study and a comparison with breast cancer mortality in the atomic bomb survivors study. *Radiat Res* 1996;145:694-707.

Howlett NG, Taniguchi T, Olson S, et al. Biallelic inactivation of BRCA2 in Fanconi anemia. *Science* 2002;297:606-9.

Huang CS, Chern HD, Chang KJ, Cheng CW, Hsu SM, Shen CY. Breast cancer risk associated with genotype polymorphism of the estrogen-metabolizing genes CYP17, CYP1A1, and COMT: a multigenic study on cancer susceptibility. *Cancer Res* 1999;59:4870-5.

Hughes NC, Janezic SA, McQueen KL, et al. Identification and characterization of variant alleles of human acetyltransferase NAT1 with defective function using p-aminosalicylate as an in-vivo and in-vitro probe. *Pharmacogenetics* 1998;8:55-66.

Hughes DJ, Ginolhac SM, Coupier I, et al. Breast cancer risk in BRCA1 and BRCA2 mutation carriers and polyglutamine repeat length in the AIB1 gene. *Int J Cancer* 2005;117:230-3.

Hughes-Davies L, Huntsman D, Ruas M, et al. EMSY links the BRCA2 pathway to sporadic breast and ovarian cancer. *Cell* 2003;115:523-35.

Hunter DJ, Willett WC. Diet, body size and breast cancer. *Epidemiol Rev* 1993;15:110-32.

Hunter DJ, Hankinson SE, Hough H, et al. A prospective study of NAT2 acetylation genotype, cigarette smoking, and risk of breast cancer. *Carcinogenesis* 1997;18:2127-32.

Hunter DJ, Kraft P, Jacobs KB, et al. A genome-wide association study identifies alleles in FGFR2 associated with risk of sporadic postmenopausal breast cancer. *Nat Genet* 2007; published online May 27, doi:10.1038/ng2075.

Huo X, Hu Z, Zhai X, et al. Common non-synonymous polymorphisms in the BRCA1 associated RING domain (BARD1) gene are associated with breast cancer susceptibility: a case-control analysis. *Breast Cancer Res Treat* 2006;102:329-37.

Husain A, He G, Venkatraman ES, Spriggs DR. BRCA1 up-regulation is associated with repair-mediated resistance to cis-diamminedichloroplatinum (II). *Cancer Res* 1998;58:1120-3.

Huusko P, Pääkkönen K, Launonen V, et al. Evidence of founder mutations in Finnish BRCA1 and BRCA2 families. *Am J Hum Genet* 1998;62:1544-8.

Huusko P, Castren K, Launonen V, et al. Germ-line TP53 mutations in Finnish cancer families exhibiting features of the Li-Fraumeni syndrome and negative for BRCA1 and BRCA2. *Cancer Genet Cytogenet* 1999;112:9-14.

Huzarski T, Cybulski C, Domagala W, et al. Pathology of breast cancer in women with constitutional CHEK2 mutations. *Breast Cancer Res Treat* 2005;90:187-9.

Håkansson S, Johannsson O, Johannsson U, et al. Moderate frequency of BRCA1 and BRCA2 germ-line mutations in Scandinavian familial breast cancer. *Am J Hum Genet* 1997;60:1068-78.

Hästbacka J, de la Chapelle A, Kaitila I, Sistonen P, Weaver A, Lander E. Linkage disequilibrium mapping in isolated founder populations: diastrophic dysplasia in Finland. *Nat Genet* 1992;2:204-11.

Höglund P, Sistonen P, Norio R, et al. Fine mapping of congenital chloride diarrhea gene by linkage disequilibrium. *Am J Hum Genet* 1995;57:95-102.

Iida A, Kurose K, Isobe R, et al. Mapping of a new target region of allelic loss to a 2-cM interval at 22q13.1 in primary breast cancer. *Genes Chromosomes Cancer* 1998;21:108-12.

International HapMap Consortium. The international HapMap project. *Nature* 2003;426:789-96.

International HapMap Consortium. A haplotype map of the human genome. *Nature* 2005;437:1299-320.

International Union Against Cancer. *TNM Classification of Malignant Tumours*. 6th ed. New York: Wiley-Liss, 2002;131-41.

Irminger-Finger I, Jefford CE. Is there more to BARD1 than BRCA1? *Nature Rev Cancer* 2006;6:38291.

Ishibe N, Hankinson SE, Colditz GA, et al. Cigarette smoking, cytochrome P450 1A1 polymorphisms, and breast cancer risk in the Nurses' Health Study. *Cancer Res* 1998;58:667-71.

Ishitobi M, Miyoshi Y, Hasegawa S, et al. Mutational analysis of BARD1 in familial breast cancer patients in Japan. *Cancer Lett* 2003;200:1-7.

Jacobs EJ, Spencer Feigelson H, Bain EB, et al. Polymorphisms in the vascular endothelial growth factor gene and breast cancer risk in the Cancer Prevention Study II cohort. *Breast Cancer Res* 2006;8:R22.

Jaiswal AK, McBride OW, Adesnik M, Nebert DW. Human dioxin-inducible cytosolic NAD(P)H:menadione oxidoreductase. *J Biol Chem* 1988;263:13572-8.

Jang J-H, Shin K-H, Park J-G. Mutations in fibroblast growth factor receptor 2 and fibroblast growth factor receptor 3 genes associated with human gastric and colorectal cancers. *Cancer Res* 2001;61:3541-3.

Jasin M. Homologous recombination of DNA damage and tumorigenesis: the BRCA connection. *Oncogene* 2002;21:8981-93.

Jazaeri AA, Yee CJ, Sotiriou C, Brantley KR, Boyd J, Liu ET. Gene expression profiles of BRCA1-linked, BRCA2-linked, and sporadic ovarian cancers. *J Natl Cancer Inst* 2002;94:990-1000.

Jefcoate CR, Liehr JG, Santen RJ, et al. Tissue-specific synthesis and oxidative metabolism of estrogens. *J Natl Cancer Inst Monogr* 2000;27:95-112.

Jeffreys AJ, Kauppi L, Neumann R. Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex. *Nat Genet* 2001;29:217-22.

Jekimovs CR, Chen X, Arnold J, et al. Low frequency of CHEK2 1100delC allele in Australian multiple-case breast cancer families: functional analysis in heterozygous individuals. *Br J Cancer* 2005;92:784-90.

Jenne DE, Reimann H, Nezu J, et al. Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat Genet* 1998;18:38-43.

Jensen RA, Thompson ME, Jetton TL, et al. BRCA1 is excreted and exhibits properties of a granin. *Nat Genet* 1996;12:303-8.

Jensen DE, Proctor M, Marquis ST, et al. BAP1: a novel ubiquitin hydrolase which binds to the BRCA1 RING finger and enhances BRCA1-mediated cell growth suppression. *Oncogene* 1998;16:1097-112.

Jernström H, Chu W, Vesprini D, et al. Genetic factors related to racial variation in plasma levels of insulin-like growth factor-1: Implications for premenopausal breast cancer risk. *Mol Genet Metab* 2001;72:144-54.

Jin Q, Hemminki K, Enquist K, et al. Vascular endothelial growth factor polymorphisms in relation to breast cancer development and prognosis. *Clin Cancer Res* 2005;11:3647-53.

Johannesdottir G, Gudmundsson J, Bergthorsson JT, et al. High prevalence of the 999del5 mutation in Icelandic breast and ovarian cancer patients. *Cancer Res* 1996;56:3663-5.

Johannsson O, Ostermeyer EA, Håkansson S, et al. Founding BRCA1 mutations in hereditary breast and ovarian cancer in Southern Sweden. *Am J Hum Genet* 1996;58:441-50.

Johannsson OT, Idvall I, Anderson C, et al. Tumour biological features of BRCA1-induced breast and ovarian cancer. *Eur J Cancer* 1997;33:362-71.

Johnson GCL, Esposito L, Barratt BJ, et al. Haplotype tagging for the identification of common disease genes. *Nat Genet* 2001;29:233-7.

Johnson KC. Accumulating evidence on passive and active smoking and breast cancer risk. *Int J Cancer* 2005;117:619-28.

- Johnson N, Fletcher O, Naceur-Lombardelli C, dos Santos Silva I, Ashworth A, Peto J. Interaction between CHEK2\*1100delC and other low-penetrance breast-cancer susceptibility genes: a familial study. *Lancet* 2005;366:1554-7.
- Jorde LB. Linkage disequilibrium as a gene-mapping tool. *Am J Hum Genet* 1995;56:11-4.
- Jorde LB. Linkage disequilibrium and the search for complex disease genes. *Genome Res* 2000;10:1435-44.
- Jorgenson E, Witte JS. A gene-centric approach to genome-wide association studies. *Nat Rev Genet* 2006;7:885-91.
- Joseph PD. The role of peroxidase-catalyzed activation of aromatic amines in breast cancer. *Mutagenesis* 1996;11:3-7.
- Junien C, Huerre C, Despoisse S, Gilgenkrantz S, Lenoir GM. c-Ha-ras1 is not deleted in del(11p13) Wilm's tumor (WAGR) and maps to 11p15.1-11p15.5. *Cytogenet Cell Genet* 1984;37:503.
- Justenhoven C, Hamann U, Pesch B, et al. ERCC2 genotypes and a corresponding haplotype are linked with breast cancer risk in a German population. *Cancer Epidemiol Biomarkers Prev* 2004;13:2059-64.
- Juul A, Bang P, Hertel NT, et al. Serum insulin-like growth factor-1 in 1030 healthy children, adolescents, and adults: relation to age, sex, stage of puberty, testicular size, and body mass index. *J Clin Endocrinol Metab* 1994;78:744-52.
- Juul A, Dalgaard P, Blum WF, et al. Serum levels of insulin-like growth factor (IGF)-binding protein-3 (IGFBP-3) in healthy infants, children, and adolescents: the relation to IGF-I, IGF-II, IGFBP-1, IGFBP-2, age, sex, body mass index, and pubertal maturation. *J Clin Endocrinol Metab* 1995;80:2534-42.
- Järvelä I. Infantile neuronal ceroid lipofuscinosis (CLN1): linkage disequilibrium in the Finnish population and evidence that variant late infantile form (variant CLN2) represents a nonallelic locus. *Genomics* 1991;10:333-7.
- Kadouri L, Kote-Jarai Z, Easton DF, et al. Polyglutamine repeat length in the AIB1 gene modifies breast cancer susceptibility in BRCA1 carriers. *Int J Cancer* 2004;108:399-403.
- Kamb A, et al. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 1994;264:436-40.
- Karhu R, Laurila E, Kallioniemi A, Syrjäkoski K. Large genomic BRCA2 rearrangements and male breast cancer. *Cancer Detect Prev* 2006;30:530-4.
- Karjalainen JM, Kellokoski JK, Mannermaa AJ, et al. Failure in post-transcriptional processing is a possible inactivation mechanism of AP-2alpha in cutaneous melanoma. *Br J Cancer* 2000;82:2015-21.

Karppinen SM, Heikkinen K, Rapakko K, Winqvist R. Mutation screening of the BARD1 gene: evidence for involvement of the Cys557Ser allele in hereditary susceptibility to breast cancer. *J Med Genet* 2004;41:e114.

Karppinen SM, Barkardottir RB, Backenhorn K, et al. Nordic collaborative study of the BARD1 Cys557Ser allele in 3956 patients with cancer: enrichment in familial BRCA1/BRCA2 mutation-negative breast cancer but not in other malignancies. *J Med Genet* 2006;43:856-62.

Kastan MB, Lim DS. The many substrates and functions of ATM. *Nat Rev Mol Cell Biol* 2000;1:179-86.

Kataoka N, Cai Q, Wen W, et al. Population-based case-control study of VEGF gene polymorphisms and breast cancer risk among Chinese women. *Cancer Epidemiol Biomarkers Prev* 2006;15:1148-52.

Keen J, Lester D, Ingelhearn C, Curtis A, Bhattacharya S. Rapid detection of single base mismatches as heteroduplexes on hydrolink gels. *Trends Genet* 1991;7:5.

Kelsey JL, Gammon MD, John EM. Reproductive factors and breast cancer. *Epidemiol Rev* 1993;15:36-47.

Kere J, Sistonen P, Holmberg C, de la Chapelle A. The gene for congenital chloride diarrhea maps close to but is distinct from the gene for cystic fibrosis transmembrane conductance regulator. *Proc Natl Acad Sci USA* 1993;90:10686-9.

Ketterer B. Protective role of glutathione and glutathione transferases in mutagenesis and carcinogenesis. *Mutat Res* 1988; 202:343-61.

Khanna KK, Keating KE, Kozlov S, et al. ATM associates with and phosphorylates p53: mapping the region of interaction. *Nat Genet* 1998;20:398-400.

Khanna KK, Jackson SP. DNA double-strand breaks: signaling, repair and the cancer connection. *Nature Genet* 2001;27:247-54.

Kilpivaara O, Bartkova J, Eerola H, et al. Correlation of CHEK2 protein expression and c.1100delC mutation status with tumor characteristics among unselected breast cancer patients. *Int J Cancer* 2005;113:575-80.

Kilpivaara O, Vahteristo P, Falck J, et al. CHEK2 variant I157T may be associated with increased breast cancer risk. *Int J Cancer* 2004;111:543-7.

Kim SU, Park SK, Yoo KS, et al. XRCC1 genetic polymorphism and breast cancer risk. *Pharmacogenetics* 2002;12:335-8.

King MC, Marks JH, Mandell JB for The New York Breast Cancer Study Group. Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* 2003;302:643-6.

Kinzler KW, Vogelstein B. Gatekeepers and caretakers. *Nature* 1997;384:761-2.

Kischkel FC, Kioschis P, Weitz S, Poustka A, Lichter P, Krammer PH. Assignment of CASP8 to human chromosome band 2q33-->q34 and Casp8 to the murine syntenic region on chromosome 1B-proximal c by in situ hybridization. *Cytogenet Cell Genet* 1998;82:95-6.

Klebanoff SJ. Oxygen metabolism and the toxic properties of phagocytes. *Ann Intern Med* 1980;93:480-9.

Kleibl Z, Novotny J, Bezdiekova D, et al. The CHEK2 c.1100delC germline mutation rarely contributes to breast cancer development in the Czech Republic. *Breast Cancer Res Treat* 2005;90:165-7.

Knudson AG Jr. Mutation and cancer: Statistical study of retinoblastoma. *Proc Nat Acad Sci USA* 1971;68:820-3.

Koonin EV, Altschul SF. ...Functional motifs... *Nat Genet* 1996;13:266-7.

Krippel P, Langsenlehner U, Renner W, et al. A common 936C/T gene polymorphism of vascular endothelial growth factor is associated with decreased breast cancer risk. *Int J Cancer* 2003;106:468-71.

Kristensen VN, Andersen TI, Lindblom A, Erikstein B, Magnus P, Borresen-Dale AL. A rare CYP19 (aromatase) variant may increase the risk of breast cancer. *Pharmacogenetics* 1998;8:43-8.

Kristensen VN, Borresen-Dale AL. Molecular epidemiology of breast cancer: genetic variation in steroid hormone metabolism. *Mut Res* 2000;462:323-33.

Krontiris TG, Devlin B, Karp DD, Robert NJ, Risch N. An association between the risk of cancer and mutations in the HRAS1 minisatellite locus. *N Engl J Med* 1993;329:517-23.

Kruglyak L. Prospects for whole-genome linkage disequilibrium mapping of common disease genes. *Nat Genet* 1999;22:139-44.

Kumar R, Höglund L, Zhao C, Försti A, Snellman E, Hemminki K. Single nucleotide polymorphisms in the XPG gene: determination of role in DNA repair and breast cancer risk. *Int J Cancer* 2003;103:671-5.

Kuschel B, Auranen A, McBride S, et al. Variants in DNA double-strand break repair genes and breast cancer susceptibility. *Hum Mol Genet* 2002;11:1399-407.

Kuschel B, Auranen A, Gregory CS, et al. Common polymorphisms in Checkpoint Kinase 2 are not associated with breast cancer. *Cancer Epid Biom Prev* 2003;12:809-12.

Kuschel B, Chenevix-Trench G, Spurdle AB, et al. Common polymorphisms in ERCC2 (Xeroderma pigmentosum D) are not associated with breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2005;14:1828-31.

Laan M, Pääbo S. demographic history and linkage disequilibrium in human populations. *Nat Genet* 1997;17:435-8.



Lahti-Domenici J, Rapakko K, Pääkkönen K, et al. Exclusion of large deletions and other rearrangements in BRCA1 and BRCA2 in Finnish breast and ovarian cancer families. *Cancer Genet Cytogenet* 2001;129:120-3.

Laitinen T, Polvi A, Rydman P, et al. Characterization of a common susceptibility locus for asthma-related traits. *Science* 2004;304:300-4.

Lakhani SR, Jacquemier J, Sloane JP, et al. Multifactorial analysis of differences between sporadic breast cancers and cancers involving BRCA1 and BRCA2 mutations. *J Natl Cancer Inst* 1998;90:1138-45.

Lakhani SR, Gusterson BA, Jacquemier J, et al. The pathology of familial breast cancer: Histological features of cancers in families not attributable to mutations in BRCA1 or BRCA2. *Clin Cancer Res* 2000;6:782-9.

Lakhani SR, Manek S, Penault-Llorca F, et al. Pathology of ovarian cancers in BRCA1 and BRCA2 carriers. *Clin Cancer Res* 2004;10:2473-81.

Laloo F, Varley J, Ellis D, et al. Prediction of pathogenic mutations in patients with early-onset breast cancer by family history. *Lancet* 2003;361:1101-2.

Laloo F, Varley J, Moran A, et al. BRCA1, BRCA2 and TP53 mutations in very early-onset breast cancer with associated risks to relatives. *Eur J Cancer* 2006;42:1143-50.

Lancaster JM, Wooster R, Mangion J, et al. BRCA2 mutations in primary breast and ovarian cancers. *Nat Genet* 1996;13:238-40.

Lane D. Cancer. p53, guardian of the genome. *Nature* 1992;358:15-6.

Lauge A, Lefebvre C, Laurent-Puig P, et al. No evidence for germline PTEN mutations in families with breast and brain tumours. *Int J Cancer* 1999;84:216-9.

Laurila E, Syrjäkoski K, Holli K, Kallioniemi A, Karhu R. Search for large genomic alterations of the BRCA1 gene in Finnish population. *Cancer Genet Cytogenet* 2005;163:57-61.

Law ML, Kao FT, Wei Q, et al. The progesterone receptor gene maps to human chromosome band 11q13, the site of the mammary oncogene int-2. *Proc Natl Acad Sci USA* 1987;84:2877-81.

Layde PM, Webster LA, Baughman AL, Wingo PA, Rubin GL, Ory HW. The independent associations of parity, age at first full term pregnancy, and duration of breastfeeding with the risk of breast cancer. Cancer and Steroid Hormone Study Group. *J Clin Epidemiol* 1989;42:963-73.

Le Marchand L. The predominance of the environment over genes in cancer causation: implications for genetic epidemiology. *Cancer Epidemiol Biomarkers Prev* 2005;14:1037-9.

Le Marchand L, Donlon T, Kolonel LN, Henderson BE, Wilkens LR. Estrogen metabolism-related genes and breast cancer risk: The multiethnic cohort study. *Cancer Epidemiol Biomarkers Prev* 2005;14:1998-2003.

Lee JS, Collins KM, Brown AL, Lee CH, Chung JH. hCds1-mediated phosphorylation of BRCA1 regulates the DNA damage response. *Nature* 2000;404:201-4.

Lee KM, Park SK, Kim SU, et al. N-acetyltransferase (NAT1, NAT2) and glutathione S-transferase (GSTM1, GSTT1) polymorphisms in breast cancer. *Cancer Lett* 2003;196:179-86.

Lee KM, Choi JY, Park SK, et al. Genetic polymorphisms of ataxia telangiectasia mutated and breast cancer. *Cancer Epidemiol Biomarkers Prev* 2005a;14:821-5.

Lee KM, Choi LY, Kang C, et al. Genetic polymorphisms of selected DNA repair genes, estrogen and progesterone receptor status, and breast cancer risk. *Clin Cancer Res* 2005b;11:4620-6.

Lee KM, Park SK, Hamajima N, et al. Genetic polymorphisms of TGF- $\beta$ 1 & TNF- $\beta$  and breast cancer risk. *Breast Cancer Res Treat* 2005c;90:149-55.

Levrn O, Attwooll C, Henry RT, et al. The BRCA1-interacting helicase BRIP1 is deficient in Fanconi anemia. *Nat Genet* 2005;37:931-3.

Levy-Lahad E, Lahad A, Eisenberg S, et al. A single nucleotide polymorphism in the RAD51 gene modifies cancer risk in BRCA2 but not BRCA1 carriers. *Proc Natl Acad Sci USA* 2001;98:3232-6.

Li FP, Fraumeni JF. Rhabdomyosarcoma in children: an epidemiologic study and identification of a familial cancer syndrome. *J Nat Cancer Inst* 1969;43:1364-73.

Li FP, Fraumeni JF Jr, Mulvihill JJ, et al. A cancer family syndrome in twenty-four kindreds. *Cancer Res* 1988;48:5358-62.

Li J, Stern DF. Regulation of CHEK2 by DNA-dependent protein kinase. *J Biol Chem* 2005;280:12041-50.

Liaw D, Marsh DJ, Li J, et al. Germline mutations of the PTEN gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nat Genet* 1997;16:64-7.

Lichtenstein P, Holm NV, Verkasalo PK, et al. Environmental and heritable factors in the causation of cancer--analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 2000;343:78-85.

Liderau R, Soussi T. absence of p53 germ-line mutations in bilateral breast cancer patients. *Hum Genet* 1992;89:250-2.

Liede A, Rehal P, Vesprini D, Jack E, Abrahamson J, Narod SA. A breast cancer patient of Scottish descent with germ-line mutations in BRCA1 and BRCA2. *Am J Hum Genet* 1998;62:1543-4.

Liehr JG. Hormone-associated cancer: mechanistic similarities between human breast cancer and estrogen-induced kidney carcinogenesis in hamsters. *Environ Health Perspect* 1997;105:565-9.

Lillie EO, Bernstein L, Ursin G. The role of androgens and polymorphisms in the androgen receptor in the epidemiology of breast cancer. *Breast Cancer Res* 2003;5:164-73.

Lin HJ, Han CY, Lin BK, Hardy S. Slow acetylator mutations in the human polymorphic N-acetyltransferase gene in 786 Asians, blacks, Hispanics, and whites: application to metabolic epidemiology. *Am J Hum Genet* 1993;52:827-34.

Lin SC, Chou YC, Wu MH, et al. Genetic variants of myeloperoxidase and catechol-O-methyltransferase and breast cancer risk. *Eur J Cancer Prev* 2005;14:257-61.

Lind C, Hochstein P, Ernster L. DT-diaphorase as a quinone reductase: a cellular control device against semiquinone and superoxide radical formation. *Arch Biochem Biophys* 1982;216:178-85.

Lippert TH, Seeger H, Mueck AO. The impact of endogenous estradiol metabolites on carcinogenesis. *Steroids* 2000;65:357-69.

Loman N, Johannsson O, Bendahl PO, Borg Å, Fernö M, Olsson H. Steroid receptors in hereditary breast carcinomas associated with BRCA1 or BRCA2 mutations or unknown susceptibility genes. *Cancer* 1998;83:310-9.

Loman N, Johannsson O, Bendahl PO, et al. Prognosis and clinical presentation of BRCA2-associated breast cancer. *Eur J Cancer* 2000;36:1365-73.

Long JR, Egan KM, Dunning L, et al. Population-based case-control study of AhR (aryl hydrocarbon receptor) and CYP1A2 polymorphisms and breast cancer. *Pharmacogenet Genomics* 2006;16:237-43.

Longy M, Lacombe D. Cowden disease. Report of a famili and review. *Ann Genet* 1996;39:35-42.

Low YL, Dunning AM, Dowsett M, et al. Implications of gene-environment interaction in studies of gene variants in breast cancer: An example of dietary isoflavones and the D356N polymorphism in the sex hormone-binding globulin gene. *Cancer Res* 2006;66:8980-3.

Lu J, Wei Q, Bondy ML, et al. Polymorphisms and haplotypes of the NBS1 gene are associated with risk of sporadic breast cancer in non-Hispanic white women  $\geq 55$  years. *Carcinogenesis* 2006;27:2209-16.

Lubahn DB, Joseph DR, Sar M, et al. The human androgen receptor: complementary deoxyribonucleic acid cloning, sequence analysis and gene expression in prostate. *Mol Endocrinol* 1988;2:1265-75.

MacMahon B, Trichopoulos D, Brown J, et al. Age at menarche, urine estrogens and breast cancer risk. *Int J Cancer* 1982;30:427-31.

MacPherson G, Healey CS, Teare MD, et al. Association of a common variant of the CASp8 gene with reduced risk of breast cancer. *J Natl Cancer Inst* 2004;96:1866-9.

Malkin D, Li FP, Strong LC, et al. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* 1990;250:1233-8.

Mannermaa A, Peltoketo H, Winqvist R, et al. Human familial and sporadic breast cancer: analysis of the coding regions of the 17 beta-hydroxysteroid dehydrogenase 2 gene (EDH17B2) using a single-strand conformation polymorphism assay. *Hum Genet* 1994;93:319-24.

Mannervik B. The isoenzymes of glutathione transferase. *Adv Enzymol Relat Areas Mol Bio* 1985;57:357-417.

Marcus JN, Watson P, Page DL, et al. Hereditary breast cancer: pathobiology, prognosis, and BRCA1 and BRCA2 gene linkage. *Cancer* 1996;77:697-709.

Marsh DJ, Coulon V, Lunetta KL, et al. Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation. *Hum Mol Genet* 1998;7:507-15.

Martucci CP, Fishman J. P450 enzymes of estrogen metabolism. *Pharmacol Ther* 1993;57:237-57.

Mateus Pereira LH, Sigurdson AJ, Doody MM, et al. CHEK2:1100delC and female breast cancer in the United States. *Int J Cancer* 2004;112:541-3.

Matsuoka S, Huang M, Elledge SJ. Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 1998;282:1893-7.

Maxam AM, Gilbert W. A new method for sequencing DNA. *Proc Natl Acad Sci* 1977;74:560-4.

McAllister KA, Haugen-Strano A, Hagevik S, et al. Characterization of the rat and mouse homologues of the BRCA2 breast cancer susceptibility gene. *Cancer Res* 1997;57:3121-5.

McBride OW, Merry D, Givol D. The gene for human p53 cellular tumor antigen is located on chromosome 17 short arm (17p13). *Proc Natl Acad Sci USA* 1986;83:130-4.

McQueen CA, Maslansky CJ, Glowinski IB, Crescenzi SB, Weber WW, Williams GM. Relationship between the genetically determined acetylator phenotype and DNA damage induced by hydralazine and 2-aminofluorene in cultured rabbit hepatocytes. *Proc Natl Acad Sci USA* 1982;79:1269-72.

Meijers-Heijboer H, Wijnen J, Vasen H, et al. The CHEK2 1100delC mutation identifies families with a hereditary breast and ovarian cancer phenotype. *Am J Hum Genet* 2003;72:1308-14.

Menard S, Casalini P, Campiglio M, Pupa SM, Tagliabue E. Role of HER2/neu in tumor progression and therapy. *Cell Mol Life Sci* 2004;61:2965-78.

Menasce LP, White GR, Harrison CJ, Boyle JM. Localization of the estrogen receptor locus (ESR) to chromosome 6q25.1 by FISH and simple post-FISH banding technique. *Genomics* 1993;17:263-5.

- Menzel HJ, Sarmanova J, Soucek P, et al. Association of NQO1 polymorphism with spontaneous breast cancer in two independent populations. *Br J Cancer* 2004;90:1989-94.
- Meraldi P, Honda R, Nigg EA. Aurora kinases link chromosome segregation and cell division to cancer susceptibility. *Curr Opin Genet Dev* 2004;14:29-36.
- Meravjer SD, Pham TM, Caduff RF, et al. Somatic mutations in the BRCA1 gene in sporadic ovarian tumours. *Nat Genet* 1995;9:439-43.
- Metsola K, Kataja V, Sillanpää P, et al. XRCC1 and XPD genetic polymorphisms, smoking and breast cancer risk in a Finnish case-control study. *Breast Cancer Res* 2005;7:R987-R997.
- Meza JE, Brzovic PS, King MC, Klevit RE. Mapping the functional domains of BRCA1. *J Biol Chem* 1999;274:5659-65.
- Miki Y, Swensen J, Shattuck-Eidens D, et al. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* 1994;266:66-71.
- Miki Y, Katagiri T, Kasumi F, Yoshimoto T, Nakamura Y. Mutation analysis in the BRCA2 gene in primary breast cancers. *Nat Genet* 1996;13:245-7.
- Millikan RC, Pittman GS, Newman B, et al. Cigarette smoking, N-acetyltransferases 1 and 2, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 1998;7:371-8.
- Millikan R, Eaton A, Worley K, et al. HER2 codon 655 polymorphism and risk of breast cancer in African Americans and whites. *Breast Cancer Res Treat* 2003;79:355-64.
- Milne RL, Ribas G, Gonzales-Neira A, et al. ERCC4 associated with breast cancer risk: A two-stage case-control study using high-throughput genotyping. *Cancer Res* 2006;66:9420-7.
- Milner J, Ponder B, Hughes-Davies L, Seltsmann M, Kouzarides T. Transcriptional activation functions in BRCA2. *Nature* 1997;386:772-3.
- Misrahi M, Atger M, d'Auriol L, et al. Complete amino acid sequence of the human progesterone receptor from cloned cDNA. *Biochem Biophys Res Commun* 1987;143:740-8.
- Mitrunen K, Jourenkova N, Kataja V, et al. Steroid metabolism gene CYP17 polymorphism and the development of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2000;9:1343-8.
- Mitrunen K, Jourenkova N, Kataja V, et al. Glutathione S-transferase M1, M3, P1 and T1 genetic polymorphisms and susceptibility to breast cancer. *Cancer Epidemiol Biomarkers Prev* 2001a;10:229-36.
- Mitrunen K, Sillanpää P, Kataja V, et al. Association between manganese superoxide dismutase (MnSOD) gene polymorphism and breast cancer risk. *Carcinogenesis* 2001b;22:827-9.
- Mitrunen K, Jourenkova N, Kataja V, et al. Polymorphic catechol-O-methyltransferase gene and breast cancer risk. *Cancer epidemiol Biomarkers Prev* 2001c;10:635-40.

- Moffa AB, Tannheimer SL, Ethier SP. Transforming potential of alternatively spliced variants of fibroblast growth factor receptor 2 in human mammary epithelial cells. *Mol Cancer Res* 2004;2:643-52.
- Mohlke KL, Lange EM, Valle TT, et al. Linkage disequilibrium between microsatellite markers extends beyond 1 cM on chromosome 20 in Finns. *Genome Res* 2001;11:1221-6.
- Moisio AL, Sistonen P, Weissenbach J, de la Chapelle A, Peltomäki P. Age and origin of two common MLH1 mutations predisposing to hereditary colon cancer. *Am J Hum Genet* 1996;59:1243-51.
- Monzon J, et al. CDKN2A mutations in multiple primary melanomas. *N Engl J Med* 1998;338:879-87.
- Moslehi R, Chu W, Karlan B, et al. BRCA1 and BRCA2 mutation analysis of 208 Ashkenazi Jewish women with ovarian cancer. *Am J Hum Genet* 2000;66:1259-72.
- Moynahan ME, Chiu JW, Koller BH, Jasin M. Brca1 controls homology-directed DNA repair. *Mol Cell* 1999;4:511-8.
- Moynahan ME, Pierce AJ, Jasin M. BRCA2 is required for homology-directed repair of chromosomal breaks. *Mol Cell* 2001;7:263-72.
- Mueller CR, Roskelley CD. Regulation of BRCA1 expression and its relationship to sporadic breast cancer. *Breast Cancer Res* 2003;5:45-52.
- Männistö S, Pietinen P, Pyy M, Palmgren J, Eskelinen M, Uusitupa M. Body-size indicators and risk of breast cancer according to menopause and estrogen-receptor status. *Int J Cancer* 1996;68:8-13.
- Männistö S. Diet, body size, and risk of breast cancer. A case-control study. Publications of the National Public Health Institute A 17/1999. Helsinki, National Public Health Institute, 1999.
- Nair J, Barbin A, Velic I, Bartsch H. Etheno DNA-base adducts from endogenous reactive species. *Mutation Research* 1999;424:59-69.
- Narod SA, Boyd J. Current understanding of the epidemiology and clinical implications of BRCA1 and BRCA2 mutations for ovarian cancer. *Curr Opin Obstet Gynecol* 2002;14:19-26.
- Nelen MR, Padberg GW, Peeters EA, et al. Localization of the gene for Cowden disease to chromosome 10q22-23. *Nat Genet* 1996;13:114-6.
- Nelson SE, Gould MN, Hampton JM, Trentham-Dietz A. A case-control study of the HER2 Ile655Val polymorphism in relation to risk of invasive breast cancer. *Breast Cancer Res* 2005;7:R357-R364.
- Netzel-Arnett S, Hooper JD, Szabo R, et al. Membrane anchored serine proteases: A rapidly expanding group of cell surface proteolytic enzymes with potential roles in cancer. *Cancer Metastasis Rev* 2003;22:237-58.

Neuhausen SL, Mazoyer S, Friedman L, et al. Haplotype and phenotype analysis of six recurrent BRCA1 mutations in 61 families: results of an international study. *Am J Hum Genet* 1996;58:271-80.

Neuhausen S, Dunning A, Steele L, et al. Role of CHEK2\*1100delC in unselected series of non-BRCA1/2 male breast cancers. *Int J Cancer* 2004;108:477-8.

Nevanlinna HR. The Finnish population structure. A genetic and genealogical study. *Hereditas* 1972;71:195-236.

Nordling M, Karlsson P, Wahlstrom J, Engwall Y, Wallgren A, Martinsson T. A large deletion disrupts the exon 3 transcription activation domain of the BRCA2 gene in a breast/ovarian cancer family. *Cancer Res* 1998;58:1372-5.

Norio R, Nevanlinna HR, Perheentupa J. Hereditary diseases in Finland; rare flora in rare soul. *Ann Clin Res* 1973;5:109-41.

Norio R. Finnish Disease Heritage II: population prehistory and genetic roots of Finns. *Hum Genet* 2003a;112:457-69.

Norio R. Finnish Disease Heritage I: characteristics, causes, background. *Hum Genet* 2003b;112:441-56.

Norio R. Finnish Disease Heritage III: the individual diseases. *Hum Genet* 2003c;112:470-526.

Nyström-Lahti M, Sistonen P, Mecklin JP, et al. Close linkage to chromosome 3p and conservation of ancestral founding haplotype in hereditary nonpolyposis colorectal cancer families. *Proc Natl Acad Sci USA* 1994;91:6054-8.

Offit K, Pierce H, Kirchhoff T, et al. Frequency of CHEK2\*1100delC in New York breast cancer cases and controls. *BMC Med Genet* 2003;4:1.

Ohayon T, Gal I, Baruch RG, Szabo C, Friedman E. CHEK2\*1100delC and male breast cancer risk in Israel. *Int J Cancer* 2004;108:479-80.

Ohayon T, Gershoni-Baruch R, Papa MZ, Menachem TD, Barzilai SE, Friedman E. The R72P P53 mutation is associated with familial breast cancer in Jewish women. *Br J Cancer* 2005;92:1144-8.

Oldenburg RA, Kroeze-Jansema K, Kraan J, et al. The CHEK2\*1100delC variant acts as a breast cancer risk modifier in non-BRCA1/BRCA2 multiple-case families. *Cancer Res* 2003;63:8153-7.

Olsen JH, Hahnemann JM, Borresen-Dale AL, et al. Cancer in patients with ataxia-telangiectasia and in their relatives in the Nordic Countries. *J Natl Cancer Inst* 2001;93:121-7.

Omura T, Morohashi K. Gene regulation of steroidogenesis. *J Steroid Biochem Mol Biol* 1995;53:19-25.

Ophoff RA, Escamilla MA, Service SK, et al. Genomewide linkage disequilibrium mapping of severe bipolar disorder in a population isolate. *Am J Hum Genet* 2002;71:565-74.

Orita M, Iwahana H, Kanazawa H, Hayashi K, Sekiya T. Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proc Natl Acad Sci USA* 1989;86:2766-70.

Osorio A, Rodriguez-Lopez R, Diez O, et al. The breast cancer low-penetrance allele 1100delC in the CHEK2 gene is not present in Spanish familial breast cancer population. *Int J Cancer* 2004;108:54-6.

Overall CM, Tam EM, Kappelhoff R, et al. Protease degradomics: mass spectrometry discovery of protease substrates and the CLIP-CHIP, a dedicated DNA microarray of all human proteases and inhibitors. *Biol Chem* 2004;385:493-504.

Papewalis J, Nikitin AY, Rajewsky MF. G to A polymorphism at amino acid codon 655 of the human erbB-2/HER2 gene. *Nucleic Acids Res* 1991;19:5452.

Parl FF. Glutathione S-transferase genotypes and cancer risk. *Cancer Letters* 2005;221:123-9.

Patel KJ, Yu VPCC, Lee H, et al. Involvement of Brca2 in DNA repair. *Mol Cell* 1998;1:347-57.

Patel JH, Loboda AP, Showe MK, Showe LC, McMahon SB. Analysis of genomic targets reveals complex functions of MYC. *Nat Rev Cancer* 2004;4:562-8.

Patel KJ. Fanconi anemia and breast cancer susceptibility. *Nat Genet* 2007;39:142-3.

Patil N, Berno AJ, Hinds DA, et al. Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. *Science* 2001;294:1719-23.

Payne AH. Hormonal regulation of cytochrome P450 enzymes, cholesterol side-chain cleavage and 17 $\alpha$ -hydroxylase/C17-20 lyase in Leydig cells. *Biol Reprod* 1990;42:399-404.

Pellegrini L, Yu DS, Lo T, et al. Insights into DNA recombination from the structure of a RAD51-BRCA2 complex. *Nature* 2002;420:287-93.

Pellegrini L, Venkitaraman A. Emerging functions of BRCA2 in DNA recombination. *Trends Biochem Sci* 2004;29:310-16.

Peltonen L, Jalanko A, Varilo T. Molecular genetics of the Finnish disease heritage. *Hum Mol Genet* 1999;8:1913-23.

Peltonen L. Positional cloning of disease genes: advantages of genetic isolates. *Hum Hered* 2000;50:66-75.

Peltonen L, Palotie A, Lange K. Use of population isolates for mapping complex traits. *Nat Rev Genet* 2000;1:182-90.



Perou CM, Sorlie T, Eisen MB, et al. Molecular portraits of human breast tumours. *Nature* 2000;406:747-52.

Peto J, Collins N, Barfoot R, et al. Prevalence of BRCA1 and BRCA2 gene mutations in patients with early-onset breast cancer. *J Natl Cancer Inst* 1999;91:943-9.

Petrij-Bosch A, Peelen T, van Vliet M, et al. BRCA1 genomic deletions are major founder mutations in Dutch breast cancer patients. *Nat Genet* 1997;17:341-5.

Petruska JM, Mosebrook DR, Jakab GJ, Trush MA. Myeloperoxidase-enhanced formation of (+-)-trans-7,8-dihydroxy-7,8-dihydrobenzo[a]pyrene-DNA adducts in lung tissue in vitro: a role of pulmonary inflammation in the bioactivation of a procarcinogen. *Carcinogenesis* 1992;13:1075-81.

Pharoah PDP, Day NE, Duffy S, Easton DF, Ponder BAJ. Family history and the risk of breast cancer: a systematic review and meta-analysis. *Int J Cancer* 1997;71:800-9.

Pharoah PDP, Easton DF, Stockton DL, Gayther S, Ponder BAJ, United Kingdom Coordinating Committee for Cancer Research (UKCCCR) Familial Ovarian Cancer Study Group: Survival in familial, BRCA1-associated, and BRCA2-associated epithelial ovarian cancer. *Cancer Res* 1999;59:868-71.

Pharoah PDP, Dunning AM, Ponder BAJ, Easton DF. Association studies for finding cancer-susceptibility genetic variants. *Nat Rev Cancer* 2004;4:850-60.

Phelan CM, Rebbeck TR, Weber BL, et al. Ovarian cancer risk in BRCA1 carriers is modified by the HRAS1 variable number of tandem repeat (VNTR) locus. *Nat Genet* 1996;12:309-11.

Phillips MS, Lawrence R, Sachidanandam R, et al. Chromosomewide distribution of haplotype blocks and the role of recombination hot spots. *Nat Genet* 2003;33:382-7.

Pike MC, Gerkins VR, Casagrande JT, Gray GE, Brown J and Henderson BE. The hormonal basis of breast cancer. *Natl Cancer Inst Monogr* 1979;53:187-93.

Pippard EC, Hall AJ, Barker DJP, Bridges BA. Cancer in homozygotes and heterozygotes of ataxia-telangiectasia and xeroderma pigmentosum in Britain. *Cancer Res* 1988;48:2929-32.

Pirinen K. Rajamaakunta asutusliikkeen aikakautena. In: *Savon historia II* pp 1534-617. Kustannuskiila OY, Pieksämäki, 1982.

Plummer SJ, Anton-Culver H, Webster L, et al. Detection of BRCA1 mutations by the protein truncation test. *Hum Mol Genet* 1995;10:1989-91.

Pooley KA, Healey CS, Smith PL, et al. Association of the progesterone receptor gene with breast cancer risk: a single-nucleotide polymorphism tagging approach. *Cancer Epidemiol Biomarkers Prev* 2006;15:675-82.

Powers JT, Hong SK, Mayhew CN, Rogers PM, Knudsen ES, Johnson DG. E2F1 uses the ATM signalling pathway to induce p53 and Chk2 phosphorylation and apoptosis. *Mol Cancer Res* 2004;2:203-14.

- Povey S, Wilson D, Edwards YH. Assignment of a human diaphorase (DIA4) to chromosome 16. *Ann Hum Genet* 1980;43:349-53.
- Preston DL, Mattsson A, Holmberg E, Shore R, Hildreth NG, Boice JD Jr. Radiation effects on breast cancer risk: a pooled analysis of eight cohorts. *Radiat Res* 2002;158:220-35.
- Pritchard JK, Przeworski M. Linkage disequilibrium in humans: Models and data. *Am J Hum Genet* 2001;69:1-14.
- Puget N, Torchard D, Serova-Sinilnikova OM, et al. A 1-kb Alu-mediated germ-line deletion removing BRCA1 exon 17. *Cancer Res* 1997;57:828-31.
- Puget N, Sinilnikova OM, Stoppa-Lyonnet D, et al. An Alu-mediated 6-kb duplication in the BRCA1 gene: A new founder mutation? *Am J Hum Genet* 1999;64:300-2.
- Pukkala E, Weiderpass E. time trends in socio-economic differences in incidence rates of cancers of the breast and female genital organs (Finland, 1971-1995). *Int J Cancer* 1999;81:56-61.
- Pukkala E, Kesminiene A, Poliakov S, et al. Breast cancer in Belarus and Ukraine after the Chernobyl accident. *Int J Cancer* 2006;119:651-8.
- Purcell S, Cherny SS, Sham PC. Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics* 2003;19:149-50.
- Pylkäs K, Tommiska J, Syrjäkoski K, et al. Evaluation of the role of Finnish ataxia-telangiectasia mutations in hereditary predisposition to breast cancer. *Carcinogenesis* 2007;28:1040-5.
- Pääkkönen K, Sauramo S, Sarantaus L, et al. Involvement of BRCA1 and BRCA2 in breast cancer in a Western Finnish sub-population. *Genet Epidemiol* 2001;20:239-46.
- Rafii S, O'Reagan P, Xinarianos G, et al. A potential role for the XRCC2 R188H polymorphic site in DNA-damage repair and breast cancer. *Hum Mol Genet* 2002;11:1433-8.
- Rahman N, Seal S, Thompson D, et al. PALB2, which encodes a BRCA2-interacting protein, is a breast cancer susceptibility gene. *Nat Genet* 2007;39:165-7.
- Ramus SJ, Kote-Jarai Z, Friedman LS, et al. Analysis of BRCA1 and BRCA2 mutations in Hungarian families with breast or breast-ovarian cancer. *Am J Hum Genet* 1997a;60:1242-6.
- Ramus SJ, Friedman LS, Gayther SA, et al. A breast/ovarian cancer patient with germline mutations in both BRCA1 and BRCA2. *Nat Genet* 1997b;15:14-5.
- Rapakko K, Allinen M, Syrjäkoski K, et al. Germline TP53 alterations in Finnish breast cancer families are rare and occur at conserved mutation-prone sites. *Br J cancer* 2001;84:116-9.
- Rashid MU, Jakubowska A, Justenhoven C, et al. German populations with infrequent CHEK2\*1100delC and minor associations with early-onset and familial breast cancer. *Eur J Cancer* 2005;41:2896-2903.

- Rebbeck TR. Inherited genetic predisposition in breast cancer. A population-based perspective. *Cancer* 1999;86:2493-2501.
- Rebbeck TR, Wang Y, Kantoff PW, et al. Modification of BRCA1- and BRCA2-associated breast cancer risk by AIB1 genotype and reproductive history. *Cancer Res* 2001;61:5420-4.
- Reich DE, Cargill M, Bolk S, et al. Linkage disequilibrium in the human genome. *Nature* 2001;411:199-204.
- Reid S, Schindler D, Hanenberg H, et al. Biallelic mutations in PALB2 cause Fanconi anemia subtype FA-N and predispose to childhood cancer. *Nat Genet* 2007;39:162-4.
- Renehan AG, Zwahlen M, Minder C, O'Dwyer ST, Shalet SM, Egger M. Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis. *Lancet* 2004;363:1346-53.
- Renwick A, Thompson D, Seal S, et al. ATM mutations that cause ataxia-telangiectasia are breast cancer susceptibility alleles. *Nat Genet* 2006;38:873-5.
- Rice JC, Massey-Brown KS, Futscher BW. Aberrant methylation of the BRCA1 CpG island promoter is associated with decreased BRCA1 mRNA in sporadic breast cancer cells. *Oncogene* 1998;17:1807-12.
- Risch NJ. Searching for genetic determinants in the new millennium. *Nature* 2000;405:847-56.
- Risch HA, McLaughlin JR, Cole DEC, et al. Prevalence and penetrance of germline BRCA1 and BRCA2 mutations in a population series of 649 women with ovarian cancer. *Am J Hum Genet* 2001;68:700-10.
- Robson ME, Chappuis PO, Satagopan J, et al. A combined analysis of outcome following breast cancer: differences in survival based on BRCA1/BRCA2 mutation status and administration of adjuvant treatment. *Breast Cancer Research* 2003;6:R8-R17.
- Rodriguez-Lopez R, Osorio A, Ribas G, et al. The variant E233G of the RAD51D gene could be a low-penetrance allele in high-risk breast cancer families without BRCA1/2 mutations. *Int J Cancer* 2004;110:845-9.
- Roest PA, Roberts RG, Sugino S, van Ommen GJ, den Dunnen JT. Protein truncation test (PTT) for rapid detection of translation-terminating mutations. *Hum Mol Genet* 1993;2:1719-21.
- Rogoff HA, Pickering MT, Frame FM, et al. Apoptosis associated with deregulated E2F activity is dependent on E2F1 and Atm/Nbs1/Chk2. *Mol Cell Biol* 2004;24:2968-77.
- Ross JS, Fletcher JA, Linette GP, et al. The Her-2/neu gene and protein in breast cancer 2003: biomarker and target of therapy. *Oncologist* 2003;8:307-25.
- Roth S, Kristo P, Auranen A, et al. A missense mutation in the BRCA2 gene in three siblings with ovarian cancer. *Br J Cancer* 1998;77:1199-202.

Rubin SC, Benjamin I, Behbakt K, et al. Clinical and pathological features of ovarian cancer in women with germ-line mutations of BRCA1. *New Engl J Med* 1996;335:1413-6.

Russell PA, Phaoah PDP, De Foy K, et al. Frequent loss of BRCA1 mRNA and protein expression in sporadic ovarian cancers. *Int J Cancer* 2000;87:317-21.

Rökman A, Baffoe-Bonnie AB, Gillanders E, et al. Hereditary prostate cancer in Finland: fine-mapping validates 3p26 as a major predisposition locus. *Hum Genet* 2005;116:43-50.

Sajantila A, Salem AH, Savolainen P, Bauer K, Gierig C, Pääbö S. Paternal and maternal DNA lineages reveal a bottleneck in the founding of the Finnish population. *Proc Natl Acad Sci USA* 1996;93:12035-9.

Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci* 1977;74:5463-7.

Sarantaus L, Huusko P, Eerola H, et al. Multiple founder effects and geographical clustering of BRCA1 and BRCA2 families in Finland. *Eur J Hum Genet* 2000;8:757-63.

Sarantaus L, Auranen A, Nevanlinna H. BRCA1 and BRCA2 mutations among Finnish ovarian carcinoma families. *Int J Oncol* 2001a;18:831-5.

Sarantaus L, Vahteristo P, Bloom E, et al. BRCA1 and BRCA2 mutations among 233 unselected Finnish ovarian carcinoma patients. *Eur J Hum Genet* 2001b;9:424-30.

Sauer MK, Andrulis IL. Identification and characterization of missense alterations in the BRCA1 associated RING domain (BARD1) gene in breast and ovarian cancer. *J Med Genet* 2005;42:633-8.

Saurin AJ, Borden KLB, Boddy MN, Freemont PS. Does this have a familiar RING? *TIBS* 1996;21:208-14.

Savitsky K, Bar-Shira A, Gilad S, et al. A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* 1995;268:1749-53.

Schaid DJ, Rowland CM, Tines DE, Jacobson RM, Poland GA. Score tests for association of traits with haplotypes when linkage phase is ambiguous. *Am J Hum Genet* 2002;70:425-34. 2002.

Schechter AL, Stern DF, Vaidyanathan L, et al. The neu oncogene: an erb-B-related gene encoding a 185-000-Mr tumour antigen. *Nature* 1984;312:523-16.

Schernhammer ES, Hankinson SE, Hunter DJ, Blouin MJ, Pollak MN. Polymorphic variation at the -202 locus in IGFBP3: influence on serum levels of insulin-like growth factors, interaction with plasma retinol and vitamin D and breast cancer risk. *Int J Cancer* 2003;107:60-4.

Schleutker J, Baffoe-Bonnie AB, Gillanders E, et al. Genome-wide scan for linkage in Finnish hereditary prostate cancer (HPC) families identifies novel susceptibility loci at 11q14 and 3p25-26. *Prostate* 2003;57:280-9.

- Schmidt MK, Tollenaar RAEM, de Kemp SR, et al. Breast cancer survival and tumor characteristics in premenopausal women carrying the CHEK2\*1100delC germline mutation. *J Clin Oncol* 2007;25:64-9.
- Schutte M, da Costa LT, Hahn SA, et al. Identification by representational difference analysis of a homozygous deletion in pancreatic carcinoma that lies within the BRCA2 region. *Proc Nat Acad Sci USA* 1995;92:5950-4.
- Schutte M, Seal S, Barfoot R, et al. Variants in CHEK2 other than 1100delC do not make a major contribution to breast cancer susceptibility. *Am J Hum Genet* 2003;72:1023-8.
- Scott CL, Jenkins MA, Southey MC, et al. Average age-specific cumulative risk of breast cancer according to type and site of germline mutations in BRCA1 and BRCA2 estimated from multiple-case breast cancer families attending Australian family cancer clinics. *Hum Genet* 2003;112:542-51.
- Scully R, Chen J, Plug A, et al. Association of BRCA1 with Rad51 in mitotic and meiotic cells. *Cell* 1997a;88:265-75.
- Scully R, Chen J, Ochs RL, et al. Dynamic changes of BRCA1 subnuclear location and phosphorylation state are initiated by DNA damage. *Cell* 1997b;90:425-35.
- Scully R, Anderson SF, Chao DM, et al. BRCA1 is a component of the RNA polymerase II holoenzyme. *PNAS* 1997c;94:5605-10.
- Scully R, Livingston DM. In search of the tumour-suppressor functions of BRCA1 and BRCA2. *Nature* 2000;408:429-32.
- Seal S, Barfoot R, Jayatilake H, et al. Evaluation of Fanconi anemia genes in familial breast cancer predisposition. *Cancer Res* 2003;63:8596-9.
- Seal S, Thompson D, Renwick A, et al. Truncating mutations in the Fanconi anemia J gene BRIP1 are low-penetrance breast cancer susceptibility alleles. *Nat Genet* 2006;38:1239-41.
- Serrano M, Hannon GJ, Beach D. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* 1993;366:704-7.
- Seth P, Lunetta KL, Bell DW, et al. Phenol sulfotransferases: Hormonal regulation, polymorphism, and age of onset of breast cancer. *Cancer Res* 2000;60:6859-63.
- Shaag A, Walsh T, Renbaum P, et al. Functional and genomic approaches reveal an ancient allele associated with breast cancer in the Ashkenazi Jewish population. *Hum Mol Genet* 2005;14:555-63.
- Shannon C, Smith IE. Breast cancer in adolescents and young women. *Eur J Cancer* 2003;39:2632-42.
- Sharan Sk, Morimatsu M, Albrecht U, et al. Embryonic lethality and radiation hypersensitivity mediated by Rad51 in mice lacking Brca2. *Nature* 1997;386:804-10.

Sheffield VC, Beck JS, Kwitek AE, Sandstrom DW, Stone EM. The sensitivity of single-strand conformation polymorphism analysis for the detection of single base substitutions. *Genomics* 1993;16:325-32.

Sheffield VC, Weber JL, Buetow KH, et al. A collection of tri- and tetranucleotide repeat markers used to generate high quality, high resolution human genome-wide linkage maps. *Hum Mol Genet* 1995;4:413-9.

Sherr CJ. The INK4a/ARF network in tumour suppression. *Nat Rev Cancer* 2001;2:731-7.

Shieh S-Y, Ahn J, Tamai K, Taya Y, Prives C. The human homologs of checkpoint kinases Chk1 and Cds1 (Chk2) phosphorylate p53 at multiple DNA damage-inducible sites. *Genes Dev* 2000;14:289-300.

Shindo M, Nakano H, Kuroyanagi H, et al. cDNA cloning, expression, subcellular localization, and chromosomal assignment of mammalian aurora homologues, aurora-related kinase (ARK) 1 and 2. *Biochem Biophys Res Comm* 1998;244:285-92.

Sidransky D, Tokino T, Helzlsouer K, et al. Inherited p53 gene mutations in breast cancer. *Cancer Res* 1992;52:2984-6.

Sigurdson AJ, Hauptmann M, Chatterjee N, et al. Kin-cohort estimates for familial breast cancer risk in relation to variants in DNA base excision repair, BRCA1 interacting and growth factor genes. *BMC Cancer* 2004;4:9.

Sillanpää P, Hirvonen A, Kataja V, et al. Vitamin D receptor gene polymorphism as an important modifier of positive family history related breast cancer risk. *Pharmacogenetics* 2004;14:239-45.

Sillanpää P, Hirvonen A, Kataja V, et al. NAT2 slow acetylator genotype as an important modifier of breast cancer risk. *Int J Cancer* 2005a;114:579-84.

Sillanpää P, Kataja V, Eskelinen M, et al. Sulfotransferase 1A1 genotype as a potential modifier of breast cancer risk among premenopausal women. *Pharmacogenet Genomics* 2005b;15:749-52.

Sillanpää P, Heikinheimo L, Kataja V, et al. CYP1A1 and CYP1B1 genetic polymorphisms, smoking and breast cancer risk in a Finnish Caucasian population. *Breast Cancer Res Treat* 2007;104:287-97. Epub 2006, Oct 25.

Simard J, Tonin P, Durocher F, et al. Common origins of BRCA1 mutations in Canadian breast and ovarian cancer families. *Nat Genet* 1994;8:392-8.

Sjalander A, Birgander R, Hallmans G, et al. p53 polymorphisms and haplotypes in breast cancer. *Carcinogenesis* 1996;17:1313-6.

Sjöblom T, Jones S, Wood LD, et al. The consensus coding sequences of human breast and colorectal cancers. *Science* 2006;314:268-74.

Slatkin M, Excoffier L. Testing for linkage disequilibrium in genotypic data using the Expectation-Maximization algorithm. *Heredity* 1996;76:377-83.

Smith SA, Easton DF, Evans DG, Ponder BA. Allele losses in the region 17q12-21 in familial breast and ovarian cancer involve the wild-type chromosome. *Nat Genet* 1992;2:128-31.

Smith-Warner SA, Spiegelman D, Yaun SS, et al. Alcohol and breast cancer in women: a pooled analysis of cohort studies. *JAMA* 1998;279:535-40.

Sodha N, Wilson C, Bullock SL, Houlston RS, Eeles RA. Analysis of familial male breast cancer for germline mutations in CHEK2. *Cancer Lett* 2004;215:187-9.

Soininen AM Pohjois-Savon asuttaminen keski- ja uuden ajan vaihteessa. In: Historiallisia tutkimuksia LVIII. Suomen Historiallinen Seura, Helsinki, 1981.

Song H, Ramus SJ, Kruger Kjaer S, et al. Tagging single nucleotide polymorphisms in the BRIP1 gene and susceptibility to breast and ovarian cancer. *PLoS ONE* 2007;2:e268.

Sorlie T, Tibshirani R, Parker J, et al. Repeated observation of breast tumor subtypes in independent gene expression data sets. *Proc Natl Acad Sci USA* 2003;100:8418-23.

Soussi T, May P. Structural aspects of the p53 protein in relation to gene evolution: A second look. *J Mol Biol* 1996;260:623-37.

Southey MC, Batten LE, McCredie MRE, et al. Estrogen receptor polymorphism at codon 325 and risk of breast cancer in women before age forty. *J Natl Cancer Inst* 1998;90:532-6.

Spain BH, Larson CJ, Shihabuddin LS, Gage FH, Verma IM. Truncated BRCA2 is cytoplasmic: Implications for cancer-linked mutations. *Proc Natl Acad Sci USA* 1999;96:13920-5.

Spinardi L, Mazars R, Theillet C. Protocols for an improved detection of point mutations by SSCP. *Nucl Acids Res* 1991;19:4009.

Spurdle AB, Hopper JD, Chen X, et al. The BRCA2 372 HH genotype is associated with risk of breast cancer in Australian women under age 60 years. *Cancer Epidemiol Biomarkers Prev* 2002;11:413-6.

Spurdle AB, Antoniou AC, Keleman L, et al. The AIB1 polyglutamine repeat does not modify breast cancer risk in BRCA1 and BRCA2 mutation carriers. *Cancer Epidemiol Biomarkers Prev* 2006;15:76-9.

Stacey SN, Sulem P, Johannsson OT, et al. The BARD1 Cys557Ser variant and breast cancer risk in Iceland. *PLOS Med* 2006;3:e217.

Stacey SN, Manolescu A, Sulem P, et al. Common variants on chromosome 2q35 and 16q12 confer susceptibility to estrogen receptor-positive breast cancer. *Nat Genet* 2007; published online May 27, doi:10.1038/ng2064

Starink TM, van der Veen JP, Arwert F, et al. The Cowden syndrome: a clinical and genetic study in 21 patients. *Clin Genet* 1986;29:222-33.

Steffen J, Nowakowska D, Niwinska A, et al. Germline mutations 657del5 of the NBS1 gene contribute significantly to the incidence of breast cancer in Central Poland. *Int J Cancer* 2006;119:472-5.

Stevens C, Smith L, La Thangue NB. Chk2 activates E2F-1 in response to DNA damage. *Nature Cell Biol* 2003;5:401-9.

Stoppa-Lyonnet D, Ansquer Y, Dreyfus H, et al. Familial invasive breast cancers: Worse outcome related to BRCA1 mutations. *J Clin Oncol* 2000;18:4053-9.

Strachan T, Read AP. *Human Molecular Genetics*. 3rd ed. New York: Garland Science, 2003.

Stratton MR, Ford D, Neuhausen S, et al. Familial male breast cancer is not linked to the BRCA1 locus on chromosome 17q. *Nat Genet* 1994;7:103-7.

Stredrick DL, Garcia-Closas M, Pineda MA, et al. The ATM missense mutation p.Ser49Cys (c.146C>G) and the risk of breast cancer. *Hum Mut* 2006;27:538-44.

Struwing JP, Hartge P, Wacholder S, et al. The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews. *N Engl J Med* 1997;336:1401-8.

Sun T, Miao X, Wang J, et al. Functional Phe31Ile polymorphism in Aurora A and risk of breast carcinoma. *Carcinogenesis* 2004;25:222-30.

Sutter TR, Tang YM, Haeyls CL, et al. Complete cDNA sequence of a human dioxin-inducible mRNA identifies a new gene subfamily of cytochrome P450 that maps to chromosome 2. *J Biol Chem* 1994;269:13092-9.

Swensen J, Hoffman M, Skolnick MH, Neuhausen SL. Identification of a 14 kb deletion involving the promoter region of BRCA1 in a breast cancer family. *Hum Mol Genet* 1997;6:1513-7.

Swift M, Reitnauer PJ, Morrell D, Chase CL. Breast and other cancers in families with ataxia-telangiectasia. *N Engl J Med* 1987;316:1289-94.

Swift M, Morrell D, Massey RB, Chase CL. Incidence of cancer in 161 families affected by ataxia-telangiectasia. *N Engl J Med* 1991;325:1831-6.

Syrjäkoski K, Vahteristo P, Eerola H, et al. Population-based study of BRCA1 and BRCA2 mutations in 1035 unselected Finnish breast cancer patients. *JNCI* 2000;92:1529-31.

Syrjäkoski K, Kuukasjärvi T, Waltering K, et al. BRCA2 mutations in 154 Finnish male breast cancer patients. *Neoplasia* 2004a;6:541-5.

Syrjäkoski K, Kuukasjärvi T, Auvinen A, Kallioniemi O-P. CHEK2 is not a risk factor for male breast cancer population. *Int J Cancer* 2004b;108:475-6.

Takahashi H, Behbakt K, McGovern PE, et al. Mutation analysis of the BRCA1 gene in ovarian cancer. *Cancer Res* 1995;55:2998-3002.



Tamimi RM, Hankinson SE, Ding S, et al. The HRAS1 variable number of tandem repeats and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12:1528-30.

Tamimi RM, Hankinson SE, Spiegelman D, Kraft P, Colditz GA, Hunter DJ. Common ataxia telangiectasia mutated haplotypes and risk of breast cancer: a nested case-control study. *Breast Cancer Res*;2004a;6:R416-R422.

Tamimi RM, Hankinson SE, Spiegelman D, Colditz G, Hunter DJ. Manganese superoxide dismutase polymorphism, plasma antioxidants, cigarette smoking, and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2004b;13:989-96.

Tang YM, Wo YYP, Stewart J, et al. Isolation and characterization of the human cytochrome P450 CYP1B1 gene. *J Biol Chem* 1996;271:28324-30.

Taniguchi T, D'Andrea AD. Molecular pathogenesis of Fanconi anemia: recent progress. *Blood* 2006;107:4223-33.

Tavtigian SV, Simard J, Rommens J, et al. The complete BRCA2 gene and mutations in chromosome 13q-linked kindreds. *Nat Genet* 1996;12:333-7.

Taylor J, Lymboura M, Pace PE, et al. An important role for BRCA1 in breast cancer progression is indicated by its loss in a large proportion of non-familial breast cancers. *Int J Cancer* 1998;79:334-42.

Teng DHF, Bogden R, Mitchell J, et al. Low incidence of BRCA2 mutations in breast carcinoma and other cancers. *Nat Genet* 1996;13:241-4.

Teraoka SN, Malone KE, Doody DR, et al. Increased frequency of ATM mutations in breast carcinoma patients with early onset disease and positive family history. *Cancer* 2001;92:479-87.

Terry MB, Gammon MD, Zhang FF, et al. Polymorphisms in the DNA repair gene XPD, polycyclic aromatic hydrocarbon-DNA adducts, cigarette smoking, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2004;13:2053-8.

Terry MB, Gammon MD, Zhang FF, et al. ADH3 genotype, alcohol intake and breast cancer risk. *Carcinogenesis* 2006;27:840-7.

Terwilliger JD, Ott J. *Handbook of Human Genetic Linkage*. Johns Hopkins University Press, Baltimore, 1994.

Thai TH, Du F, Tsou Tsan J, et al. Mutations in the BRCA1-associated RING domain (BARD1) gene in primary breast, ovarian and uterine cancers. *Hum Mol Genet* 1998;7:195-202.

Thakur S, Zhang HB, Peng Y, et al. Localization of BRCA1 and a splice variant identifies the nuclear localization signal. *Mol Cell Biol* 1997;17:444-52.

Thompson ME, Jensen RA, Obermiller PS, Page DL, Holt JT. Decreased expression of BRCA1 accelerates growth and is often present during sporadic breast cancer progression. *Nature Genet* 1995;9:444-50.

- Thompson PA, Shields PG, Freudenheim JL, et al. Genetic polymorphisms in catechol-O-methyltransferase, menopausal status, and breast cancer risk. *Cancer Res* 1998;58:2107-10.
- Thompson PA, Ambrosone C. Molecular epidemiology of genetic polymorphisms in estrogen metabolizing enzymes in human breast cancer. *J Natl Cancer Inst Monogr* 2000;27:125-34.
- Thompson D, Easton D. Variation in cancer risks, by mutation position, in BRCA2 mutation carriers. *Am J Hum Genet* 2001;68:410-9.
- Thompson D, Easton DF, the Breast Cancer Linkage Consortium. Cancer incidence in BRCA1 mutation carriers. *J Natl Cancer Inst* 2002a;94:1358-65.
- Thompson D, Easton D, the Breast Cancer Linkage Consortium. Variation in BRCA1 cancer risks by mutation position. *Cancer Epidemiol Biomarkers Prev* 2002b;11:329-36.
- Thompson D, Easton D. The genetic epidemiology of breast cancer genes. *J Mammary Gland Biol* 2004;9:221-36.
- Thompson D, Duedal S, Kirner J, et al. Cancer risks and mortality in heterozygous ATM carriers. *J Natl Cancer Inst* 2005;97:813-22.
- Thompson D, Seal S, Schutte M, et al. A multicenter study of cancer incidence in CHEK2 1100delC mutation carriers. *Cancer Epidemiol Biomarkers Prev* 2006;15:2542-5.
- Thorlacius S, Olafsdottir G, Tryggvadottir L, et al. A single BRCA2 mutation in male and female breast cancer families from Iceland with varied cancer phenotypes. *Nat Genet* 1996;13:117-9.
- Thorlacius S, Sigurdsson S, Bjarnadottir H, et al. Study of a single BRCA2 mutation with high carrier frequency in a small population. *Am J Hum Genet* 1997;60:1079-84.
- Thorlacius S, Struwing J, Hartge P, et al. Population-based study of risk of breast cancer in carriers of BRCA2 mutation. *Lancet* 1998;352:1337-9.
- Tirkkonen M, Johannsson O, Agnarsson BA, et al. Distinct somatic genetic changes associated with tumor progression in carriers of BRCA1 and BRCA2 germ-line mutations. *Cancer Res* 1997;57:1222-7.
- Toda T, Miyake M, Kobayashi K, et al. Linkage-disequilibrium mapping narrows the Fukuyama-type congenital muscular dystrophy (FCMD) candidate region to <100 kb. *Am J Hum Genet* 1996;59:1313-20.
- Tominaga K, Morisaki H, Kaneko Y, et al. Role of human Cds1 (Chk2) kinase in DNA damage checkpoint and its regulation by p53. *J Biol Chem* 1999;274:31463-7.
- Tomlinson IP, Houlston RS. Peutz-Jeghers syndrome. *J Med Genet* 1997;34:1007-11.
- Tommiska J, Eerola H, Heinonen M, et al. Breast cancer patients with p53 Pro72 homozygous genotype have a poorer survival. *Clin Cancer Res* 2005;11:5098-103.

Tretli S. Height and weight in relation to breast cancer morbidity and mortality. A prospective study of 570 000 women in Norway. *Int J Cancer* 1989;44:23-30.

Trichopoulos D, MacMahon B, Cole P. Menopause and breast cancer risk. *J Natl Cancer Inst* 1972;48:605-13.

Tsou HC, Teng DHF, Ping XL, et al. The role of MMAC1 mutations in early-onset breast cancer: causative in association with Cowden syndrome and excluded in BRCA1-negative cases. *Am J Hum Genet* 1997;61:1036-43.

Tuhkanen H, Anttila M, Kosma VM, et al. Genetic alterations in the peritumoral stromal cells of malignant and borderline epithelial ovarian tumors as indicated by allelic imbalance on chromosome 3p. *Int J Cancer* 2004;109:247-52.

Tutt A, Bertwistle D, Valentine J, et al. Mutation in Brca2 stimulates error-prone homology-directed repair of DNA double-strand breaks occurring between repeated sequences. *EMBO J* 2001;20:4704-16.

Uitterlinden AG, Fang Y, van Meurs JBJ, Pols HAP, van Leeuwen JPTM. Genetics and biology of vitamin D receptor polymorphisms. *Gene* 2004;338:143-56.

Urist M, Tanaka T, Poyurovsky MV, Prives C. p73 induction after DNA damage is regulated by checkpoint kinases Chk1 and Chk2. *Genes Devel* 2004;18:3041-54.

Vahteristo P, Eerola H, Tamminen A, Blomqvist C, Nevanlinna H. A probability model for predicting BRCA1 and BRCA2 mutations in breast and breast-ovarian cancer families. *Br J Cancer* 2001a;84:704-8.

Vahteristo P, Tamminen A, Karvinen P, et al. p53, CHK2 and CHK1 genes in Finnish families with Li-Fraumeni syndrome: further evidence of CHK2 in inherited cancer predisposition. *Cancer Res* 2001b;61:5718-22.

Vahteristo P, Bartkova J, Eerola H, et al. A CHEK2 genetic variant contributing to a substantial fraction of familial breast cancer. *Am J Hum Genet* 2002;71:432-8.

van der Lijft R, Khan PM, Vasen H, et al. Rapid detection of translation-terminating mutations at the adenomatous polyposis coli (APC) gene by direct protein truncation test. *Genomics* 1994;20:1-4.

van Gent DC, Hoeijmakers JHJ, Kanaar R. Chromosomal stability and the DNA double-stranded break connection. *Nat Rev Genet* 2001;2:196-206.

Van Rensburg CE, Van Staden AM, Anderson R, Van Rensburg EJ. Hypochlorous acid potentiates hydrogen peroxide-mediated DNA-strand breaks in human mononuclear leucocytes. *Mutat Res* 1992;265:255-61.

Vandenplas S, Grobler-Rabie A, Brebner K, et al. Blot hybridization of genomic DNA. *J Med Genet* 1984;21:164-72.

van't Veer LJ, Dai H, van der Vijver, et al. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 2002;415:530-56.

Varilo T, Savukoski M, Norio R, Santavuori P, Peltonen L, Järvelä I. The age of human mutation: Genealogical and linkage disequilibrium analysis of the CLN5 mutation in the Finnish population. *Am J Hum Genet* 1996;58:506-12.

Varilo T, Paunio T, Parker A, et al. The interval of linkage disequilibrium (LD) detected with microsatellite and SNP markers in chromosomes of Finnish populations with different histories. *Hum Mol Genet* 2003;12:51-9.

Varilo T, Peltonen L. Isolates and their potential use in complex gene mapping efforts. *Curr Opin Genet Devel* 2004;14:316-23.

Varley JM, McGown G, Thorncroft M, et al. Germ-line mutations of TP53 in Li-Fraumeni families: an extended study of 39 families. *Cancer Res* 1997;57:3245-52.

Vatsis KP, Martell KJ, Weber WW. Diverse point mutations in the human gene for polymorphic N-acetyltransferase. *Proc Natl Acad Sci USA* 1991;88:6333-7.

Vatten LJ, Kvinnsland S. Body height and risk of breast cancer. A prospective study of 23 831 Norwegian women. *Br J Cancer* 1990;61:881-5.

Vehmanen P, Friedman LS, Eerola H, et al. Low proportion of BRCA1 and BRCA2 mutations in Finnish breast cancer families: evidence for additional susceptibility genes. *Hum Mol Genet* 1997a;6:2309-15.

Vehmanen P, Friedman LS, Eerola H, et al. A low proportion of BRCA2 mutations in Finnish breast cancer families. *Am J Hum Genet* 1997b;60:1050-8.

Velasco G, Cal S, Quesada V, Sanchez LM, Lopes-Otin C. Matriptase-2, a membrane-bound mosaic serine proteinase predominantly expressed in human liver and showing degrading activity against extracellular matrix proteins. *J Biol Chem* 2002; 277:37637-46.

Venkitaraman AR. Cancer susceptibility and the functions of BRCA1 and BRCA2. *Cell* 2002;108:171-82.

Verhoog LC, Brekelmans CTM, Seynaeve C, et al. Survival and tumour characteristics of breast-cancer patients with germline mutations of BRCA1. *Lancet* 1998;351:316-21.

Verhoog LC, Berns EMJJ, Brekelmans CTM, Seynaeve C, Meijers-Heijboer EJ, Klijn JGM. Prognostic significance of germline BRCA2 mutations in hereditary breast cancer patients. *J Clin Oncol* 2000;18:119s-124s.

Visapää I, Salonen R, Varilo T, Paavola P, Peltonen L. Assignment of the locus for Hydrolethalus syndrome to a highly restricted region on 11q23-25. *Am J Hum Genet* 1999;65:1086-95.

Vogelstein B, Lane D, Levine AJ. Surfing the p53 network. *Nature* 2000;408:307-10.

Vogl FD, Taioli E, Maugard C, et al. Glutathione S-transferases M1, T1, and P1 and breast cancer: a pooled analysis. *Cancer Epidemiol Biomarkers Prev* 2004;13:1473-9.

Wagner K, Hemminki K, Grzybowska E, et al. Polymorphisms in genes involved in GH1 release and their association with breast cancer risk. *Carcinogenesis* 2006a;27:1867-75.

Walker KK, Levine AJ. Identification of a novel p53 functional domain that is necessary for efficient growth suppression. *Proc Natl Acad Sci USA* 1996;93:15335-40.

Walsh T, Casadei S, Coats KH, et al. Spectrum of mutations in BRCA1, BRCA2, CHEK2, and TP53 infamilies at high risk of breast cancer. *JAMA* 2006;295:1379-88.

Walter P, Green S, Greene G, et al. Cloning of the human estrogen receptor cDNA. *Proc Natl Acad Sci USA* 1985;82:7889-93.

Walworth NC. Cell-cycle checkpoint kinases: checking in on the cell cycle. *Curr Opin Cell Biol* 2000;12:697-704.

Wang Y, Cortez D, Yazdi P, Neff N, Elledge SJ, Qin J. BASC, a super complex of BRCA1-associated proteins involved in the recognition and repair of aberrant DNA structures. *Genes Dev* 2000;14:927-39.

Wang WW, Spurdle AB, Kolachana P, et al. A single nucleotide polymorphism in the 5' untranslated region of RAD51 and risk of cancer among BRCA1/2 mutation carriers. *Cancer Epidemiol Biomarkers Prev* 2001a;10:955-60.

Wang Q, Zhang H, Guerrette S, et al. Adenosine nucleotide modulates the physical interaction between hMSH2 and BRCA1. *Oncogene* 2001b;20:4640-9.

Wang SC, Shao R, Pao AY, Zhang S, Hung MC, Su LK. Inhibition of cancer cell growth by BRCA2. *Cancer Res* 2002;62:1311-4.

Wang J, Costantino JP, Tan-Chiu E, Wickerham DL, Paik S, Wolmark N. Lower-category benign breast disease and the risk of invasive breast cancer. *J Natl Cancer Inst* 2004;96:616-20.

Wang -Gohrke S, Becher H, Krienberg R, Runnebaum IB, Chang-Claude J. Intron 3 16 bp duplication polymorphism of p53 is associated with an increased risk for breast cancer by the age of 50 years. *Pharmacogenetics* 2002;12:269-72.

Ware MD, DeSilva D, Sinilnikova OM, Stoppa-Lyonnet D, Tavtigian SV, Mazoyer S. Does nonsense-mediated mRNA decay explain the ovarian cancer cluster region of the BRCA2 gene? *Oncogene* 2006, 25:323-8.

Weber BHF, Brohm M, Stec I, Backe J, Caffier H. A somatic truncating mutation in BRCA2 in a sporadic breast tumour. *Am J Hum Genet* 1996;59:962-4.

Weber BL, Nathanson KL. Low penetrance genes associated with increased risk of breast cancer. *Eur J Cancer* 2000;36:1193-9.

- Wei JH, Chou YF, Ou YH, et al. TTK/hMps1 participates in the regulation of DNA damage checkpoint response by phosphorylating CHK2 on threonine 68. *J Biol Chem* 2005;280:7748-57.
- Weinert T. Yeast checkpoint controls and relevance to cancer. *Cancer Surv* 1997;29:109-32.
- Weinert T. DNA damage and checkpoint pathways: molecular anatomy and interaction with repair. *Cell* 1998;94:555-8.
- Weitzel JN, Ding S, Larson GP, et al. The hHRAS1 minisatellite locus and risk of ovarian cancer. *Cancer Res* 2000;60:259-61.
- Welsh PL, Owens KN, King MC. Insights into the function of BRCA1 and BRCA2. *Trends Genet* 2000;16:69-74.
- Welsh P, King MC. BRCA1 and BRCA2 and the genetics of breast and ovarian cancer. *Hum Mol Genet* 2001;10:705-13.
- Wen W, Gao YT, Shu XO, et al. Insulin-like growth factor-I gene polymorphisms and breast cancer risk in Chinese women. *Int J Cancer* 2005;113:307-11.
- Wenham RM, Schildkraut JM, McLean K, et al. Polymorphisms in BRCA1 and BRCA2 and risk of epithelial ovarian cancer. *Clin Cancer Res* 2003;9:4396-403.
- Whelan AJ, Bartch D, Goodfellow PJ. Brief report: A familial syndrome of pancreatic cancer and melanoma with a mutation in the CDKN2 tumor-suppressor gene. *N Engl J Med* 1995;33:975-7.
- White MB, Carvalho M, Derse D, O'Brien SJ, Dean M. Detecting single base substitutions as heteroduplex polymorphisms. *Genomics* 1992;12:301-6.
- Whitehouse C, Solomon E. Current status of the molecular characterization of the ovarian cancer antigen CA125 and implications for its use in clinical screening. *Gynecol Oncol* 2003;88:S152-S157.
- Whitfield GK, Remus LS, Jurutka PW, et al. Functional relevant polymorphisms in the human nuclear vitamin D receptor gene. *Mol Cell Endocrinol* 2001;177:145-59.
- WHO. Pathology and Genetics: Tumours of the Breast and Female Genital Organs. Lyon: IARC Press, 2003.
- Wilkening S, Burwinkel B, Grzybowska E, et al. Polyglutamine repeat length in the NCOA3 does not affect risk in familial breast cancer. *Cancer Epidemiol Biomarkers Prev* 2005;14:291-2.
- Williams JA, Stone EM, Millar BC, Hewer A, Phillips DH. Pathways of heterocyclic amine activation in the breast: DNA adducts of 2-amino-3-methylimidazo[4,5-f]quinoline (IQ) formed by peroxidases and in human mammary epithelial cells and fibroblasts. *Mutagenesis* 2000;15:149-54.

- Wilson CA, Ramos L, Vilasenor MR, et al. Localization of human BRCA1 and its loss in high-grade, non-inherited breast carcinomas. *Nat Genet* 1999;21:236-40.
- Winterbourn CC. Superoxide as an intracellular radical risk. *Free Radic Biol Med* 1993;14:85-90.
- Wirtenberger M, Hemminki K, Försti A, et al. c-MYC Asn11Ser is associated with increased risk for familial breast cancer. *Int J Cancer* 2005;117:638-42.
- Wong AKC, Pero R, Ormonde PA, Tavtigian V, Bartel PL. RAD51 interacts with the evolutionarily conserved BRC motifs in the human breast cancer susceptibility gene *brca2*. *J Biol Chem* 1997;272:31941-4.
- Wooster R, Neuhausen SL, Mangion J, et al. Localization of a breast cancer susceptibility gene, BRCA2, to chromosome 13q12-13. *Science* 1994;265:2088-90.
- Wooster R, Bignell G, Lancaster J, et al. Identification of the breast cancer susceptibility gene BRCA2. *Nature* 1995;378:789-92.
- Wu LC, Wang ZW, Tsan JT, et al. Identification of a RING protein that can interact in vivo with the BRCA1 gene product. *Nat Genet* 1996;14:430-40.
- Wu X, Webster SR, Chen J. Characterization of tumor-associated Chk2 mutations. *J Biol Chem* 2001;276:2971-4.
- Xia F, Taghian DG, DeFrank JS, et al. Deficiency of human BRCA2 leads to impaired homologous recombination but maintains normal nonhomologous end joining. *Proc Natl Acad Sci USA* 2001;98:8644-9.
- Xia B, Sheng Q, Nakanishi K, et al. Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2. *Mol Cell* 2006;22:719-29.
- Xia B, Dorsman JC, Ameziane N, et al. Fanconi anemia is associated with a defect in the BRCA2 partner PALB2. *Nat Genet* 2007;39:159-61.
- Xie D, Shu XO, Deng Z, et al. population-based, case-control study of HER2 genetic polymorphism and breast cancer risk. *J Natl Cancer Inst* 2000;92:412-7.
- Xu X, Weaver Z, Linke SP, et al. Centrosome amplification and a defective G2-M cell cycle checkpoint induce genetic instability in BRCA1 exon 11 isoform-deficient cells. *Molecular Cell* 1999;3:389-95.
- Xu B, Kim ST, Kastan MB. Involvement of Brca1 in S-phase and G2-phase checkpoints after ionizing irradiation. *Mol Cell Biol* 2001;21:3445-50.
- Yang H, Jeffrey PD, Miller J, et al. BRCA2 function in DNA binding and recombination from a BRCA2-DSS1-ssDNA structure. *Science* 2002a;297:1837-48.
- Yang S, Kuo C, Bisi JE, Kim MK. PML-dependent apoptosis after DNA damage is regulated by the checkpoint kinase hCds1/Chk2. *Nature Cell Biol* 2002b;4:865-70.

Yang Q, Yoshimura G, Mori I, Sakurai T and Kakudo K. Chromosome 3p and breast cancer. *J Hum Genet* 2002c;47:453-9.

Yang-Feng TL, Schechter AL, Weinberg RA, Francke U. Oncogene from rat neuro/glioblastomas (human gene symbol NGL) is located on the proximal long arm of human chromosome 17 and EGFR is confirmed at 7p13-q11.2. *Cytogenet Cell Genet* 1985;40:784.

Yang-Feng TL, Han H, Chen KC, et al. Allelic loss in ovarian cancer. *Int J Cancer* 1993;54:546-51.

Yarden RI, Brody LC. BRCA1 interacts with components of the histone deacetylase complex. *PNAS* 1999;96:4983-8.

Yarden Y, Sliwkowski MX. Untangling the erbB signalling network. *Nat Rev Mol Cell Biol* 2001;2:127-37.

Yoshiji H, Gomez DE, Shibuya M, Thorgeirsson. Expression of vascular endothelial growth factor, its receptor, and other angiogenic factors in human breast cancer. *Cancer Res* 1996;56:2013-6.

Yu H, Rohan T. Role of insulin-like growth factor family in cancer development and progression. *J Natl Cancer Inst* 2000;92:1472-89.

Zavattari P, Deidda E, Whalen M, et al. Major factors influencing linkage disequilibrium by analysis of different chromosome regions in distinct populations: demography, chromosome recombination frequency and selection. *Hum Mol Genet* 2000;9:2947-57.

Zhang Y, Newcomb PA, Rgan KM, et al. Genetic polymorphisms in base-excision repair pathway genes and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2006;15:353-8.

Zhao JH, Sham PC. Faster haplotype frequency estimation using unrelated subjects. *Hum Hered* 2002;53:36-41.

Zheng W, Gustafson DR, Sinha R, et al. Well-done meat intake and the risk of breast cancer. *J Natl Cancer Inst* 1998;90:1724-9.

Zheng W, Deitz AC, Campbell DR, et al. N-acetyltransferase 1 genetic polymorphism, cigarette smoking, well-done meat intake, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 1999;8:233-9.

Zheng W, Xie D, Cerhan JR, Sellers TA, Wen W, Folsom AR. Sulfotransferase 1A1 polymorphism, endogenous estrogen exposure, well-done meat intake, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2001;10:89-94.

Zheng SL, Zheng W, Chang BI, et al. Joint effect of estrogen receptor beta sequence variants and endogenous estrogen exposure on breast cancer risk in Chinese women. *Cancer Res* 2003;63:7624-9.

Zhong Q, Chen CF, Li S, et al. Association of BRCA1 with the hRad50-hMre11-p95 complex and the DNA damage response. *Science* 1999;285:747-50.



Zhu BT, Conney AH. Functional role of estrogen metabolism in target cells: review and perspectives. *Carcinogenesis* 1998;19:1-27.

Ziegler RG, Hoover RN, Pike MC, et al. Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst* 1993;85:1819-27.



## **9. ORIGINAL PUBLICATIONS**

## Kuopio University Publications D. Medical Sciences

**D 400. Vasara, Anna.** Autologous chondrocyte transplantation: Properties of the repair tissue in humans and in animal models.  
2007. 92 p. Acad. Diss.

**D 401. Andrulionyte, Laura.** Transcription factors as candidate genes for type 2 diabetes: studies on peroxisome proliferator-activated receptors, hepatic nuclear factor 4 $\alpha$  and PPAR $\gamma$  coactivator 1 $\alpha$ .  
2007. 112 p. Acad. Diss.

**D 402. Raatikainen, Kaisa.** Health behavioural and social risks in obstetrics: effect on pregnancy outcome.  
2007. 100 p. Acad. Diss.

**D 403. Kinnunen, Tuure.** The role of T cell recognition in the immune response against lipocalin allergens: prospects for immunotherapy.  
2007. 76 p. Acad. Diss.

**D 404. Gratz, Silvia.** Aflatoxin binding by probiotics : experimental studies on intestinal aflatoxin transport, metabolism and toxicity.  
2007. 85 p. Acad. Diss.

**D 405. Ming, Zhiyong.** Upper limb musculoskeletal disorders with special reference to sympathetic nerve functions and tactile sensation.  
2007. 91 p. Acad. Diss.

**D 406. Timonen, Leena.** Group-based exercise training in mobility impaired older women.  
2007. 91 p. Acad. Diss.

**D 407. Mättö, Mikko.** B cell receptor signaling in human B cells.  
2007. 74 p. Acad. Diss.

**D 408. Polychronaki, Nektaria.** Biomarkers of aflatoxin exposure and a dietary intervention: studies in infants and children from Egypt and Guinea and young adults from China.  
2007. 114 p. Acad. Diss.

**D 409. Mursu, Jaakko.** The role of polyphenols in cardiovascular diseases.  
2007. 88 p. Acad. Diss.

**D 410. Wlodkowic, Donald.** Selective targeting of apoptotic pathways in follicular lymphoma cells.  
2007. 97 p. Acad. Diss.

**D 411. Skommer, Joanna.** Novel approaches to induce apoptosis in human follicular lymphoma cell lines - preclinical assessment.  
2007. 80 p. Acad. Diss.

**D 412. Kempainen, Kaarina.** Early maternal sensitivity: continuity and related risk factors.  
2007. 80 p. Acad. Diss.

**D 413. Sahlman, Janne.** Chondrodysplasias Caused by Defects in the Col2a1 Gene.  
2007. 86 p. Acad. Diss.

**D 414. Pitkänen, Leena.** Retinal pigment epithelium as a barrier in drug permeation and as a target of non-viral gene delivery.  
2007. 75 p. Acad. Diss.

**D 415. Suhonen, Kirsi.** Prognostic Role of Cell Adhesion Factors and Angiogenesis in Epithelial Ovarian Cancer.  
2007. 123 p. Acad. Diss.